



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 11, 2024 – 08:40 PM EST

PDB ID : 2G6V  
Title : The crystal structure of ribD from Escherichia coli  
Authors : Stenmark, P.; Moche, M.; Gurmu, D.; Nordlund, P.; Structural Proteomics in Europe (SPINE)  
Deposited on : 2006-02-25  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	<div> <div>20%</div> <div>54%</div> <div>28%</div> <div>5%</div> <div>12%</div> </div>
1	B	402	<div> <div>33%</div> <div>59%</div> <div>25%</div> <div>6%</div> <div>10%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5530 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Riboflavin biosynthesis protein ribD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	355	Total	C	N	O	S	Se	0	0	0
			2709	1708	488	498	6	9			
1	B	363	Total	C	N	O	S	Se	0	0	0
			2756	1730	503	511	3	9			

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-26	MET	-	expression tag	UNP P25539
A	-25	ASP	-	expression tag	UNP P25539
A	-24	TYR	-	expression tag	UNP P25539
A	-23	LYS	-	expression tag	UNP P25539
A	-22	ASP	-	expression tag	UNP P25539
A	-21	ASP	-	expression tag	UNP P25539
A	-20	ASP	-	expression tag	UNP P25539
A	-19	ASP	-	expression tag	UNP P25539
A	-18	LYS	-	expression tag	UNP P25539
A	-17	GLY	-	expression tag	UNP P25539
A	-16	SER	-	expression tag	UNP P25539
A	-15	SER	-	expression tag	UNP P25539
A	-14	THR	-	expression tag	UNP P25539
A	-13	SER	-	expression tag	UNP P25539
A	-12	LEU	-	expression tag	UNP P25539
A	-11	TYR	-	expression tag	UNP P25539
A	-10	LYS	-	expression tag	UNP P25539
A	-9	LYS	-	expression tag	UNP P25539
A	-8	ALA	-	expression tag	UNP P25539
A	-7	GLY	-	expression tag	UNP P25539
A	-6	SER	-	expression tag	UNP P25539
A	-5	GLU	-	expression tag	UNP P25539
A	-4	THR	-	expression tag	UNP P25539
A	-3	LEU	-	expression tag	UNP P25539
A	-2	TYR	-	expression tag	UNP P25539

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ILE	-	expression tag	UNP P25539
A	0	GLN	-	expression tag	UNP P25539
A	1	GLY	-	expression tag	UNP P25539
A	7	MSE	MET	modified residue	UNP P25539
A	58	MSE	MET	modified residue	UNP P25539
A	100	MSE	MET	modified residue	UNP P25539
A	127	MSE	MET	modified residue	UNP P25539
A	128	MSE	MET	modified residue	UNP P25539
A	142	MSE	MET	modified residue	UNP P25539
A	163	MSE	MET	modified residue	UNP P25539
A	285	MSE	MET	modified residue	UNP P25539
A	286	MSE	MET	modified residue	UNP P25539
A	368	SER	-	expression tag	UNP P25539
A	369	THR	-	expression tag	UNP P25539
A	370	HIS	-	expression tag	UNP P25539
A	371	HIS	-	expression tag	UNP P25539
A	372	HIS	-	expression tag	UNP P25539
A	373	HIS	-	expression tag	UNP P25539
A	374	HIS	-	expression tag	UNP P25539
A	375	HIS	-	expression tag	UNP P25539
B	-26	MET	-	expression tag	UNP P25539
B	-25	ASP	-	expression tag	UNP P25539
B	-24	TYR	-	expression tag	UNP P25539
B	-23	LYS	-	expression tag	UNP P25539
B	-22	ASP	-	expression tag	UNP P25539
B	-21	ASP	-	expression tag	UNP P25539
B	-20	ASP	-	expression tag	UNP P25539
B	-19	ASP	-	expression tag	UNP P25539
B	-18	LYS	-	expression tag	UNP P25539
B	-17	GLY	-	expression tag	UNP P25539
B	-16	SER	-	expression tag	UNP P25539
B	-15	SER	-	expression tag	UNP P25539
B	-14	THR	-	expression tag	UNP P25539
B	-13	SER	-	expression tag	UNP P25539
B	-12	LEU	-	expression tag	UNP P25539
B	-11	TYR	-	expression tag	UNP P25539
B	-10	LYS	-	expression tag	UNP P25539
B	-9	LYS	-	expression tag	UNP P25539
B	-8	ALA	-	expression tag	UNP P25539
B	-7	GLY	-	expression tag	UNP P25539
B	-6	SER	-	expression tag	UNP P25539
B	-5	GLU	-	expression tag	UNP P25539

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	THR	-	expression tag	UNP P25539
B	-3	LEU	-	expression tag	UNP P25539
B	-2	TYR	-	expression tag	UNP P25539
B	-1	ILE	-	expression tag	UNP P25539
B	0	GLN	-	expression tag	UNP P25539
B	1	GLY	-	expression tag	UNP P25539
B	7	MSE	MET	modified residue	UNP P25539
B	58	MSE	MET	modified residue	UNP P25539
B	100	MSE	MET	modified residue	UNP P25539
B	127	MSE	MET	modified residue	UNP P25539
B	128	MSE	MET	modified residue	UNP P25539
B	142	MSE	MET	modified residue	UNP P25539
B	163	MSE	MET	modified residue	UNP P25539
B	285	MSE	MET	modified residue	UNP P25539
B	286	MSE	MET	modified residue	UNP P25539
B	368	SER	-	expression tag	UNP P25539
B	369	THR	-	expression tag	UNP P25539
B	370	HIS	-	expression tag	UNP P25539
B	371	HIS	-	expression tag	UNP P25539
B	372	HIS	-	expression tag	UNP P25539
B	373	HIS	-	expression tag	UNP P25539
B	374	HIS	-	expression tag	UNP P25539
B	375	HIS	-	expression tag	UNP P25539


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	42	Total O 42 42	0	0
2	B	23	Total O 23 23	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Riboflavin biosynthesis protein ribD



#### • Molecule 1: Riboflavin biosynthesis protein ribD





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	172.60Å 172.60Å 76.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.25 – 2.60 28.25 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.25-2.60) 99.9 (28.25-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.83 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.244 , 0.285 0.237 , 0.271	Depositor DCC
$R_{free}$ test set	2010 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	71.0	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 81.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	5530	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/2753	0.66	2/3723 (0.1%)
1	B	0.41	0/2800	0.62	0/3790
All	All	0.44	0/5553	0.64	2/7513 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	LEU	CA-CB-CG	5.26	127.40	115.30
1	A	67	THR	N-CA-C	5.20	125.04	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2709	0	2698	111	0
1	B	2756	0	2735	97	0
2	A	42	0	0	6	0
2	B	23	0	0	6	0
All	All	5530	0	5433	202	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 202 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:213:GLU:HA	2:B:395:HOH:O	1.32	1.25
1:B:194:SER:HB3	1:B:233:ASP:HB3	1.14	1.11
1:B:206:VAL:H	1:B:226:GLN:HE22	1.04	1.00
1:B:194:SER:CB	1:B:233:ASP:HB3	1.99	0.91
1:B:141:ARG:HH11	1:B:294:ASN:HD21	1.20	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/402 (86%)	310 (89%)	24 (7%)	13 (4%)	2	4
1	B	359/402 (89%)	319 (89%)	23 (6%)	17 (5%)	2	2
All	All	706/804 (88%)	629 (89%)	47 (7%)	30 (4%)	2	3

5 of 30 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	GLY
1	A	105	PRO
1	A	172	THR
1	A	213	GLU
1	A	217	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/321 (88%)	252 (90%)	29 (10%)	6	12
1	B	282/321 (88%)	255 (90%)	27 (10%)	7	14
All	All	563/642 (88%)	507 (90%)	56 (10%)	6	13

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	365	VAL
1	B	360	VAL
1	B	101	GLN
1	B	339	LEU
1	B	241	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	188	HIS
1	B	257	GLN
1	B	363	HIS
1	B	294	ASN
1	B	226	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	346/402 (86%)	1.29	81 (23%) <b>2</b> <b>2</b>	74, 86, 100, 111	0
1	B	354/402 (88%)	1.76	133 (37%) <b>1</b> <b>1</b>	75, 87, 99, 106	0
All	All	700/804 (87%)	1.53	214 (30%) <b>1</b> <b>1</b>	74, 86, 100, 111	0

The worst 5 of 214 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	329	ASP	6.2
1	B	164	ALA	6.0
1	A	218	LEU	5.4
1	B	233	ASP	5.2
1	B	334	CYS	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.