



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 20, 2024 – 12:07 pm GMT

PDB ID : 9G8C
Title : Crystal structure of the photosensory core module (PCM) of a cyanophenylalanine mutant oCNF165 of the bathy phytochrome Agp2 from *Agrobacterium fabrum* in the Pfr state.
Authors : Sauthof, L.; Schmidt, A.; Scheerer, P.
Deposited on : 2024-07-23
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	507	
2	B	507	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 8767 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called histidine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	484	Total	C	N	O	S	0	15	0
			3829	2440	678	690	21			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	9IJ	TYR	engineered mutation	UNP A9CI81
A	502	HIS	-	expression tag	UNP A9CI81
A	503	HIS	-	expression tag	UNP A9CI81
A	504	HIS	-	expression tag	UNP A9CI81
A	505	HIS	-	expression tag	UNP A9CI81
A	506	HIS	-	expression tag	UNP A9CI81
A	507	HIS	-	expression tag	UNP A9CI81


- Molecule 2 is a protein called histidine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	488	Total	C	N	O	S	0	13	0
			3815	2425	674	694	22			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	165	9IJ	TYR	engineered mutation	UNP A9CI81
B	502	HIS	-	expression tag	UNP A9CI81
B	503	HIS	-	expression tag	UNP A9CI81
B	504	HIS	-	expression tag	UNP A9CI81
B	505	HIS	-	expression tag	UNP A9CI81
B	506	HIS	-	expression tag	UNP A9CI81
B	507	HIS	-	expression tag	UNP A9CI81


- Molecule 3 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: C₁₂H₂₆O₇).



Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	1
			14	8	6		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is 3-[(2Z)-2-({3-(2-carboxyethyl)-5-[(E)-(4-ethenyl-3-methyl-5-oxo-1,5-dihydro-2H-pyrrol-2-ylidene)methyl]-4-methyl-1H-pyrrol-2-yl}methylidene)-5-{(Z)-[(3E,4S)-3-ethylidene-4-methyl-5-oxopyrrolidin-2-ylidene]methyl}-4-methyl-2H-pyrrol-3-yl]propanoic acid (three-letter code: EL5) (formula: $C_{33}H_{36}N_4O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total 43	C 33	N 4	O 6	0	0
6	B	1	Total 43	C 33	N 4	O 6	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	2	Total Cl 2 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	487	Total O 487 487	0	0
8	B	471	Total O 471 471	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: histidine kinase



- Molecule 2: histidine kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.72Å 93.76Å 174.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.95 – 1.90 45.95 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (45.95-1.90) 99.7 (45.95-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.185 , 0.223 0.196 , 0.233	Depositor DCC
R_{free} test set	4873 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8767	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P6G, 9IJ, CL, MPD, EL5, PEG, MLZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/3802	0.74	0/5155
2	B	0.68	0/3805	0.73	0/5165
All	All	0.67	0/7607	0.74	0/10320

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3829	0	3799	30	0
2	B	3815	0	3748	28	0
3	A	10	0	13	4	0
3	B	9	0	11	1	0
4	A	8	0	14	3	0
4	B	8	0	14	2	0
5	A	7	0	10	1	0
5	B	35	0	50	4	0
6	A	43	0	0	0	0
6	B	43	0	0	0	0
7	B	2	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	487	0	0	2	0
8	B	471	0	0	2	0
All	All	8767	0	7659	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:419:LEU:HD22	1:A:492[B]:LEU:HD22	1.48	0.93
1:A:419:LEU:HD22	1:A:492[B]:LEU:CD2	2.02	0.88
2:B:449:GLU:HB2	5:B:606:PEG:H21	1.60	0.84
2:B:419:LEU:HA	2:B:492[B]:LEU:HD21	1.68	0.75
7:B:608:CL:CL	8:B:1134:HOH:O	2.43	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	477/507 (94%)	470 (98%)	7 (2%)	0	100	100
2	B	479/507 (94%)	473 (99%)	6 (1%)	0	100	100
All	All	956/1014 (94%)	943 (99%)	13 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	391/410 (95%)	387 (99%)	4 (1%)	73	74
2	B	389/411 (95%)	386 (99%)	3 (1%)	79	80
All	All	780/821 (95%)	773 (99%)	7 (1%)	79	77

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	397	LEU
2	B	41	LEU
2	B	373	HIS
2	B	50	LEU
1	A	390	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	35	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

24 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLZ	A	432	1	8,9,10	0.48	0	4,9,11	0.79	0
2	MLZ	B	154[B]	-	8,9,10	0.39	0	4,9,11	0.33	0
2	MLZ	B	308	2	8,9,10	0.42	0	4,9,11	1.29	1 (25%)
1	MLZ	A	180[B]	-	8,9,10	0.40	0	4,9,11	0.98	0
2	MLZ	B	461	2	8,9,10	0.46	0	4,9,11	1.03	0
2	MLZ	B	446	2	8,9,10	0.41	0	4,9,11	0.43	0
1	MLZ	A	446	1	8,9,10	0.42	0	4,9,11	0.61	0
2	MLZ	B	154[A]	-	8,9,10	0.38	0	4,9,11	1.00	0
1	MLZ	A	308	1	8,9,10	0.48	0	4,9,11	1.10	1 (25%)
1	MLZ	A	467	1	8,9,10	0.45	0	4,9,11	0.79	0
2	MLZ	B	467	2	8,9,10	0.39	0	4,9,11	0.70	0
1	MLZ	A	180[A]	-	8,9,10	0.40	0	4,9,11	0.58	0
1	MLZ	A	154[B]	1	8,9,10	0.45	0	4,9,11	0.60	0
1	MLZ	A	461	1	8,9,10	0.40	0	4,9,11	1.31	1 (25%)
2	MLZ	B	174	2	8,9,10	0.43	0	4,9,11	0.71	0
1	MLZ	A	207[B]	-	8,9,10	0.38	0	4,9,11	0.77	0
1	MLZ	A	154[A]	1	8,9,10	0.45	0	4,9,11	0.99	0
2	MLZ	B	180[B]	-	8,9,10	0.41	0	4,9,11	0.56	0
2	MLZ	B	207	2	8,9,10	0.41	0	4,9,11	0.90	0
1	MLZ	A	174	1	8,9,10	0.49	0	4,9,11	0.74	0
1	MLZ	A	207[A]	-	8,9,10	0.38	0	4,9,11	0.79	0
2	MLZ	B	432	2	8,9,10	0.48	0	4,9,11	0.72	0
2	MLZ	B	180[A]	-	8,9,10	0.41	0	4,9,11	1.04	1 (25%)
1	MLZ	A	310	1	8,9,10	0.43	0	4,9,11	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLZ	A	432	1	-	0/7/8/10	-
2	MLZ	B	154[B]	-	-	1/7/8/10	-
2	MLZ	B	308	2	-	1/7/8/10	-
1	MLZ	A	180[B]	-	-	1/7/8/10	-
2	MLZ	B	461	2	-	0/7/8/10	-
2	MLZ	B	446	2	-	0/7/8/10	-
1	MLZ	A	446	1	-	0/7/8/10	-
2	MLZ	B	154[A]	-	-	0/7/8/10	-
1	MLZ	A	308	1	-	1/7/8/10	-
1	MLZ	A	467	1	-	0/7/8/10	-
2	MLZ	B	467	2	-	0/7/8/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLZ	A	180[A]	-	-	0/7/8/10	-
1	MLZ	A	154[B]	1	-	0/7/8/10	-
1	MLZ	A	461	1	-	1/7/8/10	-
2	MLZ	B	174	2	-	1/7/8/10	-
1	MLZ	A	207[B]	-	-	1/7/8/10	-
1	MLZ	A	154[A]	1	-	0/7/8/10	-
2	MLZ	B	180[B]	-	-	0/7/8/10	-
2	MLZ	B	207	2	-	1/7/8/10	-
1	MLZ	A	174	1	-	0/7/8/10	-
1	MLZ	A	207[A]	-	-	2/7/8/10	-
2	MLZ	B	432	2	-	0/7/8/10	-
2	MLZ	B	180[A]	-	-	0/7/8/10	-
1	MLZ	A	310	1	-	2/7/8/10	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	461	MLZ	CM-NZ-CE	2.59	119.42	111.95
2	B	308	MLZ	CM-NZ-CE	2.43	118.97	111.95
1	A	308	MLZ	CM-NZ-CE	2.13	118.09	111.95
2	B	180[A]	MLZ	CM-NZ-CE	2.06	117.89	111.95

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	207[A]	MLZ	O-C-CA-CB
1	A	207[B]	MLZ	O-C-CA-CB
1	A	461	MLZ	CD-CE-NZ-CM
1	A	308	MLZ	CD-CE-NZ-CM
1	A	310	MLZ	CD-CE-NZ-CM

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPD	B	602	-	7,7,7	0.31	0	9,10,10	0.42	0
5	PEG	B	604	-	6,6,6	0.17	0	5,5,5	0.14	0
5	PEG	B	603	-	6,6,6	0.18	0	5,5,5	0.13	0
5	PEG	B	606	-	6,6,6	0.12	0	5,5,5	0.14	0
6	EL5	A	604	1	42,46,46	2.77	12 (28%)	47,67,67	1.50	7 (14%)
3	P6G	B	601	-	8,8,18	0.32	0	7,7,17	0.27	0
5	PEG	A	603	-	6,6,6	0.16	0	5,5,5	0.13	0
4	MPD	A	602	-	7,7,7	0.31	0	9,10,10	0.57	0
5	PEG	B	605[A]	-	6,6,6	0.15	0	5,5,5	0.11	0
5	PEG	B	605[B]	-	6,6,6	0.13	0	5,5,5	0.15	0
6	EL5	B	607	2	42,46,46	2.86	12 (28%)	47,67,67	1.48	8 (17%)
3	P6G	A	601	-	9,9,18	0.29	0	8,8,17	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	B	602	-	-	1/5/5/5	-
5	PEG	B	604	-	-	2/4/4/4	-
5	PEG	B	603	-	-	2/4/4/4	-
5	PEG	B	606	-	-	2/4/4/4	-
6	EL5	A	604	1	-	5/26/74/74	0/4/4/4
3	P6G	B	601	-	-	5/6/6/16	-
5	PEG	A	603	-	-	2/4/4/4	-
4	MPD	A	602	-	-	3/5/5/5	-
5	PEG	B	605[A]	-	-	3/4/4/4	-
5	PEG	B	605[B]	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EL5	B	607	2	-	8/26/74/74	0/4/4/4
3	P6G	A	601	-	-	5/7/7/16	-

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	607	EL5	C2A-C1A	-10.63	1.38	1.51
6	A	604	EL5	C2A-C1A	-9.70	1.40	1.51
6	B	607	EL5	C2A-C3A	-9.52	1.39	1.51
6	A	604	EL5	C2A-C3A	-9.40	1.39	1.51
6	A	604	EL5	CHC-C4B	6.12	1.40	1.35

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	607	EL5	CHD-C1D-C2D	3.94	134.75	126.97
6	A	604	EL5	CHD-C1D-ND	-3.87	117.39	130.40
6	A	604	EL5	CHD-C1D-C2D	3.80	134.47	126.97
6	B	607	EL5	CHD-C1D-ND	-3.73	117.86	130.40
6	B	607	EL5	C2A-C3A-C4A	-3.63	103.83	107.81

There are no chirality outliers.

5 of 40 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	604	EL5	C2D-C1D-CHD-C4C
6	A	604	EL5	ND-C1D-CHD-C4C
6	A	604	EL5	NC-C4C-CHD-C1D
6	B	607	EL5	C2C-C1C-CHC-C4B
6	B	607	EL5	NC-C1C-CHC-C4B

There are no ring outliers.

7 monomers are involved in 15 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	602	MPD	2	0
5	B	606	PEG	3	0
3	B	601	P6G	1	0
5	A	603	PEG	1	0
4	A	602	MPD	3	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	605[B]	PEG	1	0
3	A	601	P6G	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand EL5 B 607




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	473/507 (93%)	0.01	17 (3%)	46	48	12, 23, 42, 64	12 (2%)
2	B	478/507 (94%)	0.19	17 (3%)	46	48	12, 25, 46, 70	11 (2%)
All	All	951/1014 (93%)	0.10	34 (3%)	46	48	12, 24, 44, 70	23 (2%)

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	55	GLY	4.7
2	B	229	VAL	3.9
2	B	6	TYR	3.8
2	B	79	THR	3.6
1	A	80	VAL	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MLZ	B	446	10/11	0.87	0.12	33,34,36,39	0
1	MLZ	A	154[B]	10/11	0.88	0.10	18,19,19,20	10
1	MLZ	A	154[A]	10/11	0.88	0.10	18,20,22,22	10
1	MLZ	A	446	10/11	0.92	0.09	33,35,37,40	0
2	MLZ	B	207	10/11	0.93	0.09	23,29,38,38	0
2	MLZ	B	308	10/11	0.93	0.09	20,23,30,30	0
1	MLZ	A	461	10/11	0.93	0.09	17,22,33,34	0
2	MLZ	B	461	10/11	0.93	0.10	21,25,39,40	0
2	MLZ	B	154[A]	10/11	0.94	0.10	16,17,18,19	7

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MLZ	B	154[B]	10/11	0.94	0.10	16,17,19,19	7
2	MLZ	B	174	10/11	0.94	0.09	15,18,27,28	0
1	MLZ	A	207[B]	10/11	0.94	0.09	22,24,29,30	7
1	MLZ	A	308	10/11	0.94	0.08	18,21,25,26	0
2	MLZ	B	432	10/11	0.94	0.08	18,20,28,30	0
1	MLZ	A	174	10/11	0.94	0.09	14,15,25,25	0
1	MLZ	A	207[A]	10/11	0.94	0.09	22,24,29,30	7
1	MLZ	A	310	10/11	0.95	0.09	20,23,30,30	0
2	MLZ	B	467	10/11	0.95	0.07	19,21,27,27	0
1	MLZ	A	467	10/11	0.96	0.07	16,19,24,24	0
1	MLZ	A	432	10/11	0.96	0.07	16,19,30,31	0
1	MLZ	A	180[A]	10/11	0.96	0.08	18,19,19,20	7
1	MLZ	A	180[B]	10/11	0.96	0.08	18,18,19,19	7
2	MLZ	B	180[A]	10/11	0.96	0.07	14,15,16,16	7
2	MLZ	B	180[B]	10/11	0.96	0.07	14,15,16,16	7

6.3 Carbohydrates

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PEG	B	605[A]	7/7	0.58	0.29	49,56,60,61	7
5	PEG	B	605[B]	7/7	0.58	0.29	28,31,33,34	7
3	P6G	B	601	9/19	0.69	0.22	35,40,44,46	0
5	PEG	B	606	7/7	0.69	0.16	52,53,55,57	0
5	PEG	B	603	7/7	0.77	0.17	40,50,57,59	0
5	PEG	A	603	7/7	0.77	0.20	54,55,59,59	0
3	P6G	A	601	10/19	0.78	0.18	24,36,52,54	0
5	PEG	B	604	7/7	0.81	0.18	55,57,59,59	0
4	MPD	A	602	8/8	0.90	0.11	23,25,26,27	0
6	EL5	A	604	43/43	0.92	0.09	17,20,25,27	0
4	MPD	B	602	8/8	0.93	0.08	20,21,23,25	0
6	EL5	B	607	43/43	0.93	0.08	18,21,24,30	0
7	CL	B	609	1/1	0.94	0.19	64,64,64,64	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	CL	B	608	1/1	0.95	0.07	38,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P6G B 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around P6G A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EL5 A 604:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around EL5 B 607:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.