



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 08:03 AM EDT

PDB ID : 4GZ7  
Title : The crystal structure of Apo-dihydropyrimidinase from Tetraodon nigroviridis  
Authors : Hsien, Y.C.; Chen, M.C.; Hsu, C.C.; Chan, S.I.; Yang, Y.S.; Chen, C.J.  
Deposited on : 2012-09-06  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	500	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MHA	A	601	-	X	X	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MHA	A	603	-	X	X	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dihydropyrimidinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	492	Total	C	N	O	S	0	0	0
			3805	2396	673	712	24			

- Molecule 2 is (CARBAMOYLMETHYL-CARBOXYMETHYL-AMINO)-ACETIC ACID (three-letter code: MHA) (formula: C<sub>6</sub>H<sub>10</sub>N<sub>2</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			13	6	2	5		
2	A	1	Total	C	N	O	0	0
			13	6	2	5		
2	A	1	Total	C	N	O	0	0
			13	6	2	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	232	Total 232	O 232	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: dihydropyrimidinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	160.82Å 160.82Å 94.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 24.51 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.00-2.00) 99.4 (24.51-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.22 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.159 , 0.195 0.159 , 0.199	Depositor DCC
$R_{free}$ test set	3145 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.1	Xtriage
Anisotropy	0.598	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4076	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MHA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.20	7/3888 (0.2%)	1.04	18/5265 (0.3%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	460	HIS	CG-CD2	6.11	1.46	1.35
1	A	281	TRP	CD2-CE2	5.77	1.48	1.41
1	A	160	TYR	CE2-CZ	5.42	1.45	1.38
1	A	160	TYR	CG-CD1	5.39	1.46	1.39
1	A	204	SER	CB-OG	5.25	1.49	1.42
1	A	170	GLU	CD-OE1	5.16	1.31	1.25
1	A	376	SER	CA-CB	5.09	1.60	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	390	ARG	NE-CZ-NH1	7.82	124.21	120.30
1	A	167	ARG	NE-CZ-NH1	-7.34	116.63	120.30
1	A	448	ARG	NE-CZ-NH1	6.76	123.68	120.30
1	A	46	ARG	NE-CZ-NH1	-6.75	116.92	120.30
1	A	114	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	A	13	LYS	CD-CE-NZ	6.10	125.72	111.70
1	A	228	ARG	NE-CZ-NH2	-6.07	117.26	120.30
1	A	439	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	A	441	ARG	NE-CZ-NH1	5.83	123.22	120.30
1	A	318	LEU	CB-CG-CD1	-5.66	101.38	111.00
1	A	470	LYS	CD-CE-NZ	-5.54	98.96	111.70
1	A	369	ARG	NE-CZ-NH1	-5.45	117.58	120.30
1	A	408	ARG	NE-CZ-NH1	5.43	123.02	120.30
1	A	441	ARG	NE-CZ-NH2	-5.29	117.66	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	473	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	163	LEU	CA-CB-CG	5.19	127.24	115.30
1	A	46	ARG	CA-CB-CG	-5.17	102.04	113.40
1	A	420	ASP	CB-CG-OD1	5.13	122.92	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3805	0	3753	52	0
2	A	39	0	25	23	0
3	A	232	0	0	5	0
All	All	4076	0	3778	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293:MET:HB2	2:A:601:MHA:HN22	1.21	1.03
2:A:603:MHA:N2	2:A:603:MHA:C3	2.23	1.01
1:A:213:HIS:ND1	2:A:603:MHA:N2	2.14	0.95
1:A:16:ASN:HD21	1:A:391:ILE:H	1.02	0.94
2:A:603:MHA:N2	2:A:603:MHA:HC32	1.55	0.93
2:A:603:MHA:HN21	2:A:603:MHA:C4	1.82	0.91
1:A:415:HIS:HD2	1:A:422:ASN:HD22	1.22	0.87
1:A:466:GLU:HG3	1:A:470:LYS:HE3	1.57	0.85
2:A:601:MHA:O3	2:A:601:MHA:HC11	1.83	0.76
1:A:293:MET:CB	2:A:601:MHA:HN22	2.00	0.73
1:A:104:GLN:HG2	1:A:105:SER:N	2.04	0.73
2:A:603:MHA:C3	2:A:603:MHA:HN21	1.91	0.73
2:A:603:MHA:N2	2:A:603:MHA:C4	2.53	0.71

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:466:GLU:CG	1:A:470:LYS:HE3	2.19	0.70
1:A:16:ASN:ND2	1:A:391:ILE:H	1.83	0.70
1:A:8:LEU:HD23	1:A:10:LYS:HD2	1.73	0.70
2:A:601:MHA:C6	2:A:601:MHA:C2	2.73	0.67
1:A:48:ILE:HD12	1:A:48:ILE:N	2.10	0.66
1:A:344:GLY:HA2	2:A:601:MHA:HN21	1.60	0.65
2:A:601:MHA:C5	2:A:601:MHA:O2	2.44	0.65
1:A:448:ARG:HH11	1:A:448:ARG:HG2	1.63	0.63
2:A:603:MHA:HC31	3:A:864:HOH:O	1.98	0.63
1:A:177:HIS:HD2	3:A:818:HOH:O	1.82	0.61
1:A:372:ALA:HA	1:A:376:SER:HB2	1.81	0.61
1:A:213:HIS:CE1	2:A:603:MHA:HN22	2.13	0.61
1:A:448:ARG:HH11	1:A:448:ARG:CG	2.14	0.60
1:A:48:ILE:N	1:A:48:ILE:CD1	2.64	0.60
1:A:101:GLN:HG3	1:A:104:GLN:HB2	1.83	0.60
1:A:322:ASP:OD1	2:A:601:MHA:O2	2.21	0.58
1:A:344:GLY:HA2	2:A:601:MHA:N2	2.18	0.58
1:A:244:HIS:CE1	1:A:295:PRO:HD3	2.40	0.55
1:A:445:GLU:OE2	1:A:450:LYS:NZ	2.39	0.55
1:A:285:TRP:CZ2	1:A:335:LYS:HA	2.42	0.54
1:A:448:ARG:HG2	1:A:448:ARG:NH1	2.22	0.54
1:A:7:ILE:HG23	1:A:48:ILE:HD13	1.89	0.54
1:A:101:GLN:HG3	1:A:104:GLN:CB	2.38	0.54
2:A:601:MHA:O2	2:A:601:MHA:HC52	2.08	0.53
1:A:47:VAL:C	1:A:48:ILE:HD12	2.29	0.52
2:A:601:MHA:C6	2:A:601:MHA:O2	2.59	0.51
1:A:401:ILE:HD13	1:A:435:VAL:HB	1.91	0.51
2:A:603:MHA:N2	2:A:603:MHA:O3	2.42	0.50
1:A:15:VAL:HB	1:A:56:ILE:HG22	1.94	0.49
1:A:120:LYS:HD2	3:A:903:HOH:O	2.13	0.48
1:A:323:ASN:HD22	1:A:348:VAL:H	1.61	0.47
1:A:198:GLY:C	1:A:216:CYS:HB2	2.35	0.47
1:A:74:ARG:NH2	3:A:797:HOH:O	2.46	0.45
1:A:415:HIS:CD2	1:A:422:ASN:HD22	2.14	0.45
1:A:322:ASP:OD2	2:A:601:MHA:O2	2.34	0.45
1:A:294:GLY:O	2:A:601:MHA:O1	2.35	0.44
1:A:139:LYS:O	1:A:177:HIS:HE1	2.00	0.43
1:A:146:ALA:HA	1:A:151:VAL:O	2.19	0.43
1:A:87:LEU:HD21	1:A:123:CYS:HB3	2.00	0.43
1:A:74:ARG:NH2	3:A:819:HOH:O	2.48	0.43
1:A:40:ARG:HA	1:A:40:ARG:HD3	1.81	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:ASN:OD1	1:A:287:HIS:HE1	2.03	0.42
1:A:40:ARG:O	1:A:42:PRO:HD3	2.18	0.42
1:A:448:ARG:CG	1:A:448:ARG:NH1	2.78	0.42
1:A:415:HIS:HD2	1:A:422:ASN:ND2	2.03	0.42
1:A:101:GLN:H	1:A:101:GLN:HG2	1.61	0.42
1:A:338:PHE:HE2	2:A:603:MHA:HC52	1.84	0.41
1:A:448:ARG:HH11	1:A:448:ARG:CB	2.33	0.41
1:A:214:GLU:HG2	1:A:296:PRO:HB2	2.02	0.41
1:A:322:ASP:CG	2:A:601:MHA:O2	2.60	0.40
1:A:89:GLY:HA3	1:A:349:GLU:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	490/500 (98%)	474 (97%)	15 (3%)	1 (0%)	47 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	383	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	404/410 (98%)	397 (98%)	7 (2%)	60 65

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	THR
1	A	102	LYS
1	A	244	HIS
1	A	415	HIS
1	A	448	ARG
1	A	466	GLU
1	A	492	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	177	HIS
1	A	287	HIS
1	A	290	GLN
1	A	323	ASN
1	A	343	ASN
1	A	415	HIS
1	A	417	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MHA	A	602	-	12,12,12	1.56	2 (16%)	15,15,15	1.37	3 (20%)
2	MHA	A	601	-	12,12,12	2.02	5 (41%)	15,15,15	1.47	4 (26%)
2	MHA	A	603	-	12,12,12	2.06	5 (41%)	15,15,15	3.07	9 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MHA	A	602	-	-	0/12/12/12	-
2	MHA	A	601	-	-	7/12/12/12	-
2	MHA	A	603	-	-	9/12/12/12	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	603	MHA	O3-C4	3.86	1.43	1.30
2	A	601	MHA	C5-C6	3.53	1.58	1.52
2	A	603	MHA	C5-C6	3.22	1.58	1.52
2	A	601	MHA	O1-C2	-3.15	1.20	1.30
2	A	602	MHA	C3-C4	3.15	1.57	1.51
2	A	601	MHA	C1-C2	2.87	1.57	1.51
2	A	603	MHA	C1-C2	2.50	1.56	1.51
2	A	601	MHA	O4-C4	2.38	1.29	1.22
2	A	601	MHA	C1-N1	2.27	1.51	1.47
2	A	603	MHA	C3-C4	2.21	1.55	1.51
2	A	603	MHA	O5-C6	-2.21	1.17	1.23
2	A	602	MHA	C5-N1	2.10	1.51	1.47

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	603	MHA	O4-C4-C3	-6.17	96.59	122.38
2	A	603	MHA	C5-N1-C3	-4.70	100.63	111.66
2	A	603	MHA	C5-N1-C1	-4.67	100.69	111.66
2	A	603	MHA	O3-C4-O4	4.30	134.38	123.33
2	A	603	MHA	O3-C4-C3	3.93	128.86	113.38
2	A	601	MHA	C5-N1-C1	2.92	118.50	111.66
2	A	603	MHA	C5-C6-N2	2.81	121.27	115.61
2	A	602	MHA	C6-C5-N1	-2.58	109.70	114.42
2	A	601	MHA	O5-C6-C5	2.36	124.91	120.77
2	A	601	MHA	C3-N1-C1	2.18	116.76	111.66
2	A	603	MHA	O5-C6-N2	-2.15	116.80	122.53
2	A	603	MHA	C4-C3-N1	2.08	120.37	113.77
2	A	602	MHA	O2-C2-C1	-2.08	113.68	122.38
2	A	601	MHA	C2-C1-N1	-2.08	107.18	113.77
2	A	602	MHA	O1-C2-O2	2.03	128.55	123.33
2	A	603	MHA	C3-N1-C1	2.01	116.36	111.66

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	MHA	C2-C1-N1-C5
2	A	601	MHA	C6-C5-N1-C1
2	A	601	MHA	N1-C3-C4-O3
2	A	603	MHA	C6-C5-N1-C3
2	A	603	MHA	N1-C1-C2-O1
2	A	603	MHA	N1-C3-C4-O3
2	A	603	MHA	N1-C3-C4-O4
2	A	601	MHA	N1-C3-C4-O4
2	A	603	MHA	N1-C1-C2-O2
2	A	601	MHA	C4-C3-N1-C1
2	A	603	MHA	C2-C1-N1-C5
2	A	603	MHA	C4-C3-N1-C5
2	A	603	MHA	C4-C3-N1-C1
2	A	601	MHA	C2-C1-N1-C3
2	A	603	MHA	C2-C1-N1-C3
2	A	601	MHA	C4-C3-N1-C5

There are no ring outliers.

2 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	MHA	13	0
2	A	603	MHA	10	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	492/500 (98%)	-0.58	8 (1%) 72 70	9, 16, 37, 70	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	ALA	4.4
1	A	72	GLY	4.2
1	A	71	MET	3.3
1	A	495	ALA	2.8
1	A	40	ARG	2.7
1	A	490	ALA	2.6
1	A	492	GLU	2.6
1	A	487	GLU	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MHA	A	603	13/13	0.77	0.29	27,38,46,47	0
2	MHA	A	601	13/13	0.86	0.26	27,40,48,49	0
2	MHA	A	602	13/13	0.92	0.15	23,38,45,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.