



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 1, 2025 – 11:30 pm BST

PDB ID : 1H0C / pdb_00001h0c
Title : The crystal structure of human alanine:glyoxylate aminotransferase
Authors : Zhang, X.; Danpure, C.J.; Roe, S.M.; Pearl, L.H.
Deposited on : 2002-06-17
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	<div> <div>11%</div> <div>65%</div> <div>29%</div> <div>..</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE–GLYOXYLATE AMINOTRANSFERASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	386	2953	1888	513	536	16	0	0	1

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	15	8	1	5	1	0	0

- Molecule 3 is (AMINOXY)ACETIC ACID (CCD ID: AOA) (formula: $C_2H_5NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			6	2	1	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	143	Total	O	0	0
			143	143		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SERINE–GLYOXYLATE AMINOTRANSFERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	90.33Å 90.33Å 142.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.49 – 2.50 47.49 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.3 (47.49-2.50) 98.3 (47.49-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.57 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.231 , 0.286 0.225 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.695	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3153	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AOA, PLP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/3022	0.81	0/4104

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2953	0	2969	103	0
2	A	15	0	7	1	0
3	A	6	0	5	0	0
4	A	36	0	48	1	0
5	A	143	0	0	7	1
All	All	3153	0	3029	103	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:LEU:HD12	1:A:136:LEU:H	1.12	1.10
1:A:5:LYS:H	1:A:5:LYS:HD2	1.30	0.94
1:A:208:GLN:H	1:A:208:GLN:HE21	1.11	0.92
1:A:25:LEU:H	1:A:32:ASN:HD21	1.20	0.85
1:A:107:ILE:HG21	1:A:351:LEU:HD13	1.59	0.83
1:A:378:GLU:HG2	1:A:381:ARG:HH12	1.43	0.83
1:A:83:HIS:HB3	5:A:2044:HOH:O	1.81	0.81
1:A:136:LEU:H	1:A:136:LEU:CD1	1.86	0.81
1:A:32:ASN:HD22	1:A:367:ASN:HD21	1.31	0.76
1:A:378:GLU:HG2	1:A:381:ARG:NH1	2.00	0.76
1:A:208:GLN:HE21	1:A:208:GLN:N	1.83	0.75
1:A:106:GLY:HA3	1:A:157:GLU:OE1	1.86	0.75
1:A:36:ARG:HD3	1:A:278:LEU:HD11	1.67	0.74
1:A:25:LEU:H	1:A:32:ASN:ND2	1.86	0.74
1:A:28:PRO:O	1:A:209:LYS:HE3	1.88	0.73
1:A:128:LYS:HE2	1:A:133:HIS:O	1.89	0.72
1:A:208:GLN:H	1:A:208:GLN:NE2	1.87	0.70
1:A:101:LEU:HB2	1:A:148:PRO:HG2	1.76	0.68
1:A:329:GLY:O	1:A:389:LYS:HA	1.97	0.65
1:A:369:THR:O	1:A:373:VAL:HG23	1.96	0.64
1:A:307:LEU:HD22	1:A:327:PRO:HG3	1.79	0.64
1:A:297:TYR:CD2	1:A:370:ARG:HG2	2.32	0.64
1:A:98:ASP:O	1:A:99:SER:HB3	1.98	0.63
1:A:136:LEU:HD12	1:A:136:LEU:N	1.98	0.62
1:A:237:PRO:HA	5:A:2080:HOH:O	1.99	0.61
1:A:250:PHE:O	1:A:258:ARG:HD3	2.00	0.61
1:A:150:LEU:HD11	1:A:181:LEU:HB2	1.84	0.59
1:A:125:PRO:O	1:A:127:THR:N	2.35	0.58
1:A:68:PHE:O	1:A:69:GLN:HB2	2.01	0.58
1:A:95:GLU:OE1	1:A:233:ARG:HB2	2.04	0.58
1:A:147:LYS:N	1:A:148:PRO:HD3	2.18	0.57
1:A:197:ARG:HG3	5:A:2067:HOH:O	2.06	0.56
1:A:243:ASP:OD1	1:A:245:LYS:HB2	2.06	0.55
1:A:55:GLN:O	1:A:59:GLU:HG3	2.09	0.53
1:A:175:ARG:HG3	1:A:175:ARG:NH1	2.23	0.53
1:A:254:ASP:HB3	1:A:256:GLN:H	1.73	0.53
1:A:127:THR:HG22	1:A:128:LYS:N	2.24	0.53
1:A:143:LEU:HD13	1:A:176:TYR:HB2	1.91	0.52
1:A:98:ASP:O	1:A:99:SER:CB	2.58	0.52
1:A:113:VAL:HG13	1:A:123:VAL:HG11	1.92	0.52
1:A:175:ARG:CG	1:A:175:ARG:HH11	2.22	0.52
1:A:174:HIS:CD2	1:A:225:LYS:HE2	2.45	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:GLU:HG3	1:A:160:THR:OG1	2.10	0.51
1:A:338:TYR:CE1	1:A:342:HIS:CD2	2.99	0.51
1:A:45:MET:HE2	1:A:267:ILE:HD11	1.93	0.51
1:A:371:GLU:O	1:A:374:ASP:HB2	2.10	0.51
1:A:141:GLU:O	1:A:144:ALA:HB3	2.11	0.50
1:A:175:ARG:HG3	1:A:175:ARG:HH11	1.76	0.50
1:A:312:LYS:O	1:A:313:ASP:CB	2.60	0.50
1:A:327:PRO:HB2	1:A:330:TYR:CD1	2.47	0.50
1:A:77:VAL:HB	1:A:251:TRP:CZ2	2.47	0.50
1:A:295:ALA:O	1:A:299:HIS:CD2	2.65	0.49
1:A:104:ALA:HA	1:A:109:GLY:HA3	1.93	0.49
1:A:36:ARG:HD3	1:A:278:LEU:CD1	2.39	0.49
1:A:348:MET:HB3	5:A:2130:HOH:O	2.13	0.49
1:A:76:LEU:HD12	1:A:76:LEU:C	2.33	0.49
1:A:371:GLU:OE1	1:A:371:GLU:N	2.40	0.48
1:A:307:LEU:HD21	1:A:384:LEU:HD21	1.95	0.48
1:A:143:LEU:HD11	1:A:173:CYS:HA	1.96	0.47
1:A:314:PRO:HG3	1:A:317:ARG:NH1	2.28	0.47
1:A:312:LYS:O	1:A:313:ASP:HB2	2.14	0.47
1:A:285:GLU:HB2	5:A:2113:HOH:O	2.13	0.47
1:A:146:HIS:O	1:A:147:LYS:C	2.54	0.47
1:A:78:ILE:HB	1:A:218:SER:OG	2.15	0.46
1:A:127:THR:CG2	1:A:128:LYS:N	2.79	0.46
1:A:48:SER:HB2	5:A:2096:HOH:O	2.16	0.46
1:A:339:VAL:HG12	1:A:345:ILE:O	2.15	0.46
1:A:146:HIS:HB2	1:A:148:PRO:HG3	1.97	0.46
1:A:164:GLN:HA	1:A:165:PRO:HD3	1.75	0.45
1:A:140:GLU:HG3	1:A:176:TYR:OH	2.16	0.45
1:A:5:LYS:H	1:A:5:LYS:CD	2.11	0.45
1:A:65:GLN:O	1:A:69:GLN:N	2.47	0.44
1:A:246:TRP:HB3	1:A:261:HIS:CD2	2.52	0.44
1:A:216:GLY:HA3	5:A:2075:HOH:O	2.17	0.44
1:A:302:LEU:HD13	1:A:324:VAL:HG22	2.00	0.44
1:A:5:LYS:HD2	1:A:5:LYS:N	2.13	0.43
1:A:247:LEU:HD12	1:A:247:LEU:HA	1.74	0.43
1:A:318:LEU:HD12	1:A:319:PRO:HD2	2.00	0.43
1:A:37:ILE:HD11	1:A:278:LEU:HD22	2.01	0.43
1:A:44:GLN:HG3	1:A:45:MET:N	2.33	0.43
1:A:302:LEU:HD13	1:A:324:VAL:CG2	2.48	0.43
1:A:107:ILE:O	1:A:110:GLN:HB2	2.18	0.43
1:A:385:GLN:HE21	1:A:385:GLN:HB3	1.65	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:PRO:HD3	1:A:212:ASN:HB3	2.01	0.43
1:A:191:THR:CG2	1:A:319:PRO:HD2	2.48	0.43
1:A:107:ILE:HG13	4:A:1397:GOL:O3	2.19	0.43
1:A:183:ASP:OD1	2:A:1391:PLP:H2A2	2.19	0.43
1:A:171:GLU:O	1:A:175:ARG:HB2	2.19	0.42
1:A:191:THR:HG22	1:A:319:PRO:HG2	2.00	0.42
1:A:384:LEU:HD12	1:A:384:LEU:HA	1.87	0.42
1:A:169:PHE:O	1:A:170:GLY:C	2.56	0.42
1:A:209:LYS:HB3	1:A:210:ALA:H	1.77	0.42
1:A:60:ILE:O	1:A:64:ILE:HG13	2.19	0.42
1:A:143:LEU:HD23	1:A:143:LEU:HA	1.85	0.42
1:A:155:HIS:CG	1:A:166:LEU:HD11	2.55	0.42
1:A:326:VAL:HA	1:A:327:PRO:HD3	1.83	0.42
1:A:191:THR:HG21	1:A:319:PRO:HD2	2.03	0.41
1:A:101:LEU:O	1:A:151:LEU:HD12	2.20	0.41
1:A:350:GLY:HA3	1:A:355:THR:HA	2.02	0.41
1:A:317:ARG:O	1:A:319:PRO:HD3	2.21	0.41
1:A:151:LEU:O	1:A:181:LEU:HB3	2.20	0.41
1:A:124:HIS:HA	1:A:125:PRO:HD3	1.83	0.41
1:A:234:LYS:HA	1:A:234:LYS:HD2	1.74	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2043:HOH:O	5:A:2043:HOH:O[7_555]	1.23	0.97

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	382/392 (97%)	346 (91%)	31 (8%)	5 (1%)	10 19

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	ASP
1	A	328	ALA
1	A	126	MET
1	A	145	GLN
1	A	388	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/326 (97%)	298 (94%)	18 (6%)	17	35

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	18	LEU
1	A	20	ILE
1	A	95	GLU
1	A	136	LEU
1	A	146	HIS
1	A	147	LYS
1	A	155	HIS
1	A	175	ARG
1	A	208	GLN
1	A	211	LEU
1	A	234	LYS
1	A	247	LEU
1	A	278	LEU
1	A	323	THR
1	A	348	MET
1	A	351	LEU
1	A	384	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	ASN
1	A	72	ASN
1	A	155	HIS
1	A	208	GLN
1	A	299	HIS
1	A	342	HIS
1	A	385	GLN
1	A	386	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	1393	-	5,5,5	0.66	0	5,5,5	0.73	0
3	AOA	A	1392	-	3,5,5	1.14	0	3,5,5	1.43	0
4	GOL	A	1395	-	5,5,5	0.31	0	5,5,5	0.59	0
4	GOL	A	1394	-	5,5,5	0.71	0	5,5,5	0.84	0
4	GOL	A	1398	-	5,5,5	0.55	0	5,5,5	0.80	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	1397	-	5,5,5	0.36	0	5,5,5	0.71	0
2	PLP	A	1391	1	15,15,16	4.55	7 (46%)	20,22,23	1.13	2 (10%)
4	GOL	A	1396	-	5,5,5	0.53	0	5,5,5	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1393	-	-	0/4/4/4	-
3	AOA	A	1392	-	-	2/2/3/3	-
4	GOL	A	1395	-	-	0/4/4/4	-
4	GOL	A	1394	-	-	0/4/4/4	-
4	GOL	A	1398	-	-	0/4/4/4	-
4	GOL	A	1397	-	-	0/4/4/4	-
2	PLP	A	1391	1	-	0/6/6/8	0/1/1/1
4	GOL	A	1396	-	-	0/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1391	PLP	C5-C4	11.78	1.53	1.40
2	A	1391	PLP	C3-C2	10.14	1.51	1.40
2	A	1391	PLP	C6-N1	4.79	1.44	1.34
2	A	1391	PLP	P-O1P	4.32	1.64	1.50
2	A	1391	PLP	P-O3P	3.50	1.68	1.54
2	A	1391	PLP	P-O4P	2.37	1.67	1.60
2	A	1391	PLP	O4P-C5A	-2.01	1.37	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1391	PLP	O3P-P-O4P	2.46	113.27	106.73
2	A	1391	PLP	C5-C6-N1	-2.25	120.08	123.82

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1392	AOA	O1-C1-C2-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	1392	AOA	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1397	GOL	1	0
2	A	1391	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	386/392 (98%)	0.66	42 (10%)	12 11	18, 39, 69, 82	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	136	LEU	5.5
1	A	391	LYS	5.5
1	A	238	PHE	4.4
1	A	143	LEU	3.6
1	A	120	GLY	3.6
1	A	144	ALA	3.5
1	A	97	GLY	3.4
1	A	148	PRO	3.4
1	A	4	HIS	3.2
1	A	142	GLY	3.2
1	A	141	GLU	3.1
1	A	176	TYR	3.1
1	A	22	ASN	3.0
1	A	390	LYS	3.0
1	A	134	TYR	3.0
1	A	175	ARG	2.9
1	A	131	GLY	2.8
1	A	5	LYS	2.8
1	A	145	GLN	2.8
1	A	101	LEU	2.7
1	A	127	THR	2.7
1	A	329	GLY	2.7
1	A	123	VAL	2.6
1	A	235	THR	2.6
1	A	304	ALA	2.5
1	A	93	VAL	2.5
1	A	137	GLN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	135	THR	2.4
1	A	172	LEU	2.4
1	A	128	LYS	2.4
1	A	168	GLY	2.4
1	A	387	CYS	2.4
1	A	234	LYS	2.3
1	A	138	GLU	2.3
1	A	328	ALA	2.3
1	A	232	SER	2.3
1	A	305	LEU	2.3
1	A	333	ARG	2.3
1	A	388	PRO	2.2
1	A	383	ALA	2.2
1	A	20	ILE	2.1
1	A	132	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	1397	6/6	0.72	0.26	65,68,68,70	0
4	GOL	A	1393	6/6	0.79	0.21	67,69,72,73	0
4	GOL	A	1398	6/6	0.79	0.19	45,49,51,52	0
4	GOL	A	1396	6/6	0.82	0.21	56,59,60,60	0
4	GOL	A	1395	6/6	0.85	0.20	63,63,64,66	0
3	AOA	A	1392	6/6	0.85	0.16	49,51,55,57	0
4	GOL	A	1394	6/6	0.87	0.16	45,49,50,52	0
2	PLP	A	1391	15/16	0.95	0.13	31,41,44,45	0

6.5 Other polymers ⓘ

There are no such residues in this entry.