



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 1, 2025 – 10:41 pm BST

PDB ID : 6H1A / pdb_00006h1a
Title : Crystal structure of VX surrogate NEMP inhibited recombinant human bile salt activated lipase
Authors : Touvrey, C.; Brazzolotto, X.; Nachon, F.
Deposited on : 2018-07-11
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

| | | |
|--------------------------------|---|--|
| MolProbity | : | 4.02b-467 |
| Mogul | : | 1.8.4, CSD as541be (2020) |
| Xtriage (Phenix) | : | 1.13 |
| EDS | : | 3.0 |
| Percentile statistics | : | 20231227.v01 (using entries in the PDB archive December 27th 2023) |
| CCP4 | : | 9.0.003 (Gargrove) |
| Density-Fitness | : | 1.0.11 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.42 |


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 164625 | 2888 (1.76-1.76) |
| Clashscore | 180529 | 3097 (1.76-1.76) |
| Ramachandran outliers | 177936 | 3072 (1.76-1.76) |
| Sidechain outliers | 177891 | 3072 (1.76-1.76) |
| RSRZ outliers | 164620 | 2887 (1.76-1.76) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 1 | A | 547 | <div> <div>6%</div> <div>88%</div> <div>7%</div> </div> |

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bile salt-activated lipase.

| Mol | Chain | Residues | Atoms | | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|----|---------|---------|-------|
| 1 | A | 523 | Total | C | N | O | P | S | 0 | 6 | 0 |
| | | | 4104 | 2638 | 686 | 764 | 1 | 15 | | | |

There are 17 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| A | -13 | HIS | - | expression tag | UNP P19835 |
| A | -12 | HIS | - | expression tag | UNP P19835 |
| A | -11 | HIS | - | expression tag | UNP P19835 |
| A | -10 | HIS | - | expression tag | UNP P19835 |
| A | -9 | HIS | - | expression tag | UNP P19835 |
| A | -8 | HIS | - | expression tag | UNP P19835 |
| A | -7 | HIS | - | expression tag | UNP P19835 |
| A | -6 | HIS | - | expression tag | UNP P19835 |
| A | -5 | GLU | - | expression tag | UNP P19835 |
| A | -4 | ASN | - | expression tag | UNP P19835 |
| A | -3 | LEU | - | expression tag | UNP P19835 |
| A | -2 | TYR | - | expression tag | UNP P19835 |
| A | -1 | PHE | - | expression tag | UNP P19835 |
| A | 0 | GLN | - | expression tag | UNP P19835 |
| A | 1 | SER | - | expression tag | UNP P19835 |
| A | 186 | ASP | ASN | engineered mutation | UNP P19835 |
| A | 298 | ASP | ALA | engineered mutation | UNP P19835 |

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 2 | A | 11 | Total | Zn | 0 | 0 |
| | | | 11 | 11 | | |

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 3 | A | 1 | Total | C | O | 0 | 0 |
| | | | 4 | 2 | 2 | | |
| 3 | A | 1 | Total | C | O | 0 | 0 |
| | | | 4 | 2 | 2 | | |
| 3 | A | 1 | Total | C | O | 0 | 0 |
| | | | 4 | 2 | 2 | | |
| 3 | A | 1 | Total | C | O | 0 | 0 |
| | | | 4 | 2 | 2 | | |
| 3 | A | 1 | Total | C | O | 0 | 0 |
| | | | 4 | 2 | 2 | | |
| 3 | A | 1 | Total | C | O | 0 | 0 |
| | | | 4 | 2 | 2 | | |

- Molecule 4 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|-----|---------|---------|
| 4 | A | 196 | Total | O | 0 | 0 |
| | | | 196 | 196 | | |

- Molecule 1: Bile salt-activated lipase



4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 21 21 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 56.37Å 97.74Å 110.83Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 48.21 – 1.75 48.21 – 1.75 | Depositor EDS |
| % Data completeness (in resolution range) | 98.9 (48.21-1.75) 99.1 (48.21-1.75) | Depositor EDS |
| R_{merge} | 0.07 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.23 (at 1.75Å) | Xtriage |
| Refinement program | PHENIX | Depositor |
| R, R_{free} | 0.194 , 0.224 0.194 , 0.224 | Depositor DCC |
| R_{free} test set | 3053 reflections (4.88%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 26.5 | Xtriage |
| Anisotropy | 0.610 | Xtriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.32 , 28.2 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$ | Xtriage |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| F_o, F_c correlation | 0.96 | EDS |
| Total number of atoms | 4339 | wwPDB-VP |
| Average B, all atoms (Å ²) | 34.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ACT, SVX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|-------------|-------------|-------------|
| | | RMSZ | $\# Z > 5$ | RMSZ | $\# Z > 5$ |
| 1 | A | 0.37 | 0/4214 | 0.54 | 0/5731 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 4104 | 0 | 4047 | 29 | 0 |
| 2 | A | 11 | 0 | 0 | 0 | 0 |
| 3 | A | 28 | 0 | 21 | 1 | 0 |
| 4 | A | 196 | 0 | 0 | 8 | 0 |
| All | All | 4339 | 0 | 4068 | 29 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|--------------------|------------------|--------------------------|-------------------|
| 1:A:438:ASP:OD1 | 4:A:701:HOH:O | 2.04 | 0.76 |
| 1:A:46:GLN:NE2 | 4:A:704:HOH:O | 2.15 | 0.71 |
| 1:A:299:ASP:OD1 | 4:A:702:HOH:O | 2.09 | 0.70 |
| 1:A:281:MET:HE2 | 1:A:327:ILE:HG21 | 1.74 | 0.69 |
| 1:A:285:VAL:O | 4:A:703:HOH:O | 2.14 | 0.66 |
| 1:A:63:ARG:CZ | 1:A:75:TYR:OH | 2.50 | 0.60 |
| 1:A:358:ARG:NH1 | 4:A:707:HOH:O | 2.36 | 0.57 |
| 1:A:53:GLY:O | 4:A:705:HOH:O | 2.17 | 0.57 |
| 1:A:317[B]:ASN:ND2 | 4:A:701:HOH:O | 2.37 | 0.57 |
| 1:A:317[A]:ASN:ND2 | 1:A:436:ALA:O | 2.38 | 0.56 |
| 1:A:193:GLU:OE1 | 1:A:439:ILE:HD11 | 2.06 | 0.56 |
| 1:A:355:LYS:HG3 | 1:A:526:TYR:CZ | 2.43 | 0.53 |
| 1:A:63:ARG:HD3 | 1:A:75:TYR:CE1 | 2.45 | 0.52 |
| 1:A:493:THR:HG22 | 4:A:708:HOH:O | 2.10 | 0.52 |
| 1:A:281:MET:HE3 | 1:A:327:ILE:HD13 | 1.92 | 0.50 |
| 1:A:281:MET:CE | 1:A:327:ILE:HG21 | 2.46 | 0.44 |
| 1:A:72:ASP:OD1 | 1:A:73:SER:N | 2.51 | 0.44 |
| 1:A:419[A]:SER:HB2 | 1:A:502:THR:HG22 | 2.00 | 0.44 |
| 1:A:513:SER:HB3 | 1:A:516:THR:HG21 | 1.99 | 0.43 |
| 1:A:498:TYR:CE2 | 1:A:514:LEU:HB2 | 2.54 | 0.43 |
| 1:A:134:GLY:HA3 | 1:A:469:THR:HG21 | 2.01 | 0.43 |
| 1:A:75:TYR:HE1 | 3:A:616:ACT:OXT | 2.03 | 0.42 |
| 1:A:434:ASP:OD1 | 1:A:437:ASP:HB2 | 2.20 | 0.42 |
| 1:A:298:ASP:OD2 | 1:A:299:ASP:N | 2.53 | 0.42 |
| 1:A:279:TYR:HB3 | 1:A:280:PRO:HD2 | 2.03 | 0.41 |
| 1:A:446:PRO:HA | 1:A:453:TYR:CD1 | 2.55 | 0.41 |
| 1:A:513:SER:HB3 | 1:A:516:THR:CG2 | 2.51 | 0.41 |
| 1:A:513:SER:C | 1:A:516:THR:HG23 | 2.41 | 0.41 |
| 1:A:398:GLU:OE2 | 1:A:519:LEU:HD22 | 2.21 | 0.40 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 1 | A | 524/547 (96%) | 509 (97%) | 15 (3%) | 0 | 100 | 100 |

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1 | A | 432/451 (96%) | 426 (99%) | 6 (1%) | 62 | 49 |

All (6) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 75 | TYR |
| 1 | A | 191 | PHE |
| 1 | A | 271 | LYS |
| 1 | A | 358 | ARG |
| 1 | A | 468 | TRP |
| 1 | A | 512 | ARG |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | $\# Z > 2$ | Counts | RMSZ | $\# Z > 2$ |
| 1 | SVX | A | 194 | 1 | 10,11,12 | 2.37 | 3 (30%) | 9,14,16 | 0.95 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|-------|
| 1 | SVX | A | 194 | 1 | - | 2/9/12/14 | - |

All (3) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 1 | A | 194 | SVX | P1-C4 | -5.49 | 1.63 | 1.77 |
| 1 | A | 194 | SVX | P1-OG | 3.89 | 1.63 | 1.58 |
| 1 | A | 194 | SVX | P1-O5 | 2.87 | 1.61 | 1.58 |

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 1 | A | 194 | SVX | N-CA-CB-OG |
| 1 | A | 194 | SVX | C2-O5-P1-O6 |

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 11 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | $\# Z > 2$ | Counts | RMSZ | $\# Z > 2$ |
| 3 | ACT | A | 618 | 2 | 3,3,3 | 1.47 | 1 (33%) | 3,3,3 | 1.48 | 0 |
| 3 | ACT | A | 615 | 2 | 3,3,3 | 1.21 | 0 | 3,3,3 | 1.46 | 0 |
| 3 | ACT | A | 613 | 2 | 3,3,3 | 1.62 | 1 (33%) | 3,3,3 | 1.29 | 0 |
| 3 | ACT | A | 614 | 2 | 3,3,3 | 1.40 | 1 (33%) | 3,3,3 | 1.47 | 0 |
| 3 | ACT | A | 617 | - | 3,3,3 | 1.39 | 1 (33%) | 3,3,3 | 1.42 | 0 |
| 3 | ACT | A | 612 | 2 | 3,3,3 | 1.13 | 0 | 3,3,3 | 1.26 | 0 |
| 3 | ACT | A | 616 | - | 3,3,3 | 1.37 | 0 | 3,3,3 | 1.32 | 0 |

All (4) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 3 | A | 613 | ACT | CH3-C | 2.37 | 1.59 | 1.49 |
| 3 | A | 618 | ACT | CH3-C | 2.21 | 1.58 | 1.49 |
| 3 | A | 617 | ACT | CH3-C | 2.08 | 1.57 | 1.49 |
| 3 | A | 614 | ACT | CH3-C | 2.02 | 1.57 | 1.49 |

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 3 | A | 616 | ACT | 1 | 0 |

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|---------------|--------|---------------|-----------------------|--------|
| 1 | A | 522/547 (95%) | 0.28 | 31 (5%) 29 33 | 12, 32, 58, 86 | 6 (1%) |

All (31) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | A | 436 | ALA | 5.7 |
| 1 | A | 433 | ALA | 4.8 |
| 1 | A | 274 | LEU | 4.3 |
| 1 | A | 421 | PRO | 4.1 |
| 1 | A | 285 | VAL | 4.0 |
| 1 | A | 119 | PHE | 3.9 |
| 1 | A | 275 | ALA | 3.8 |
| 1 | A | 7 | TYR | 3.5 |
| 1 | A | 69 | ILE | 3.5 |
| 1 | A | 75 | TYR | 3.3 |
| 1 | A | 0 | GLN | 3.3 |
| 1 | A | 435 | HIS | 3.0 |
| 1 | A | 120 | LEU | 3.0 |
| 1 | A | 533 | THR | 2.8 |
| 1 | A | 273 | PRO | 2.8 |
| 1 | A | 318 | ASN | 2.7 |
| 1 | A | 324 | PHE | 2.7 |
| 1 | A | 454 | ARG | 2.5 |
| 1 | A | 72 | ASP | 2.5 |
| 1 | A | 22 | LEU | 2.4 |
| 1 | A | 1 | SER | 2.4 |
| 1 | A | 323 | ILE | 2.3 |
| 1 | A | 276 | GLY | 2.3 |
| 1 | A | 117 | ALA | 2.2 |
| 1 | A | 282 | LEU | 2.1 |
| 1 | A | 372 | ALA | 2.1 |
| 1 | A | 528 | ALA | 2.1 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | A | 420 | HIS | 2.1 |
| 1 | A | 277 | LEU | 2.0 |
| 1 | A | 335 | GLY | 2.0 |
| 1 | A | 434 | ASP | 2.0 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 1 | SVX | A | 194 | 12/13 | 0.96 | 0.08 | 22,25,37,37 | 0 |

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 3 | ACT | A | 616 | 4/4 | 0.81 | 0.16 | 55,57,58,59 | 0 |
| 2 | ZN | A | 610 | 1/1 | 0.84 | 0.15 | 88,88,88,88 | 0 |
| 3 | ACT | A | 615 | 4/4 | 0.88 | 0.13 | 46,47,56,62 | 0 |
| 2 | ZN | A | 607 | 1/1 | 0.88 | 0.09 | 65,65,65,65 | 0 |
| 3 | ACT | A | 617 | 4/4 | 0.88 | 0.13 | 47,48,56,64 | 0 |
| 3 | ACT | A | 618 | 4/4 | 0.88 | 0.11 | 40,52,56,65 | 0 |
| 3 | ACT | A | 612 | 4/4 | 0.92 | 0.09 | 29,29,29,35 | 0 |
| 3 | ACT | A | 614 | 4/4 | 0.93 | 0.10 | 33,35,40,41 | 0 |
| 2 | ZN | A | 604 | 1/1 | 0.94 | 0.10 | 61,61,61,61 | 0 |
| 3 | ACT | A | 613 | 4/4 | 0.95 | 0.10 | 29,30,34,37 | 0 |
| 2 | ZN | A | 608 | 1/1 | 0.95 | 0.06 | 77,77,77,77 | 0 |
| 2 | ZN | A | 611 | 1/1 | 0.97 | 0.09 | 56,56,56,56 | 0 |
| 2 | ZN | A | 605 | 1/1 | 0.97 | 0.11 | 55,55,55,55 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|-----------------------------|-------|
| 2 | ZN | A | 609 | 1/1 | 0.98 | 0.05 | 63,63,63,63 | 0 |
| 2 | ZN | A | 602 | 1/1 | 0.99 | 0.02 | 29,29,29,29 | 0 |
| 2 | ZN | A | 603 | 1/1 | 0.99 | 0.02 | 26,26,26,26 | 0 |
| 2 | ZN | A | 601 | 1/1 | 1.00 | 0.02 | 28,28,28,28 | 0 |
| 2 | ZN | A | 606 | 1/1 | 1.00 | 0.01 | 26,26,26,26 | 0 |

6.5 Other polymers [i](#)

There are no such residues in this entry.