



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 9, 2025 – 01:20 pm BST

PDB ID : 9H39 / pdb\_00009h39  
Title : Crystal structure of Lotus japonicus CHIP13 extracellular domain in complex with a nanobody  
Authors : Gysel, K.; Andersen, K.R.  
Deposited on : 2024-10-16  
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div><div>3%</div><div></div><div>91%</div><div>5%</div></div>
1	B	227	<div><div>%</div><div></div><div>91%</div><div>...</div></div>
2	C	128	<div><div>%</div><div></div><div>84%</div><div>11%</div><div>5%</div></div>
2	D	128	<div><div>3%</div><div></div><div>85%</div><div>9%</div><div>...</div></div>
3	N	3	<div><div></div><div>100%</div></div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	E	2	 100%
4	M	2	 100%
5	F	4	 25%  75%
6	G	3	 100%

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LysM type receptor kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	10	0
			1695	1061	270	354	10			
1	B	218	Total	C	N	O	S	0	11	0
			1725	1081	274	360	10			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	254	HIS	-	expression tag	UNP D3KU00
A	255	HIS	-	expression tag	UNP D3KU00
A	256	HIS	-	expression tag	UNP D3KU00
A	257	HIS	-	expression tag	UNP D3KU00
A	258	HIS	-	expression tag	UNP D3KU00
A	259	HIS	-	expression tag	UNP D3KU00
B	254	HIS	-	expression tag	UNP D3KU00
B	255	HIS	-	expression tag	UNP D3KU00
B	256	HIS	-	expression tag	UNP D3KU00
B	257	HIS	-	expression tag	UNP D3KU00
B	258	HIS	-	expression tag	UNP D3KU00
B	259	HIS	-	expression tag	UNP D3KU00

- Molecule 2 is a protein called anti-LYS13 VHH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	122	Total	C	N	O	S	0	8	0
			999	621	180	194	4			
2	D	123	Total	C	N	O	S	0	2	0
			979	606	179	190	4			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	N	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	M	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	G	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	0	0
			14	8	1	5		
7	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	178	Total	O	0	0
			178	178		
8	B	158	Total	O	0	0
			158	158		
8	C	167	Total	O	0	0
			167	167		
8	D	79	Total	O	0	0
			79	79		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LysM type receptor kinase





- Molecule 1: LysM type receptor kinase



- Molecule 2: anti-LYS13 VHH



- Molecule 2: anti-LYS13 VHH



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1  
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  100%

MAG1  
MAG2

- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  25% 75%

MAG1  
MAG2  
BMA3  
FUC4

- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1  
MAG2  
FUC3



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.90Å 55.09Å 109.34Å 90.00° 98.91° 90.00°	Depositor
Resolution (Å)	49.08 – 1.66 49.08 – 1.66	Depositor EDS
% Data completeness (in resolution range)	98.5 (49.08-1.66) 98.5 (49.08-1.66)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.09 (at 1.66Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.182 , 0.225 0.183 , 0.225	Depositor DCC
$R_{free}$ test set	93339 reflections (2.79%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.0	Xtriage
Anisotropy	0.436	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6190	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, FUC, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.14	0/1750	0.34	0/2393
1	B	0.13	0/1786	0.33	0/2442
2	C	0.16	0/1039	0.39	0/1408
2	D	0.13	0/1005	0.38	0/1362
All	All	0.14	0/5580	0.35	0/7605

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1695	0	1641	5	0
1	B	1725	0	1675	8	0
2	C	999	0	989	10	0
2	D	979	0	949	8	0
3	N	39	0	34	0	0
4	E	28	0	25	0	0
4	M	28	0	25	0	0
5	F	49	0	43	0	0
6	G	38	0	34	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	28	0	26	0	0
8	A	178	0	0	1	0
8	B	158	0	0	2	0
8	C	167	0	0	2	0
8	D	79	0	0	2	0
All	All	6190	0	5441	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:4:LEU:HD22	2:D:24:THR:HG22	1.66	0.78
1:A:109:MET:HG2	1:A:248:LEU:HD12	1.68	0.74
2:D:83:THR:HG23	2:D:85:GLU:H	1.58	0.67
2:C:4[A]:LEU:HD22	2:C:24[A]:THR:HG22	1.77	0.66
2:C:53:ARG:NH1	8:C:201:HOH:O	2.29	0.65
1:B:69:LYS:HG3	1:B:109:MET:HE3	1.78	0.65
2:D:33:THR:HG22	2:D:52:SER:HA	1.79	0.64
1:B:126:ASN:OD1	1:B:126:ASN:N	2.41	0.54
1:B:210:ILE:HD11	1:B:250:ILE:HG12	1.94	0.49
1:B:227:LYS:NZ	8:B:408:HOH:O	2.46	0.49
2:C:24[A]:THR:HG23	8:C:267:HOH:O	2.13	0.48
2:C:63:VAL:HG13	2:C:67:PHE:HB2	1.96	0.48
1:B:109:MET:HE1	1:B:197:LEU:HB2	1.98	0.46
1:A:70[A]:SER:OG	1:A:105:ILE:O	2.31	0.46
1:B:70[A]:SER:OG	1:B:105:ILE:O	2.31	0.46
2:D:32:ASN:ND2	2:D:104:GLU:OE2	2.43	0.45
1:B:74[A]:GLU:OE1	8:B:401:HOH:O	2.20	0.45
2:D:24:THR:HG23	8:D:243:HOH:O	2.16	0.44
2:C:13:GLN:HA	2:C:117[B]:SER:OG	2.17	0.44
2:C:30:ARG:HD2	2:C:32:ASN:OD1	2.18	0.44
2:D:33:THR:HG21	8:D:277:HOH:O	2.18	0.43
1:A:162:ASP:OD1	1:A:163:ALA:N	2.52	0.43
2:C:49:ALA:HA	2:C:58:ASN:O	2.19	0.43
2:D:12:VAL:HG11	2:D:18:LEU:HG	1.99	0.42
2:C:34:MET:HG2	2:C:94:VAL:HG22	2.02	0.42
1:A:190:PHE:CD1	1:A:239:PRO:HB2	2.55	0.42
2:D:93:LEU:HB2	2:D:108:TRP:CZ3	2.55	0.41
2:C:4[A]:LEU:CD2	2:C:24[A]:THR:HG22	2.49	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:ASP:OD2	8:A:301:HOH:O	2.22	0.41
1:B:109:MET:HG2	1:B:248:LEU:HD12	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/227 (99%)	217 (97%)	7 (3%)	0	100	100
1	B	228/227 (100%)	222 (97%)	6 (3%)	0	100	100
2	C	128/128 (100%)	125 (98%)	3 (2%)	0	100	100
2	D	123/128 (96%)	120 (98%)	3 (2%)	0	100	100
All	All	703/710 (99%)	684 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/204 (100%)	203 (99%)	2 (1%)	73	59
1	B	209/204 (102%)	207 (99%)	2 (1%)	73	59
2	C	111/109 (102%)	109 (98%)	2 (2%)	54	32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	D	106/109 (97%)	101 (95%)	5 (5%)	22 5
All	All	631/626 (101%)	620 (98%)	11 (2%)	54 35

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	MET
1	A	200	GLN
1	B	109	MET
1	B	126	ASN
2	C	63	VAL
2	C	119	HIS
2	D	4	LEU
2	D	38	ARG
2	D	54	ASP
2	D	83	THR
2	D	93	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	ASN
1	B	54	ASN
2	D	13	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PCA	A	33	1	7,8,9	0.62	0	9,10,12	1.21	0
1	PCA	B	33	1	7,8,9	0.63	0	9,10,12	1.22	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	33	1	-	0/0/11/13	0/1/1/1
1	PCA	B	33	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	33	PCA	CB-CA-N	2.02	109.11	103.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	E	1	1,4	14,14,15	0.73	0	17,19,21	0.95	1 (5%)
4	NAG	E	2	4	14,14,15	0.65	0	17,19,21	1.20	1 (5%)
5	NAG	F	1	5,1	14,14,15	0.72	0	17,19,21	1.30	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	F	2	5	14,14,15	0.75	0	17,19,21	1.06	1 (5%)
5	BMA	F	3	5	11,11,12	0.87	0	15,15,17	2.78	6 (40%)
5	FUC	F	4	5	10,10,11	0.86	0	14,14,16	0.92	0
6	NAG	G	1	6,1	14,14,15	0.73	0	17,19,21	1.29	1 (5%)
6	NAG	G	2	6	14,14,15	0.74	0	17,19,21	1.05	2 (11%)
6	FUC	G	3	6	10,10,11	0.90	1 (10%)	14,14,16	0.80	0
4	NAG	M	1	1,4	14,14,15	0.76	0	17,19,21	1.18	1 (5%)
4	NAG	M	2	4	14,14,15	0.76	0	17,19,21	0.90	1 (5%)
3	NAG	N	1	1,3	14,14,15	0.74	0	17,19,21	1.38	3 (17%)
3	NAG	N	2	3	14,14,15	0.74	0	17,19,21	1.52	4 (23%)
3	BMA	N	3	3	11,11,12	1.04	1 (9%)	15,15,17	2.19	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
5	NAG	F	1	5,1	-	0/6/23/26	0/1/1/1
5	NAG	F	2	5	-	2/6/23/26	0/1/1/1
5	BMA	F	3	5	-	0/2/19/22	0/1/1/1
5	FUC	F	4	5	-	-	0/1/1/1
6	NAG	G	1	6,1	-	0/6/23/26	0/1/1/1
6	NAG	G	2	6	-	0/6/23/26	0/1/1/1
6	FUC	G	3	6	-	-	0/1/1/1
4	NAG	M	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	M	2	4	-	2/6/23/26	0/1/1/1
3	NAG	N	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	N	2	3	-	1/6/23/26	0/1/1/1
3	BMA	N	3	3	-	2/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	N	3	BMA	C2-C3	2.15	1.55	1.52
6	G	3	FUC	O5-C1	-2.14	1.40	1.43

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	3	BMA	C1-O5-C5	8.44	123.63	112.19
3	N	3	BMA	C3-C4-C5	4.38	118.06	110.24
6	G	1	NAG	C1-O5-C5	4.27	117.98	112.19
3	N	3	BMA	C2-C3-C4	4.02	117.85	110.89
5	F	1	NAG	C1-O5-C5	3.98	117.58	112.19
3	N	3	BMA	C1-O5-C5	3.66	117.15	112.19
4	M	1	NAG	C1-O5-C5	3.37	116.76	112.19
3	N	2	NAG	O4-C4-C3	-3.34	102.62	110.35
4	E	2	NAG	C1-O5-C5	3.23	116.57	112.19
5	F	3	BMA	C3-C4-C5	3.12	115.81	110.24
3	N	1	NAG	C1-O5-C5	3.05	116.32	112.19
3	N	1	NAG	O4-C4-C3	-2.92	103.60	110.35
5	F	3	BMA	O5-C1-C2	2.66	114.88	110.77
3	N	2	NAG	C4-C3-C2	2.63	114.88	111.02
3	N	2	NAG	C2-N2-C7	2.60	126.61	122.90
3	N	1	NAG	C4-C3-C2	2.43	114.57	111.02
3	N	2	NAG	O5-C1-C2	-2.43	107.46	111.29
4	E	1	NAG	C1-O5-C5	2.37	115.40	112.19
5	F	3	BMA	C2-C3-C4	2.35	114.97	110.89
3	N	3	BMA	O3-C3-C2	-2.29	105.62	109.99
3	N	3	BMA	O4-C4-C3	-2.27	105.09	110.35
5	F	3	BMA	O5-C5-C4	2.12	115.98	110.83
4	M	2	NAG	C1-O5-C5	2.10	115.03	112.19
5	F	3	BMA	O4-C4-C3	-2.08	105.53	110.35
5	F	2	NAG	C2-N2-C7	2.05	125.82	122.90
6	G	2	NAG	C1-O5-C5	2.04	114.96	112.19
6	G	2	NAG	C2-N2-C7	2.01	125.76	122.90

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	N	3	BMA	O5-C5-C6-O6
4	M	2	NAG	O5-C5-C6-O6
5	F	2	NAG	C4-C5-C6-O6
5	F	2	NAG	O5-C5-C6-O6
3	N	3	BMA	C4-C5-C6-O6
4	M	1	NAG	O5-C5-C6-O6
4	M	2	NAG	C4-C5-C6-O6
3	N	2	NAG	C3-C2-N2-C7
3	N	1	NAG	O5-C5-C6-O6




There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




## Oligosaccharide Chain E









## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	NAG	B	302	1	14,14,15	0.73	0	17,19,21	1.17	1 (5%)
7	NAG	B	301	1	14,14,15	0.69	0	17,19,21	1.06	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	B	302	1	-	0/6/23/26	0/1/1/1
7	NAG	B	301	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	301	NAG	O5-C5-C6	2.05	110.41	107.20
7	B	302	NAG	C2-N2-C7	2.04	125.81	122.90

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	301	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/227 (94%)	0.22	6 (2%) 55 59	23, 47, 73, 99	10 (4%)
1	B	217/227 (95%)	0.32	3 (1%) 73 77	20, 50, 75, 118	11 (5%)
2	C	122/128 (95%)	0.12	1 (0%) 82 86	18, 43, 68, 101	8 (6%)
2	D	123/128 (96%)	0.63	4 (3%) 49 53	37, 58, 95, 130	2 (1%)
All	All	677/710 (95%)	0.31	14 (2%) 63 67	18, 49, 81, 130	31 (4%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	250	ILE	5.2
2	C	119	HIS	3.6
2	D	120	HIS	2.9
2	D	119	HIS	2.9
1	B	118	CYS	2.8
1	A	61	SER	2.6
2	D	83	THR	2.4
1	A	248	LEU	2.4
2	D	47	LEU	2.3
1	A	161	TYR	2.3
1	A	152	CYS	2.3
1	A	246	ALA	2.3
1	A	44	CYS	2.2
1	B	248	LEU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	PCA	B	33	8/9	0.92	0.11	41,53,62,62	0
1	PCA	A	33	8/9	0.94	0.08	34,42,49,50	0

### 6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	N	1	14/15	-	-	39,45,57,60	0
3	NAG	N	2	14/15	-	-	67,83,89,108	0
3	BMA	N	3	11/12	-	-	109,118,126,127	0
5	NAG	F	2	14/15	0.53	0.14	66,78,97,107	0
4	NAG	E	2	14/15	0.79	0.12	94,110,122,124	0
4	NAG	M	1	14/15	-	-	38,48,54,61	0
4	NAG	M	2	14/15	-	-	50,75,82,82	0
6	NAG	G	2	14/15	0.80	0.12	84,104,111,116	0
5	NAG	F	1	14/15	0.85	0.10	45,56,63,64	0
5	BMA	F	3	11/12	-	-	113,118,123,123	0
5	FUC	F	4	10/11	-	-	60,62,68,69	0
6	NAG	G	1	14/15	0.94	0.08	67,78,90,98	0
4	NAG	E	1	14/15	0.94	0.09	62,71,83,98	0
6	FUC	G	3	10/11	-	-	58,79,84,86	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain N:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain M:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	NAG	B	302	14/15	0.62	0.13	92,102,116,119	0
7	NAG	B	301	14/15	0.66	0.15	67,100,112,112	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.