



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 28, 2024 – 05:02 am GMT

PDB ID : 5HCA
Title : Globular Domain of the Entamoeba histolytica calreticulin in complex with glucose
Authors : Moreau, C.P.; Gaboriaud, C.
Deposited on : 2016-01-04
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	<div> <div>5%</div> <div>87%</div> <div>8%</div> <div>..</div> </div>
1	B	274	<div> <div>5%</div> <div>85%</div> <div>11%</div> <div>.</div> </div>
1	C	274	<div> <div>3%</div> <div>87%</div> <div>9%</div> <div>..</div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6825 atoms, of which 7 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calreticulin,Calreticulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	265	Total	C	N	O	S	8	5	0
			2176	1381	353	433	9			
1	A	266	Total	C	N	O	S	0	3	0
			2164	1371	349	436	8			
1	B	264	Total	C	N	O	S	0	6	0
			2176	1380	350	436	10			

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	12	MET	-	initiating methionine	UNP F2VN92
C	204	GLY	-	linker	UNP F2VN92
C	205	SER	-	linker	UNP F2VN92
C	206	GLY	-	linker	UNP F2VN92
C	361	GLY	-	expression tag	UNP F2VN92
C	362	HIS	-	expression tag	UNP F2VN92
C	363	HIS	-	expression tag	UNP F2VN92
C	364	HIS	-	expression tag	UNP F2VN92
C	365	HIS	-	expression tag	UNP F2VN92
C	366	HIS	-	expression tag	UNP F2VN92
C	367	HIS	-	expression tag	UNP F2VN92
A	12	MET	-	initiating methionine	UNP F2VN92
A	204	GLY	-	linker	UNP F2VN92
A	205	SER	-	linker	UNP F2VN92
A	206	GLY	-	linker	UNP F2VN92
A	361	GLY	-	expression tag	UNP F2VN92
A	362	HIS	-	expression tag	UNP F2VN92
A	363	HIS	-	expression tag	UNP F2VN92
A	364	HIS	-	expression tag	UNP F2VN92
A	365	HIS	-	expression tag	UNP F2VN92
A	366	HIS	-	expression tag	UNP F2VN92
A	367	HIS	-	expression tag	UNP F2VN92
B	12	MET	-	initiating methionine	UNP F2VN92

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	204	GLY	-	linker	UNP F2VN92
B	205	SER	-	linker	UNP F2VN92
B	206	GLY	-	linker	UNP F2VN92
B	361	GLY	-	expression tag	UNP F2VN92
B	362	HIS	-	expression tag	UNP F2VN92
B	363	HIS	-	expression tag	UNP F2VN92
B	364	HIS	-	expression tag	UNP F2VN92
B	365	HIS	-	expression tag	UNP F2VN92
B	366	HIS	-	expression tag	UNP F2VN92
B	367	HIS	-	expression tag	UNP F2VN92

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is AMMONIUM ION (three-letter code: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	H	N	0	0
			5	4	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 7 is beta-D-glucopyranose (three-letter code: BGC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	104	Total 104	O 104	0	0
8	A	77	Total 77	O 77	0	0
8	B	74	Total 74	O 74	0	0

- Molecule 1: Calreticulin, Calreticulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.36Å 143.44Å 171.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.23 – 2.15 43.23 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.2 (43.23-2.15) 95.5 (43.23-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.16Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.193 , 0.221 0.195 , 0.222	Depositor DCC
R_{free} test set	4821 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6825	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACT, NH4, BGC, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/2207	0.55	1/2971 (0.0%)
1	B	0.42	0/2219	0.56	0/2985
1	C	0.49	0/2219	0.58	1/2987 (0.0%)
All	All	0.44	0/6645	0.57	2/8943 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	LEU	CA-CB-CG	-5.22	103.29	115.30
1	C	89	LEU	CA-CB-CG	-5.04	103.72	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2164	0	2089	24	0
1	B	2176	0	2103	25	0
1	C	2176	0	2111	16	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	C	5	0	0	0	0
4	A	1	4	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	4	3	3	0	0
7	A	12	0	12	1	0
8	A	77	0	0	1	0
8	B	74	0	0	3	0
8	C	104	0	0	0	0
All	All	6818	7	6318	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 61 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:205:SER:HB2	8:B:514:HOH:O	1.78	0.82
1:A:143:TYR:HB2	1:A:195:MET:HE3	1.67	0.76
1:C:183:GLU:OE1	1:C:186:GLU:HG2	1.89	0.73
1:C:200:GLU:HA	1:B:200:GLU:OE2	1.87	0.72
1:B:193:TRP:HB3	1:B:195[A]:MET:HG3	1.73	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	267/274 (97%)	259 (97%)	7 (3%)	1 (0%)	30 27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	268/274 (98%)	257 (96%)	11 (4%)	0	100	100
1	C	268/274 (98%)	252 (94%)	15 (6%)	1 (0%)	30	27
All	All	803/822 (98%)	768 (96%)	33 (4%)	2 (0%)	37	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	290	ALA
1	A	201	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/239 (98%)	231 (98%)	4 (2%)	56	61
1	B	236/239 (99%)	232 (98%)	4 (2%)	56	61
1	C	236/239 (99%)	232 (98%)	4 (2%)	56	61
All	All	707/717 (99%)	695 (98%)	12 (2%)	56	61

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	331	ILE
1	B	25	ARG
1	B	151	ARG
1	B	48	TRP
1	C	186	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 5 are monoatomic and 1 is modelled with single atom - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	403	-	4,4,4	0.15	0	6,6,6	0.14	0
3	SO4	B	402	-	4,4,4	0.14	0	6,6,6	0.19	0
7	BGC	A	407	-	12,12,12	0.59	0	17,17,17	1.88	5 (29%)
6	ACT	A	406	-	3,3,3	0.88	0	3,3,3	0.66	0
3	SO4	C	402	-	4,4,4	0.14	0	6,6,6	0.06	0
3	SO4	B	401	-	4,4,4	0.11	0	6,6,6	0.14	0
3	SO4	A	402	-	4,4,4	0.15	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	BGC	A	407	-	-	1/2/22/22	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
7	A	407	BGC	O5-C5-C6	3.56	115.29	106.44
7	A	407	BGC	C4-C3-C2	-3.56	104.61	110.82
7	A	407	BGC	O1-C1-O5	2.82	118.83	110.38
7	A	407	BGC	C1-C2-C3	-2.69	104.74	110.31
7	A	407	BGC	O2-C2-C3	2.34	115.76	110.35

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	407	BGC	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	407	BGC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/274 (97%)	0.20	15 (5%) 31 37	21, 51, 77, 119	3 (1%)
1	B	264/274 (96%)	0.30	13 (4%) 36 41	21, 51, 77, 130	6 (2%)
1	C	265/274 (96%)	0.03	8 (3%) 52 58	19, 45, 71, 117	5 (1%)
All	All	795/822 (96%)	0.17	36 (4%) 39 44	19, 49, 75, 130	14 (1%)

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	201	ILE	10.6
1	B	201	ILE	8.0
1	B	205	SER	5.9
1	C	360	ASN	5.8
1	A	360	ASN	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	C	402	5/5	0.81	0.11	126,126,126,126	0
6	ACT	A	406	4/4	0.82	0.19	56,62,74,74	0
3	SO4	B	402	5/5	0.88	0.12	80,80,82,86	0
3	SO4	B	401	5/5	0.90	0.10	86,86,87,90	0
7	BGC	A	407	12/12	0.91	0.12	29,44,49,54	0
3	SO4	A	403	5/5	0.93	0.09	76,76,77,81	0
4	NH4	A	401	1/1	0.93	0.25	62,75,75,75	0
3	SO4	A	402	5/5	0.96	0.06	81,81,81,82	0
2	CA	B	403	1/1	0.97	0.21	66,66,66,66	0
2	CA	C	401	1/1	0.97	0.17	62,62,62,62	0
2	CA	A	404	1/1	0.97	0.15	73,73,73,73	0
5	CL	B	404	1/1	0.98	0.10	48,48,48,48	0
5	CL	A	405	1/1	0.99	0.03	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.