



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 9, 2024 – 10:30 am GMT

PDB ID : 6HGA
Title : Crystal Structure of the human IL-17RC D2-D3-D4 domains in complex with an anti-APP tag Fab
Authors : Rondeau, J.M.; Goepfert, A.
Deposited on : 2018-08-23
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	264	<div> <div>13%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>.</div> </div> </div>
2	H	222	<div> <div>97%</div> <div>.</div> </div>
3	L	219	<div> <div>2%</div> <div>96%</div> <div>.</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5551 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-17 receptor C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	263	Total	C	N	O	S	0	0	0
			2089	1328	367	381	13			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	263	GLN	ASN	engineered mutation	UNP Q8NAC3
B	307	ARG	GLN	variant	UNP Q8NAC3
B	349	GLN	ASN	engineered mutation	UNP Q8NAC3
B	372	GLN	ASN	engineered mutation	UNP Q8NAC3
B	406	GLN	ASN	engineered mutation	UNP Q8NAC3
B	468	GLU	-	expression tag	UNP Q8NAC3
B	469	PHE	-	expression tag	UNP Q8NAC3
B	470	ARG	-	expression tag	UNP Q8NAC3
B	471	HIS	-	expression tag	UNP Q8NAC3

- Molecule 2 is a protein called anti-APP-tag Fab heavy-chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	0	0
			1677	1060	274	336	7			

- Molecule 3 is a protein called anti-APP-tag Fab light-chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	219	Total	C	N	O	S	0	0	0
			1697	1058	290	342	7			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	36	Total	O	0	0
			36	36		
5	L	38	Total	O	0	0
			38	38		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-17 receptor C



- Molecule 2: anti-APP-tag Fab heavy-chain



- Molecule 3: anti-APP-tag Fab light-chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	250.91Å 250.91Å 78.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	82.13 – 2.60 82.13 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (82.13-2.60) 100.0 (82.13-2.60)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.62Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.205 , 0.232 0.214 , 0.238	Depositor DCC
R_{free} test set	2262 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	65.8	Xtriage
Anisotropy	0.443	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 86.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5551	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.50	0/2149	0.71	0/2941
2	H	0.52	0/1716	0.72	0/2352
3	L	0.50	0/1735	0.71	0/2354
All	All	0.50	0/5600	0.71	0/7647

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2089	0	2047	19	0
2	H	1677	0	1628	2	0
3	L	1697	0	1641	5	0
4	B	14	0	13	0	0
5	H	36	0	0	0	0
5	L	38	0	0	0	0
All	All	5551	0	5329	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:398:LEU:HD11	1:B:412:LEU:HB2	1.77	0.66
1:B:300:LEU:HD12	1:B:319:PRO:HD2	1.77	0.66
1:B:371:VAL:HB	1:B:379:LEU:HB2	1.77	0.66
1:B:397:LEU:HD12	1:B:439:LEU:HD13	1.77	0.66
3:L:88:LEU:HD11	3:L:111:LEU:HG	1.90	0.54
1:B:401:ARG:HH21	1:B:454:ALA:HB2	1.72	0.53
1:B:235:LEU:HD12	1:B:275:VAL:HG12	1.93	0.50
1:B:471:HIS:CE1	3:L:96:ASN:OD1	2.67	0.48
1:B:401:ARG:HB2	1:B:452:LEU:HA	1.96	0.48
3:L:189:ASP:O	3:L:193:ARG:HG3	2.14	0.48
2:H:190:THR:HG22	2:H:192:SER:H	1.81	0.46
1:B:218:GLY:C	1:B:220:ASN:H	2.19	0.45
3:L:141:LEU:HD21	3:L:151:VAL:HG22	1.98	0.45
1:B:314:TRP:HE3	1:B:356:LEU:HD23	1.81	0.45
1:B:297:HIS:CD2	1:B:322:LEU:HD11	2.52	0.45
1:B:306:LEU:HB3	1:B:381:GLU:HG3	1.99	0.45
1:B:351:THR:OG1	1:B:354:LYS:HB2	2.17	0.45
1:B:385:ALA:O	1:B:389:GLY:HA3	2.18	0.44
1:B:325:GLU:HB3	1:B:347:TRP:HB3	2.01	0.43
1:B:435:LEU:HD21	1:B:460:MET:SD	2.59	0.43
2:H:41:GLN:HB2	2:H:47:LEU:HD23	2.01	0.42
1:B:341:LEU:O	1:B:345:LEU:HB2	2.19	0.41
1:B:256:GLY:HA2	1:B:258:GLN:HG2	2.03	0.41
1:B:306:LEU:HD11	1:B:314:TRP:HB3	2.03	0.41
3:L:15:LEU:HG	3:L:111:LEU:HD21	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	261/264 (99%)	245 (94%)	15 (6%)	1 (0%)	30	52
2	H	220/222 (99%)	216 (98%)	3 (1%)	1 (0%)	25	47
3	L	217/219 (99%)	214 (99%)	3 (1%)	0	100	100
All	All	698/705 (99%)	675 (97%)	21 (3%)	2 (0%)	37	59

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	178	SER
1	B	219	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	234/234 (100%)	226 (97%)	8 (3%)	32	58
2	H	194/194 (100%)	193 (100%)	1 (0%)	86	95
3	L	197/197 (100%)	197 (100%)	0	100	100
All	All	625/625 (100%)	616 (99%)	9 (1%)	62	82

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	235	LEU
1	B	236	SER
1	B	258	GLN
1	B	399	GLU
1	B	409	LEU
1	B	412	LEU
1	B	427	ARG
1	B	464	ILE
2	H	179	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	222	HIS
1	B	241	GLN
3	L	18	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	H	1	2	7,8,9	0.79	0	9,10,12	2.29	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	PCA	CB-CA-C	-6.07	104.35	112.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	501	1	14,14,15	0.34	0	17,19,21	0.94	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	501	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	NAG	C1-O5-C5	2.83	116.03	112.19
4	B	501	NAG	O5-C1-C2	-2.31	107.63	111.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	263/264 (99%)	0.91	35 (13%) 8 6	70, 111, 151, 192	0
2	H	221/222 (99%)	-0.10	1 (0%) 87 84	50, 65, 90, 121	0
3	L	219/219 (100%)	0.04	4 (1%) 67 62	56, 72, 96, 122	0
All	All	703/705 (99%)	0.32	40 (5%) 30 25	50, 82, 137, 192	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	209	LEU	7.3
1	B	424	ALA	5.1
1	B	403	PRO	4.0
1	B	452	LEU	3.8
1	B	354	LYS	3.6
1	B	210	PRO	3.4
1	B	259	ILE	3.4
1	B	214	VAL	3.3
1	B	400	THR	3.2
1	B	224	VAL	3.2
1	B	314	TRP	3.2
1	B	246	PRO	3.0
1	B	414	PRO	3.0
1	B	369	VAL	2.8
1	B	379	LEU	2.6
3	L	24	ARG	2.5
1	B	308	LEU	2.5
1	B	407	ARG	2.5
1	B	358	PHE	2.5
1	B	355	VAL	2.5
2	H	137	THR	2.5
1	B	245	PRO	2.4
1	B	359	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	409	LEU	2.4
1	B	402	GLY	2.4
3	L	65	ASP	2.4
1	B	309	LEU	2.3
1	B	311	LEU	2.3
1	B	301	TRP	2.2
1	B	352	VAL	2.2
1	B	216	ALA	2.2
1	B	321	SER	2.1
1	B	326	ALA	2.1
1	B	226	ASN	2.1
1	B	303	ALA	2.1
1	B	406	GLN	2.1
1	B	418	THR	2.0
3	L	165	LEU	2.0
3	L	82	ARG	2.0
1	B	242	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	H	1	8/9	0.86	0.14	100,101,107,110	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	B	501	14/15	0.79	0.15	154,157,160,163	0

6.5 Other polymers [i](#)

There are no such residues in this entry.