



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 01:17 PM EDT

PDB ID : 6HHV
Title : Structure of T. thermophilus AspRS in Complex with 5'-O-(N-(L-aspartyl)-sulfamoyl)N3-methyluridine
Authors : De Graef, S.; Pang, L.; Strelkov, S.V.; Weeks, S.D.
Deposited on : 2018-08-29
Resolution : 2.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	581	<div> <div>5%</div> <div>92%</div> <div>6% ..</div> </div>
1	B	581	<div> <div>15%</div> <div>92%</div> <div>8%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9212 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aspartate-tRNA(Asp) ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	574	Total	C	N	O	S	0	0	0
			4536	2903	813	809	11			
1	B	580	Total	C	N	O	S	0	0	0
			4455	2848	796	800	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P36419
B	0	GLY	-	expression tag	UNP P36419

- Molecule 2 is 5'-O-(N-(L-aspartyl)-sulfamoyl)N3-methyluridine (three-letter code: G5Q) (formula: C₁₄H₂₀N₄O₁₁S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			30	14	4	11	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	S	0	0
			30	14	4	11	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	91	Total	O	0	0
			91	91		
3	B	70	Total	O	0	0
			70	70		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aspartate-tRNA(Asp) ligase



• Molecule 1: Aspartate-tRNA(Asp) ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.22Å 112.83Å 88.53Å 90.00° 104.38° 90.00°	Depositor
Resolution (Å)	67.28 – 2.18 67.28 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (67.28-2.18) 99.8 (67.28-2.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.192 , 0.226 0.196 , 0.230	Depositor DCC
R_{free} test set	2325 reflections (2.56%)	wwPDB-VP
Wilson B-factor (Å ²)	50.6	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9212	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: G5Q

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/4644	0.48	0/6294
1	B	0.28	0/4564	0.49	0/6205
All	All	0.29	0/9208	0.49	0/12499

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4536	0	4511	21	0
1	B	4455	0	4300	25	0
2	A	30	0	0	0	0
2	B	30	0	0	0	0
3	A	91	0	0	0	0
3	B	70	0	0	1	0
All	All	9212	0	8811	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:ARG:HH12	1:B:139:ARG:HH21	1.42	0.68
1:B:408:ALA:HA	1:B:413:LEU:HD13	1.79	0.63
1:A:349:LEU:HD11	1:A:387:LEU:HB3	1.83	0.59
1:A:332:LYS:O	1:A:336:GLU:HG3	2.03	0.59
1:A:154:ILE:HD13	1:A:238:LEU:HD22	1.88	0.56
1:B:409:ASP:OD1	1:B:415:ARG:NH2	2.40	0.55
1:B:430:GLU:HB2	1:B:441:MET:HG2	1.89	0.53
1:A:553:GLU:OE1	1:A:553:GLU:N	2.42	0.52
1:B:138:ARG:NH1	1:B:142:GLU:OE1	2.43	0.51
1:B:326:PRO:HA	1:B:384:ASP:OD2	2.10	0.50
1:B:78:ARG:HD2	1:B:90:VAL:O	2.10	0.50
1:B:204:LYS:HD2	1:B:241:GLU:HB2	1.94	0.50
1:B:351:TRP:HA	1:B:386:LEU:O	2.12	0.50
1:B:172:THR:OG1	1:B:173:LYS:N	2.42	0.49
1:B:306:PHE:O	1:B:315:GLN:HG2	2.12	0.49
1:A:40:ARG:HD2	1:A:114:TRP:CD1	2.49	0.48
1:A:297:LEU:HB3	1:A:323:LEU:HD11	1.95	0.47
1:A:207:LEU:HD22	1:A:212:LEU:HD12	1.96	0.47
1:B:25:VAL:HG21	1:B:63:VAL:CG1	2.45	0.47
1:A:284:GLU:OE1	1:A:321:LYS:NZ	2.48	0.47
1:A:389:VAL:HG23	1:A:400:LEU:HD13	1.96	0.46
1:B:226:ASP:O	1:B:228:ARG:N	2.38	0.46
1:A:488:ARG:NH2	1:A:516:GLU:OE1	2.47	0.46
1:B:443:HIS:HB2	1:B:444:PRO:HD2	2.00	0.44
1:A:74:LEU:HD11	1:A:76:ARG:HD3	2.00	0.44
1:B:139:ARG:NH1	3:B:701:HOH:O	2.30	0.44
1:A:139:ARG:O	1:A:139:ARG:HD3	2.17	0.44
1:A:165:GLN:HA	1:A:216:PHE:O	2.18	0.44
1:A:343:ARG:HD3	1:A:343:ARG:HA	1.61	0.43
1:B:180:ARG:HB2	1:B:224:ASP:HB3	2.00	0.43
1:A:326:PRO:HA	1:A:384:ASP:OD1	2.19	0.42
1:A:174:SER:HB2	1:A:195:ALA:HB2	2.01	0.42
1:B:202:LEU:HD21	1:B:445:PHE:CZ	2.53	0.42
1:A:139:ARG:HD3	1:A:139:ARG:C	2.40	0.42
1:B:137:ARG:NH1	1:B:139:ARG:HH21	2.11	0.42
1:A:329:LEU:HD23	1:A:333:GLU:HB3	2.03	0.41
1:A:572:GLU:OE2	1:B:180:ARG:NH1	2.53	0.41
1:A:204:LYS:HD3	1:A:239:ASP:OD1	2.21	0.41
1:B:40:ARG:HD2	1:B:114:TRP:CD1	2.56	0.41
1:B:440:TYR:HD2	1:B:497:LEU:HD22	1.86	0.41
1:B:158:LEU:HD23	1:B:158:LEU:HA	1.93	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:277:LEU:O	1:B:423:VAL:HA	2.20	0.40
1:A:74:LEU:HD11	1:A:76:ARG:NH1	2.36	0.40
1:B:25:VAL:HG21	1:B:63:VAL:HG11	2.02	0.40
1:B:279:TYR:CE2	1:B:283:MET:HE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	570/581 (98%)	556 (98%)	14 (2%)	0	100	100
1	B	576/581 (99%)	558 (97%)	17 (3%)	1 (0%)	47	52
All	All	1146/1162 (99%)	1114 (97%)	31 (3%)	1 (0%)	51	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	227	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	461/483 (95%)	454 (98%)	7 (2%)	65	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	434/483 (90%)	429 (99%)	5 (1%)	71	81
All	All	895/966 (93%)	883 (99%)	12 (1%)	69	79

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLU
1	A	72	LYS
1	A	74	LEU
1	A	139	ARG
1	A	215	TYR
1	A	343	ARG
1	A	421	LEU
1	B	2	ARG
1	B	76	ARG
1	B	174	SER
1	B	215	TYR
1	B	231	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	569	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	G5Q	B	601	-	31,31,31	0.64	1 (3%)	42,46,46	0.80	2 (4%)
2	G5Q	A	601	-	31,31,31	0.68	1 (3%)	42,46,46	0.80	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G5Q	B	601	-	-	2/22/39/39	0/2/2/2
2	G5Q	A	601	-	-	1/22/39/39	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	G5Q	C-N3S	2.91	1.42	1.37
2	B	601	G5Q	C-N3S	2.72	1.42	1.37

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	G5Q	O-C-CA	2.88	126.35	120.28
2	B	601	G5Q	O-C-N3S	-2.72	118.05	122.98
2	A	601	G5Q	O-C-N3S	-2.54	118.38	122.98
2	A	601	G5Q	O-C-CA	2.18	124.86	120.28

There are no chirality outliers.


All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	G5Q	N-CA-CD-CG
2	B	601	G5Q	N-CA-CD-CG
2	B	601	G5Q	C5'-O5'-S-N3S

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	574/581 (98%)	0.44	28 (4%) 29 31	39, 59, 94, 119	0
1	B	580/581 (99%)	0.89	90 (15%) 2 2	41, 61, 125, 147	0
All	All	1154/1162 (99%)	0.67	118 (10%) 6 7	39, 60, 114, 147	0

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	360	SER	8.9
1	B	363	VAL	7.6
1	B	303	GLY	7.5
1	B	365	LYS	7.3
1	B	382	PRO	7.3
1	B	304	PRO	7.2
1	B	306	PHE	6.6
1	B	357	GLY	6.6
1	B	367	LEU	6.5
1	B	373	ALA	6.5
1	B	386	LEU	6.3
1	B	361	GLY	6.2
1	B	387	LEU	6.1
1	B	362	GLY	6.1
1	B	374	LEU	6.0
1	B	380	ALA	5.9
1	B	326	PRO	5.9
1	B	317	ALA	5.8
1	B	322	ALA	5.8
1	B	351	TRP	5.6
1	B	359	PHE	5.5
1	B	328	ALA	5.4
1	B	354	VAL	5.3
1	B	308	GLN	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	352	ALA	5.1
1	B	227	LEU	4.8
1	B	366	PHE	4.8
1	B	335	ALA	4.7
1	B	305	LEU	4.7
1	B	388	PHE	4.7
1	B	302	VAL	4.5
1	B	313	VAL	4.4
1	B	385	THR	4.4
1	B	352	ALA	4.4
1	B	355	GLU	4.4
1	B	311	PHE	4.3
1	B	369	PRO	4.3
1	B	358	GLY	4.3
1	B	412	GLY	4.2
1	B	314	PHE	4.1
1	B	325	LEU	4.1
1	B	370	VAL	4.1
1	B	307	ARG	4.1
1	B	356	GLU	4.1
1	A	359	PHE	4.0
1	B	389	VAL	4.0
1	B	348	GLY	3.9
1	B	375	LEU	3.9
1	B	411	LEU	3.9
1	A	356	GLU	3.8
1	B	323	LEU	3.8
1	B	364	ALA	3.8
1	B	378	THR	3.7
1	B	324	ALA	3.6
1	B	320	VAL	3.5
1	B	399	ALA	3.5
1	B	349	LEU	3.5
1	B	350	ALA	3.5
1	B	377	ALA	3.5
1	B	329	LEU	3.5
1	B	229	ALA	3.5
1	B	332	LYS	3.5
1	B	384	ASP	3.4
1	B	390	ALA	3.4
1	B	376	GLN	3.4
1	B	301	GLU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	312	ARG	3.3
1	B	309	SER	3.2
1	A	373	ALA	3.2
1	A	226	ASP	3.1
1	B	400	LEU	3.1
1	A	442	HIS	3.0
1	A	375	LEU	3.0
1	A	371	ARG	2.9
1	A	383	GLY	2.9
1	B	413	LEU	2.9
1	B	336	GLU	2.8
1	B	500	GLY	2.8
1	B	327	LYS	2.8
1	B	319	SER	2.7
1	A	437	ALA	2.7
1	B	431	TRP	2.7
1	B	321	LYS	2.7
1	B	368	GLU	2.6
1	B	383	GLY	2.6
1	B	440	TYR	2.6
1	B	379	GLU	2.5
1	B	331	ARG	2.5
1	A	326	PRO	2.5
1	B	316	GLU	2.5
1	B	346	ALA	2.5
1	A	440	TYR	2.5
1	A	325	LEU	2.4
1	B	580	PRO	2.4
1	A	357	GLY	2.4
1	A	351	TRP	2.4
1	B	406	ARG	2.4
1	B	297	LEU	2.4
1	B	437	ALA	2.3
1	A	312	ARG	2.3
1	B	462	ARG	2.3
1	A	360	SER	2.3
1	B	176	PRO	2.3
1	A	379	GLU	2.3
1	B	166	VAL	2.2
1	A	354	VAL	2.2
1	B	410	LEU	2.2
1	A	31	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	74	LEU	2.1
1	A	367	LEU	2.1
1	A	382	PRO	2.1
1	A	332	LYS	2.1
1	A	385	THR	2.1
1	A	377	ALA	2.0
1	B	230	ASP	2.0
1	B	439	THR	2.0
1	A	430	GLU	2.0
1	A	386	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	G5Q	B	601	30/30	0.97	0.13	40,51,56,58	0
2	G5Q	A	601	30/30	0.98	0.15	43,48,58,59	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around G5Q B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around G5Q A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.