



## wwPDB EM Validation Summary Report ⓘ

Jul 9, 2025 – 12:10 PM JST

PDB ID : 8HMZ / pdb\_00008hmz  
EMDB ID : EMD-34905  
Title : Cryo-EM structure of the human post-catalytic TSEN/pre-tRNA complex  
Authors : Zhang, X.; Yang, F.; Zhan, X.; Shi, Y.  
Deposited on : 2022-12-06  
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4-5-2 with Phenix2.0rc1  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : **FAILED**  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	485	
2	B	330	
3	C	546	
4	D	213	
5	5	72	
6	3	42	
7	E	445	

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA-splicing endonuclease subunit Sen2.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	313	Total	C	N	O	S	0	0
			2602	1680	443	466	13		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP Q8NCE0
A	-18	ALA	-	expression tag	UNP Q8NCE0
A	-17	SER	-	expression tag	UNP Q8NCE0
A	-16	ASP	-	expression tag	UNP Q8NCE0
A	-15	TYR	-	expression tag	UNP Q8NCE0
A	-14	LYS	-	expression tag	UNP Q8NCE0
A	-13	ASP	-	expression tag	UNP Q8NCE0
A	-12	ASP	-	expression tag	UNP Q8NCE0
A	-11	ASP	-	expression tag	UNP Q8NCE0
A	-10	ASP	-	expression tag	UNP Q8NCE0
A	-9	LYS	-	expression tag	UNP Q8NCE0
A	-8	ALA	-	expression tag	UNP Q8NCE0
A	-7	SER	-	expression tag	UNP Q8NCE0
A	-6	ASP	-	expression tag	UNP Q8NCE0
A	-5	GLU	-	expression tag	UNP Q8NCE0
A	-4	VAL	-	expression tag	UNP Q8NCE0
A	-3	ASP	-	expression tag	UNP Q8NCE0
A	-2	ALA	-	expression tag	UNP Q8NCE0
A	-1	GLY	-	expression tag	UNP Q8NCE0
A	0	THR	-	expression tag	UNP Q8NCE0

- Molecule 2 is a protein called tRNA-splicing endonuclease subunit Sen34.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	264	Total	C	N	O	S	0	0
			2062	1301	388	369	4		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	initiating methionine	UNP Q9BSV6
B	-18	ALA	-	expression tag	UNP Q9BSV6
B	-17	SER	-	expression tag	UNP Q9BSV6
B	-16	ASP	-	expression tag	UNP Q9BSV6
B	-15	TYR	-	expression tag	UNP Q9BSV6
B	-14	LYS	-	expression tag	UNP Q9BSV6
B	-13	ASP	-	expression tag	UNP Q9BSV6
B	-12	ASP	-	expression tag	UNP Q9BSV6
B	-11	ASP	-	expression tag	UNP Q9BSV6
B	-10	ASP	-	expression tag	UNP Q9BSV6
B	-9	LYS	-	expression tag	UNP Q9BSV6
B	-8	ALA	-	expression tag	UNP Q9BSV6
B	-7	SER	-	expression tag	UNP Q9BSV6
B	-6	ASP	-	expression tag	UNP Q9BSV6
B	-5	GLU	-	expression tag	UNP Q9BSV6
B	-4	VAL	-	expression tag	UNP Q9BSV6
B	-3	ASP	-	expression tag	UNP Q9BSV6
B	-2	ALA	-	expression tag	UNP Q9BSV6
B	-1	GLY	-	expression tag	UNP Q9BSV6
B	0	THR	-	expression tag	UNP Q9BSV6

- Molecule 3 is a protein called tRNA-splicing endonuclease subunit Sen54.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	285	Total	C	N	O	S	0	0
			2263	1440	405	412	6		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-19	MET	-	initiating methionine	UNP Q7Z6J9
C	-18	ALA	-	expression tag	UNP Q7Z6J9
C	-17	SER	-	expression tag	UNP Q7Z6J9
C	-16	ASP	-	expression tag	UNP Q7Z6J9
C	-15	TYR	-	expression tag	UNP Q7Z6J9
C	-14	LYS	-	expression tag	UNP Q7Z6J9
C	-13	ASP	-	expression tag	UNP Q7Z6J9
C	-12	ASP	-	expression tag	UNP Q7Z6J9
C	-11	ASP	-	expression tag	UNP Q7Z6J9
C	-10	ASP	-	expression tag	UNP Q7Z6J9
C	-9	LYS	-	expression tag	UNP Q7Z6J9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-8	ALA	-	expression tag	UNP Q7Z6J9
C	-7	SER	-	expression tag	UNP Q7Z6J9
C	-6	ASP	-	expression tag	UNP Q7Z6J9
C	-5	GLU	-	expression tag	UNP Q7Z6J9
C	-4	VAL	-	expression tag	UNP Q7Z6J9
C	-3	ASP	-	expression tag	UNP Q7Z6J9
C	-2	ALA	-	expression tag	UNP Q7Z6J9
C	-1	GLY	-	expression tag	UNP Q7Z6J9
C	0	THR	-	expression tag	UNP Q7Z6J9

- Molecule 4 is a protein called Chromosome 1 open reading frame 19, isoform CRA\_a.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	126	Total	C	N	O	S	0	0
			984	631	154	191	8		

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-37	MET	-	initiating methionine	UNP A0A2U3TZM3
D	-36	ALA	-	expression tag	UNP A0A2U3TZM3
D	-35	SER	-	expression tag	UNP A0A2U3TZM3
D	-34	SER	-	expression tag	UNP A0A2U3TZM3
D	-33	ALA	-	expression tag	UNP A0A2U3TZM3
D	-32	TRP	-	expression tag	UNP A0A2U3TZM3
D	-31	SER	-	expression tag	UNP A0A2U3TZM3
D	-30	HIS	-	expression tag	UNP A0A2U3TZM3
D	-29	PRO	-	expression tag	UNP A0A2U3TZM3
D	-28	GLN	-	expression tag	UNP A0A2U3TZM3
D	-27	PHE	-	expression tag	UNP A0A2U3TZM3
D	-26	GLU	-	expression tag	UNP A0A2U3TZM3
D	-25	LYS	-	expression tag	UNP A0A2U3TZM3
D	-24	GLY	-	expression tag	UNP A0A2U3TZM3
D	-23	GLY	-	expression tag	UNP A0A2U3TZM3
D	-22	GLY	-	expression tag	UNP A0A2U3TZM3
D	-21	SER	-	expression tag	UNP A0A2U3TZM3
D	-20	GLY	-	expression tag	UNP A0A2U3TZM3
D	-19	GLY	-	expression tag	UNP A0A2U3TZM3
D	-18	GLY	-	expression tag	UNP A0A2U3TZM3
D	-17	SER	-	expression tag	UNP A0A2U3TZM3
D	-16	GLY	-	expression tag	UNP A0A2U3TZM3
D	-15	GLY	-	expression tag	UNP A0A2U3TZM3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-14	SER	-	expression tag	UNP A0A2U3TZM3
D	-13	ALA	-	expression tag	UNP A0A2U3TZM3
D	-12	TRP	-	expression tag	UNP A0A2U3TZM3
D	-11	SER	-	expression tag	UNP A0A2U3TZM3
D	-10	HIS	-	expression tag	UNP A0A2U3TZM3
D	-9	PRO	-	expression tag	UNP A0A2U3TZM3
D	-8	GLN	-	expression tag	UNP A0A2U3TZM3
D	-7	PHE	-	expression tag	UNP A0A2U3TZM3
D	-6	GLU	-	expression tag	UNP A0A2U3TZM3
D	-5	LYS	-	expression tag	UNP A0A2U3TZM3
D	-4	GLY	-	expression tag	UNP A0A2U3TZM3
D	-3	SER	-	expression tag	UNP A0A2U3TZM3
D	-2	ALA	-	expression tag	UNP A0A2U3TZM3
D	-1	ALA	-	expression tag	UNP A0A2U3TZM3
D	0	ALA	-	expression tag	UNP A0A2U3TZM3

- Molecule 5 is a RNA chain called Pre-tRNA 5' END.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	5	45	Total	C	N	O	P	0	0
			968	430	174	318	46		

- Molecule 6 is a RNA chain called Pre-tRNA 3' END.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	3	37	Total	C	N	O	P	0	0
			788	353	142	257	36		

- Molecule 7 is a protein called Polyribonucleotide 5'-hydroxyl-kinase Clp1.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	E	379	Total	C	N	O	0	0
			1869	1111	379	379		

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-19	MET	-	initiating methionine	UNP Q92989
E	-18	ALA	-	expression tag	UNP Q92989
E	-17	SER	-	expression tag	UNP Q92989
E	-16	ASP	-	expression tag	UNP Q92989
E	-15	TYR	-	expression tag	UNP Q92989

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	-14	LYS	-	expression tag	UNP Q92989
E	-13	ASP	-	expression tag	UNP Q92989
E	-12	ASP	-	expression tag	UNP Q92989
E	-11	ASP	-	expression tag	UNP Q92989
E	-10	ASP	-	expression tag	UNP Q92989
E	-9	LYS	-	expression tag	UNP Q92989
E	-8	ALA	-	expression tag	UNP Q92989
E	-7	SER	-	expression tag	UNP Q92989
E	-6	ASP	-	expression tag	UNP Q92989
E	-5	GLU	-	expression tag	UNP Q92989
E	-4	VAL	-	expression tag	UNP Q92989
E	-3	ASP	-	expression tag	UNP Q92989
E	-2	ALA	-	expression tag	UNP Q92989
E	-1	GLY	-	expression tag	UNP Q92989
E	0	THR	-	expression tag	UNP Q92989


- Molecule 8 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
8	5	3	Total Mg 3 3	0
8	3	3	Total Mg 3 3	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tRNA-splicing endonuclease subunit Sen2




- Molecule 2: tRNA-splicing endonuclease subunit Sen34




- Molecule 3: tRNA-splicing endonuclease subunit Sen54








- Molecule 4: Chromosome 1 open reading frame 19, isoform CRA\_a




- Molecule 5: Pre-tRNA 5' END




- Molecule 6: Pre-tRNA 3' END



- Molecule 7: Polyribonucleotide 5'-hydroxyl-kinase Clp1





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	309195	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50.00	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A23, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.18	0/2665	0.47	0/3596
2	B	0.18	0/2108	0.43	0/2856
3	C	0.37	0/2319	0.85	23/3140 (0.7%)
4	D	0.17	0/1004	0.42	0/1366
5	5	0.41	2/1053 (0.2%)	0.43	2/1638 (0.1%)
6	3	0.14	0/881	0.23	0/1372
7	E	0.24	0/1866	0.65	1/2592 (0.0%)
All	All	0.26	2/11896 (0.0%)	0.57	26/16560 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	5	51	C	P-OP2	-7.96	1.33	1.49
5	5	51	C	P-OP1	-5.85	1.37	1.49

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	412	GLY	CA-C-N	10.44	141.48	121.54
3	C	412	GLY	C-N-CA	10.44	141.48	121.54
5	5	51	C	O4'-C1'-C2'	-9.37	96.43	105.80
3	C	414	ALA	O-C-N	8.34	133.69	122.59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	420	VAL	CA-C-N	8.32	137.43	121.54

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	412	GLY	Mainchain
3	C	414	ALA	Mainchain
3	C	521	PRO	Mainchain
3	C	522	GLN	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2602	0	2626	54	0
2	B	2062	0	2091	34	0
3	C	2263	0	2234	44	0
4	D	984	0	987	23	0
5	5	968	0	484	17	0
6	3	788	0	400	7	0
7	E	1869	0	827	20	0
8	3	3	0	0	0	0
8	5	3	0	0	0	0
All	All	11542	0	9649	171	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 171 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:E:17:LEU:O	7:E:70:GLY:HA2	1.77	0.84
5:5:36:U:O2	5:5:47:A:N6	2.10	0.84
1:A:311:LEU:HD21	3:C:414:ALA:HB2	1.61	0.81

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23:LEU:HB2	1:A:26:PRO:HD2	1.67	0.77
7:E:25:PHE:HA	7:E:81:VAL:O	1.85	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/485 (63%)	276 (90%)	29 (9%)	2 (1%)	19	49
2	B	260/330 (79%)	250 (96%)	10 (4%)	0	100	100
3	C	281/546 (52%)	259 (92%)	19 (7%)	3 (1%)	12	37
4	D	124/213 (58%)	118 (95%)	5 (4%)	1 (1%)	16	45
7	E	373/445 (84%)	346 (93%)	27 (7%)	0	100	100
All	All	1345/2019 (67%)	1249 (93%)	90 (7%)	6 (0%)	32	60

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	17	GLU
3	C	413	GLN
3	C	421	LEU
3	C	524	VAL
4	D	162	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/428 (67%)	269 (93%)	19 (7%)	14	39
2	B	212/261 (81%)	201 (95%)	11 (5%)	19	50
3	C	242/464 (52%)	227 (94%)	15 (6%)	15	43
4	D	113/171 (66%)	107 (95%)	6 (5%)	19	49
All	All	855/1324 (65%)	804 (94%)	51 (6%)	18	44

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	234	LEU
3	C	425	SER
4	D	111	THR
2	B	268	THR
3	C	413	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	522	GLN
3	C	507	HIS
2	B	205	GLN
3	C	454	GLN
2	B	186	GLN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	5	43/72 (59%)	17 (39%)	0
6	3	36/42 (85%)	1 (2%)	0
All	All	79/114 (69%)	18 (22%)	0

5 of 18 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	5	16	U
5	5	17	G
5	5	18	G

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
5	5	19	A
5	5	20	U

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	A23	5	52	5	19,28,29	1.41	3 (15%)	19,43,46	1.85	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	A23	5	52	5	-	0/3/35/36	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	5	52	A23	O5'-C5'	-2.85	1.37	1.44
5	5	52	A23	C5'-C4'	-2.81	1.42	1.51
5	5	52	A23	C5-C4	2.50	1.47	1.40

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	5	52	A23	O2C-PC-O1C	3.52	121.25	109.89
5	5	52	A23	N3-C2-N1	-3.14	123.78	128.68

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	5	52	A23	C5'-C4'-C3'	2.89	123.96	114.40
5	5	52	A23	C4-C5-N7	-2.67	106.62	109.40
5	5	52	A23	O3'-PC-O1C	-2.46	109.25	115.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	5	52	A23	3	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.