



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 02:38 AM EDT

PDB ID : 5HQ2  
Title : Structural model of Set8 histone H4 Lys20 methyltransferase bound to nucleosome core particle  
Authors : Tavarekere, G.; McGinty, R.K.; Tan, S.  
Deposited on : 2016-01-21  
Resolution : 4.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 4.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1055 (5.20-3.80)
Clashscore	141614	1123 (5.20-3.80)
Ramachandran outliers	138981	1069 (5.20-3.80)
Sidechain outliers	138945	1050 (5.20-3.80)
RSRZ outliers	127900	1101 (5.30-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	 64% 10% 27%
2	B	102	 75% 24%
3	G	129	 75% 6% 19%
4	H	122	 70% 7% 22%
5	I	149	 48% 51%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	J	149	<div><div></div><div>46%</div><div></div><div>52%</div></div>
7	K	483	<div><div></div><div>82%</div><div>5%</div><div>13%</div></div>
8	M	202	<div><div></div><div>26%</div><div>78%</div><div>21%</div></div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7738 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	99	Total	C	N	O	0	0	0
			495	297	99	99			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	conflict	UNP P84233

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	78	Total	C	N	O	0	0	0
			384	228	78	78			

- Molecule 3 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	105	Total	C	N	O	0	0	0
			516	306	105	105			

- Molecule 4 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	95	Total	C	N	O	0	0	0
			473	283	95	95			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	29	THR	SER	conflict	UNP P02281

- Molecule 5 is a DNA chain called DNA (149-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	73	Total	C	N	O	P	0	0	0
			1487	707	271	436	73			

- Molecule 6 is a DNA chain called DNA (149-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	J	72	Total	C	N	O	P	0	0	0
			1484	704	274	434	72			

- Molecule 7 is a protein called Guanine nucleotide exchange factor SRM1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	K	422	Total	C	N	O	0	0	0
			2112	1268	422	422			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	0	GLY	-	expression tag	UNP P21827
K	1	SER	-	expression tag	UNP P21827


- Molecule 8 is a protein called N-lysine methyltransferase SETD8.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	M	160	Total	C	N	O	0	0	0
			787	467	160	160			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	151	GLY	-	expression tag	UNP Q9NQR1
M	152	SER	-	expression tag	UNP Q9NQR1
M	347	PHE	HIS	conflict	UNP Q9NQR1





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.73Å 300.75Å 182.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.00 – 4.50 44.07 – 4.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.00-4.50) 99.5 (44.07-4.50)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 4.45Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.339 , 0.397 0.345 , 0.397	Depositor DCC
$R_{free}$ test set	845 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	213.1	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.15 , 301.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	7738	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	221.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/495	0.60	0/691
2	B	0.47	0/384	0.71	0/532
3	G	0.44	0/515	0.58	0/714
4	H	0.45	0/473	0.68	1/659 (0.2%)
5	I	0.30	0/1665	0.73	0/2562
6	J	0.33	1/1664 (0.1%)	0.71	0/2566
7	K	0.37	0/2130	0.56	0/2968
8	M	0.35	0/786	0.50	0/1091
All	All	0.37	1/8112 (0.0%)	0.64	1/11783 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	J	-71	DA	C1'-N9	-5.27	1.39	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	101	GLY	N-CA-C	6.40	129.09	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	495	0	234	9	0
2	B	384	0	186	3	0
3	G	516	0	249	4	0
4	H	473	0	227	6	0
5	I	1487	0	821	4	0
6	J	1484	0	812	3	0
7	K	2112	0	1078	17	0
8	M	787	0	362	0	0
All	All	7738	0	3969	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:I:70:DC:H2''	5:I:71:DT:H71	1.52	0.89
3:G:42:ARG:CB	4:H:85:THR:CB	2.59	0.80
5:I:71:DT:O4	6:J:-71:DA:N1	2.25	0.70
5:I:70:DC:H2''	5:I:71:DT:C7	2.22	0.68
7:K:266:PRO:HA	7:K:267:ARG:CB	2.27	0.64
7:K:266:PRO:CA	7:K:267:ARG:CB	2.75	0.63
1:A:42:ARG:N	5:I:70:DC:OP1	2.33	0.62
7:K:266:PRO:HB3	7:K:267:ARG:CB	2.31	0.61
4:H:94:ALA:O	4:H:98:LEU:N	2.35	0.60
7:K:230:LYS:CB	7:K:281:GLU:O	2.50	0.60
3:G:93:LEU:O	3:G:96:LEU:N	2.35	0.59
1:A:88:ALA:HB2	2:B:83:ALA:HA	1.84	0.58
7:K:268:PRO:HB2	7:K:269:PHE:CB	2.33	0.58
7:K:267:ARG:CB	7:K:268:PRO:CD	2.82	0.58
4:H:49:THR:CB	4:H:50:GLY:CA	2.81	0.58
3:G:95:LYS:O	3:G:98:GLY:N	2.36	0.57
1:A:88:ALA:CB	2:B:83:ALA:HA	2.36	0.56
1:A:72:ARG:O	1:A:75:ALA:HB3	2.09	0.53
1:A:67:PHE:O	1:A:71:VAL:N	2.36	0.53
1:A:88:ALA:HB2	2:B:83:ALA:CA	2.41	0.51
1:A:95:ALA:O	1:A:99:TYR:N	2.39	0.50
7:K:72:ARG:CB	7:K:73:PRO:CD	2.90	0.49
7:K:266:PRO:CB	7:K:267:ARG:CB	2.92	0.48
7:K:72:ARG:CB	7:K:73:PRO:HD2	2.45	0.46
1:A:93:GLN:O	1:A:94:GLU:C	2.54	0.46
7:K:264:LEU:O	7:K:265:ASP:CB	2.64	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:95:LYS:O	3:G:96:LEU:C	2.55	0.45
4:H:48:ASP:HA	4:H:49:THR:HA	1.81	0.45
7:K:47:ASP:N	7:K:463:GLY:O	2.51	0.43
7:K:435:VAL:CB	7:K:436:PRO:HD2	2.48	0.43
4:H:53:SER:N	6:J:-54:DC:OP1	2.52	0.42
7:K:363:PRO:HB3	7:K:434:GLU:CB	2.50	0.42
7:K:468:ASP:O	7:K:472:GLU:N	2.52	0.42
6:J:-57:DT:H2''	6:J:-56:DG:C8	2.54	0.42
1:A:125:GLN:O	1:A:128:ARG:N	2.53	0.42
7:K:329:VAL:O	7:K:345:GLN:N	2.53	0.41
7:K:268:PRO:HB2	7:K:269:PHE:CA	2.51	0.41
4:H:49:THR:CB	4:H:50:GLY:HA2	2.50	0.41
7:K:267:ARG:CB	7:K:268:PRO:HD2	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/135 (72%)	77 (79%)	19 (20%)	1 (1%)	15	54
2	B	76/102 (74%)	59 (78%)	16 (21%)	1 (1%)	12	48
3	G	103/129 (80%)	82 (80%)	18 (18%)	3 (3%)	4	32
4	H	93/122 (76%)	77 (83%)	15 (16%)	1 (1%)	14	52
7	K	414/483 (86%)	316 (76%)	88 (21%)	10 (2%)	6	36
8	M	158/202 (78%)	137 (87%)	18 (11%)	3 (2%)	8	41
All	All	941/1173 (80%)	748 (80%)	174 (18%)	19 (2%)	7	40

All (19) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
7	K	265	ASP
7	K	267	ARG
8	M	280	LYS
7	K	70	VAL
7	K	386	THR
3	G	26	PRO
7	K	28	HIS
7	K	326	ASP
1	A	87	SER
7	K	291	ASN
4	H	47	PRO
7	K	77	PRO
7	K	364	LYS
2	B	29	ILE
3	G	43	VAL
8	M	250	ILE
8	M	284	VAL
7	K	242	VAL
3	G	114	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1/110 (1%)	1 (100%)	0	100	100
2	B	1/78 (1%)	1 (100%)	0	100	100
4	H	1/102 (1%)	1 (100%)	0	100	100
7	K	22/400 (6%)	22 (100%)	0	100	100
All	All	25/690 (4%)	25 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	99/135 (73%)	-0.88	0 100 100	42, 88, 196, 253	0
2	B	78/102 (76%)	-0.76	0 100 100	38, 86, 140, 180	0
3	G	105/129 (81%)	-0.84	0 100 100	39, 103, 172, 195	0
4	H	95/122 (77%)	-0.87	0 100 100	54, 100, 211, 276	0
5	I	73/149 (48%)	-0.12	1 (1%) 75 66	136, 222, 304, 352	0
6	J	72/149 (48%)	-0.31	0 100 100	141, 216, 296, 318	0
7	K	422/483 (87%)	-0.49	1 (0%) 95 93	120, 236, 335, 381	0
8	M	160/202 (79%)	1.43	52 (32%) 0 1	201, 444, 500, 500	0
All	All	1104/1471 (75%)	-0.29	54 (4%) 29 25	38, 205, 488, 500	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
8	M	222	LEU	10.9
8	M	220	ILE	7.4
8	M	221	ASP	6.6
8	M	259	GLU	6.5
8	M	343	SER	6.4
8	M	289	GLU	6.1
8	M	251	GLU	5.6
8	M	344	ILE	5.5
8	M	305	CYS	5.3
8	M	320	LEU	5.1
8	M	345	GLU	5.0
8	M	336	TYR	4.9
8	M	260	ALA	4.7
8	M	224	ASP	4.7
8	M	235	GLN	4.5
8	M	255	ALA	4.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
8	M	252	ILE	4.0
8	M	340	SER	3.9
8	M	236	PHE	3.9
8	M	261	LEU	3.8
8	M	306	GLN	3.8
8	M	263	ALA	3.6
8	M	219	LYS	3.5
5	I	-39	DT	3.4
8	M	196	ALA	3.3
8	M	248	ASP	3.1
8	M	233	THR	3.1
8	M	328	ALA	3.1
8	M	337	GLY	2.9
8	M	262	TYR	2.9
8	M	256	LYS	2.9
8	M	250	ILE	2.8
8	M	258	ARG	2.8
8	M	346	ALA	2.8
8	M	264	GLN	2.7
8	M	338	ASP	2.6
7	K	451	LEU	2.5
8	M	207	ASP	2.5
8	M	232	ALA	2.4
8	M	342	ALA	2.4
8	M	321	ILE	2.4
8	M	276	GLN	2.3
8	M	197	GLU	2.3
8	M	234	LYS	2.3
8	M	278	LEU	2.3
8	M	314	GLY	2.2
8	M	347	PHE	2.2
8	M	335	ASP	2.1
8	M	223	ILE	2.1
8	M	316	PRO	2.1
8	M	290	THR	2.1
8	M	265	ASP	2.0
8	M	313	ASP	2.0
8	M	194	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.