



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 02:51 AM EDT

PDB ID : 5HQE  
Title : Crystal structure of human dihydroorotate dehydrogenase (DHODH) with compound 18T  
Authors : Huang, J.; Wu, D.; Lu, Q.  
Deposited on : 2016-01-21  
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4693 (1.64-1.60)
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)
RSRZ outliers	127900	4609 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	<div> <div>7%</div> <div>81%</div> <div>12%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ACT	A	409	-	-	X	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dihydroorotate dehydrogenase (quinone), mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	365	Total	C	N	O	S	0	0	0
			2790	1749	517	520	4			

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	7	MET	-	initiating methionine	UNP Q02127
A	8	GLY	-	expression tag	UNP Q02127
A	9	HIS	-	expression tag	UNP Q02127
A	10	HIS	-	expression tag	UNP Q02127
A	11	HIS	-	expression tag	UNP Q02127
A	12	HIS	-	expression tag	UNP Q02127
A	13	HIS	-	expression tag	UNP Q02127
A	14	HIS	-	expression tag	UNP Q02127
A	15	HIS	-	expression tag	UNP Q02127
A	16	HIS	-	expression tag	UNP Q02127
A	17	HIS	-	expression tag	UNP Q02127
A	18	HIS	-	expression tag	UNP Q02127
A	19	SER	-	expression tag	UNP Q02127
A	20	SER	-	expression tag	UNP Q02127
A	21	GLY	-	expression tag	UNP Q02127
A	22	HIS	-	expression tag	UNP Q02127
A	23	ILE	-	expression tag	UNP Q02127
A	24	ASP	-	expression tag	UNP Q02127
A	25	ASP	-	expression tag	UNP Q02127
A	26	ASP	-	expression tag	UNP Q02127
A	27	ASP	-	expression tag	UNP Q02127
A	28	LYS	-	expression tag	UNP Q02127
A	29	HIS	-	expression tag	UNP Q02127

- Molecule 2 is methyl (2Z)-cyano[3-(2-fluoro-4-methoxyphenyl)-4-oxo-1,3-thiazolidin-2-ylidene]acetate (three-letter code: 64B) (formula: C<sub>14</sub>H<sub>11</sub>FN<sub>2</sub>O<sub>4</sub>S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			22	14	1	2	4	1		

- Molecule 3 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula:  $C_{17}H_{21}N_4O_9P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 4 is OROTIC ACID (three-letter code: ORO) (formula:  $C_5H_4N_2O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	5	2	4		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	162	Total	O	0	0
			162	162		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase (quinone), mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.62Å 90.62Å 123.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.24 – 1.62 39.24 – 1.62	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.24-1.62) 99.9 (39.24-1.62)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.23 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.160 , 0.182 0.174 , 0.195	Depositor DCC
$R_{free}$ test set	3743 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 54.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3043	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ORO, 64B, ACT, SO4, FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.33	9/2836 (0.3%)	1.36	25/3832 (0.7%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	266	GLU	CD-OE2	-6.95	1.18	1.25
1	A	344	GLU	CD-OE2	6.89	1.33	1.25
1	A	195	TYR	CE1-CZ	-6.84	1.29	1.38
1	A	319	GLU	CD-OE1	6.60	1.32	1.25
1	A	315	GLN	CD-OE1	-5.90	1.10	1.24
1	A	337	SER	CB-OG	-5.50	1.35	1.42
1	A	298	ARG	CZ-NH1	5.35	1.40	1.33
1	A	282	VAL	C-O	-5.20	1.13	1.23
1	A	333	VAL	CB-CG1	-5.02	1.42	1.52

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	311	ASP	CB-CG-OD1	10.66	127.89	118.30
1	A	136	ARG	NE-CZ-NH2	-9.82	115.39	120.30
1	A	207	ASP	CB-CG-OD2	8.91	126.32	118.30
1	A	133	ARG	NE-CZ-NH2	-7.94	116.33	120.30
1	A	136	ARG	NE-CZ-NH1	7.93	124.27	120.30
1	A	393	ASP	CB-CG-OD1	7.38	124.94	118.30
1	A	207	ASP	CB-CG-OD1	-6.89	112.10	118.30
1	A	393	ASP	CB-CG-OD2	-6.71	112.26	118.30
1	A	347	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	A	249	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	A	61	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	A	298	ARG	NE-CZ-NH1	-6.09	117.25	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	GLU	OE1-CD-OE2	-5.90	116.22	123.30
1	A	117	GLU	OE1-CD-OE2	5.85	130.31	123.30
1	A	272	VAL	CG1-CB-CG2	5.74	120.09	110.90
1	A	78	MET	CA-CB-CG	5.57	122.76	113.30
1	A	275	LEU	CB-CG-CD1	5.54	120.42	111.00
1	A	170	LYS	CD-CE-NZ	-5.45	99.16	111.70
1	A	149	PHE	CB-CG-CD1	5.29	124.51	120.80
1	A	133	ARG	NH1-CZ-NH2	5.25	125.18	119.40
1	A	158	GLU	OE1-CD-OE2	-5.22	117.04	123.30
1	A	318	ARG	NE-CZ-NH1	5.19	122.89	120.30
1	A	133	ARG	NE-CZ-NH1	-5.15	117.73	120.30
1	A	99	ASP	CB-CG-OD1	-5.14	113.67	118.30
1	A	113	PHE	CB-CG-CD2	5.03	124.32	120.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2790	0	2849	18	0
2	A	22	0	0	2	0
3	A	31	0	19	0	0
4	A	11	0	3	0	0
5	A	15	0	0	0	0
6	A	12	0	9	3	0
7	A	162	0	0	2	0
All	All	3043	0	2880	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:315:GLN:HE22	1:A:318:ARG:HE	1.31	0.78
1:A:189:VAL:O	1:A:189:VAL:HG12	1.92	0.68
1:A:173:GLU:HG2	7:A:578:HOH:O	1.93	0.68
2:A:401:64B:O5	2:A:401:64B:S12	2.56	0.63
1:A:315:GLN:NE2	1:A:318:ARG:HE	2.00	0.56
1:A:189:VAL:O	1:A:189:VAL:CG1	2.55	0.54
1:A:261:THR:HG21	6:A:409:ACT:H1	1.90	0.54
1:A:200:ARG:NH1	1:A:240:GLU:OE1	2.42	0.53
1:A:123:PRO:HA	1:A:154:LEU:HG	1.92	0.52
1:A:200:ARG:NH1	1:A:240:GLU:CD	2.63	0.52
1:A:200:ARG:HH11	1:A:240:GLU:CD	2.15	0.49
1:A:263:GLN:HB3	6:A:409:ACT:H3	1.95	0.49
1:A:129:ASN:HB3	1:A:130:PRO:HD2	1.96	0.47
1:A:70:ARG:HD2	1:A:72:ARG:HH11	1.79	0.47
1:A:261:THR:OG1	6:A:409:ACT:H1	2.15	0.45
1:A:235:THR:O	1:A:239:GLN:HG3	2.16	0.44
1:A:192:ALA:HB1	1:A:236:LYS:HD3	1.99	0.44
1:A:86:HIS:HB3	7:A:568:HOH:O	2.19	0.42
2:A:401:64B:C14	2:A:401:64B:C2	2.98	0.41
1:A:282:VAL:HA	1:A:283:THR:HA	1.83	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/390 (93%)	351 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	294/317 (93%)	289 (98%)	5 (2%)	60 36

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	ARG
1	A	307	LYS
1	A	356	TYR
1	A	360	THR
1	A	396	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	315	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	ORO	A	403	-	9,11,11	1.91	3 (33%)	8,15,15	2.15	2 (25%)
5	SO4	A	406	-	4,4,4	0.92	0	6,6,6	0.92	0
2	64B	A	401	-	22,23,23	2.50	9 (40%)	25,32,32	2.38	11 (44%)
6	ACT	A	407	-	3,3,3	1.06	0	3,3,3	1.04	0
6	ACT	A	409	-	3,3,3	1.23	0	3,3,3	1.56	1 (33%)
5	SO4	A	404	-	4,4,4	0.78	0	6,6,6	0.41	0
6	ACT	A	408	-	3,3,3	0.83	0	3,3,3	0.83	0
3	FMN	A	402	-	33,33,33	1.43	4 (12%)	48,50,50	1.22	6 (12%)
5	SO4	A	405	-	4,4,4	0.61	0	6,6,6	1.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	64B	A	401	-	-	0/16/31/31	0/2/2/2
4	ORO	A	403	-	-	4/4/4/4	0/1/1/1
3	FMN	A	402	-	-	1/18/18/18	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	64B	C2-C1	-5.80	1.31	1.43
2	A	401	64B	C10-N9	-5.78	1.32	1.40
3	A	402	FMN	C9A-C5A	5.02	1.49	1.41
2	A	401	64B	C2-N3	3.38	1.20	1.14
3	A	402	FMN	C8-C7	3.31	1.49	1.40
4	A	403	ORO	C2-N1	-3.25	1.31	1.38
4	A	403	ORO	O72-C7	-3.13	1.21	1.30
2	A	401	64B	C11-S12	-3.03	1.75	1.80
2	A	401	64B	O21-C17	2.93	1.43	1.37
2	A	401	64B	C18-C17	-2.61	1.34	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	FMN	C4A-N5	2.51	1.35	1.30
2	A	401	64B	C8-N9	-2.45	1.34	1.39
3	A	402	FMN	C4-N3	-2.45	1.34	1.38
2	A	401	64B	C14-N9	2.41	1.48	1.44
2	A	401	64B	O6-C4	2.29	1.38	1.33
4	A	403	ORO	C2-N3	-2.28	1.33	1.38

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	64B	C14-N9-C10	-7.68	113.78	122.67
4	A	403	ORO	C5-C4-N3	-4.64	118.66	124.08
2	A	401	64B	O5-C4-C1	-3.27	118.33	125.17
2	A	401	64B	C18-C19-C14	-3.13	120.76	123.34
2	A	401	64B	O13-C10-N9	3.03	126.72	124.01
2	A	401	64B	C10-C11-S12	-2.77	103.11	107.59
4	A	403	ORO	C7-C6-N1	2.77	120.56	116.48
3	A	402	FMN	C4-C4A-N5	2.74	122.13	118.23
2	A	401	64B	C15-C14-C19	2.66	122.62	117.30
3	A	402	FMN	C4A-C10-N1	-2.65	118.58	124.73
2	A	401	64B	C11-C10-N9	2.52	112.70	111.22
2	A	401	64B	C2-C1-C4	-2.45	111.80	117.17
2	A	401	64B	C16-C15-C14	-2.40	114.27	119.19
3	A	402	FMN	C10-N1-C2	2.24	121.38	116.90
3	A	402	FMN	O4-C4-C4A	-2.18	120.81	126.60
2	A	401	64B	O6-C4-C1	2.10	115.37	111.41
3	A	402	FMN	C4A-C10-N10	2.07	119.51	116.48
3	A	402	FMN	C4A-C4-N3	2.06	118.43	113.19
6	A	409	ACT	O-C-CH3	-2.05	114.37	122.33
2	A	401	64B	C15-C16-C17	2.02	122.20	119.73

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	403	ORO	N1-C6-C7-O72
4	A	403	ORO	C5-C6-C7-O71
4	A	403	ORO	C5-C6-C7-O72
4	A	403	ORO	N1-C6-C7-O71
3	A	402	FMN	C4'-C5'-O5'-P


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	64B	2	0
6	A	409	ACT	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	365/390 (93%)	-0.03	26 (7%) <b>16</b> <b>14</b>	8, 17, 37, 76	12 (3%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	211	VAL	7.0
1	A	70	ARG	6.3
1	A	213	VAL	5.7
1	A	72	ARG	5.6
1	A	71	ALA	5.3
1	A	32	THR	4.2
1	A	33	GLY	4.2
1	A	362	TRP	3.9
1	A	73	PHE	3.7
1	A	212	ASN	3.7
1	A	219	ALA	3.6
1	A	189	VAL	3.4
1	A	74	GLN	3.3
1	A	37	PHE	3.3
1	A	224	LEU	3.2
1	A	226	GLY	3.1
1	A	313	SER	2.9
1	A	396	ARG	2.8
1	A	134	VAL	2.7
1	A	223	SER	2.5
1	A	36	ARG	2.2
1	A	229	GLU	2.1
1	A	218	THR	2.1
1	A	187	THR	2.1
1	A	225	GLN	2.0
1	A	220	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	ACT	A	407	4/4	0.33	0.27	33,41,50,53	0
6	ACT	A	408	4/4	0.56	0.15	57,61,63,65	0
6	ACT	A	409	4/4	0.90	0.10	25,27,43,45	0
5	SO4	A	404	5/5	0.93	0.32	62,63,66,68	0
2	64B	A	401	22/22	0.95	0.10	19,24,28,31	0
3	FMN	A	402	31/31	0.96	0.09	9,11,12,13	0
4	ORO	A	403	11/11	0.96	0.06	11,12,15,15	0
5	SO4	A	406	5/5	0.98	0.17	29,36,40,49	0
5	SO4	A	405	5/5	0.99	0.16	29,29,38,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 64B A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.