



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 28, 2024 – 07:50 PM EDT

PDB ID : 3HTM
Title : Structures of SPOP-Substrate Complexes: Insights into Molecular Architectures of BTB-Cul3 Ubiquitin Ligases: SPOBTTB/3-box
Authors : Zhuang, M.; Walden, H.; Schulman, B.A.
Deposited on : 2009-06-11
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	<div> <div>70%</div> <div>18%</div> <div>11%</div> </div>
1	B	172	<div> <div>66%</div> <div>23%</div> <div>11%</div> </div>
1	C	172	<div> <div>68%</div> <div>20%</div> <div>12%</div> </div>
1	D	172	<div> <div>73%</div> <div>14%</div> <div>12%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Speckle-type POZ protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	Se	0	0	0
			1191	748	199	230	7	7			
1	B	153	Total	C	N	O	S	Se	0	0	0
			1191	747	198	232	7	7			
1	C	152	Total	C	N	O	S	Se	0	0	0
			1187	745	198	231	7	6			
1	D	152	Total	C	N	O	S	Se	0	0	0
			1187	745	198	231	7	6			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	166	GLY	-	expression tag	UNP O43791
A	167	SER	-	expression tag	UNP O43791
A	168	GLY	-	expression tag	UNP O43791
A	169	GLY	-	expression tag	UNP O43791
A	170	SER	-	expression tag	UNP O43791
A	171	GLY	-	expression tag	UNP O43791
A	330	THR	-	linker	UNP O43791
A	331	ASP	-	linker	UNP O43791
A	332	VAL	-	linker	UNP O43791
A	333	LEU	-	linker	UNP O43791
A	334	GLU	-	linker	UNP O43791
A	335	THR	-	linker	UNP O43791
A	336	SER	-	linker	UNP O43791
A	337	GLY	-	linker	UNP O43791
B	166	GLY	-	expression tag	UNP O43791
B	167	SER	-	expression tag	UNP O43791
B	168	GLY	-	expression tag	UNP O43791
B	169	GLY	-	expression tag	UNP O43791
B	170	SER	-	expression tag	UNP O43791
B	171	GLY	-	expression tag	UNP O43791
B	330	THR	-	linker	UNP O43791

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	331	ASP	-	linker	UNP O43791
B	332	VAL	-	linker	UNP O43791
B	333	LEU	-	linker	UNP O43791
B	334	GLU	-	linker	UNP O43791
B	335	THR	-	linker	UNP O43791
B	336	SER	-	linker	UNP O43791
B	337	GLY	-	linker	UNP O43791
C	166	GLY	-	expression tag	UNP O43791
C	167	SER	-	expression tag	UNP O43791
C	168	GLY	-	expression tag	UNP O43791
C	169	GLY	-	expression tag	UNP O43791
C	170	SER	-	expression tag	UNP O43791
C	171	GLY	-	expression tag	UNP O43791
C	330	THR	-	linker	UNP O43791
C	331	ASP	-	linker	UNP O43791
C	332	VAL	-	linker	UNP O43791
C	333	LEU	-	linker	UNP O43791
C	334	GLU	-	linker	UNP O43791
C	335	THR	-	linker	UNP O43791
C	336	SER	-	linker	UNP O43791
C	337	GLY	-	linker	UNP O43791
D	166	GLY	-	expression tag	UNP O43791
D	167	SER	-	expression tag	UNP O43791
D	168	GLY	-	expression tag	UNP O43791
D	169	GLY	-	expression tag	UNP O43791
D	170	SER	-	expression tag	UNP O43791
D	171	GLY	-	expression tag	UNP O43791
D	330	THR	-	linker	UNP O43791
D	331	ASP	-	linker	UNP O43791
D	332	VAL	-	linker	UNP O43791
D	333	LEU	-	linker	UNP O43791
D	334	GLU	-	linker	UNP O43791
D	335	THR	-	linker	UNP O43791
D	336	SER	-	linker	UNP O43791
D	337	GLY	-	linker	UNP O43791

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	82	Total O 82 82	0	0
2	B	93	Total O 93 93	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	73	Total 73	O 73	0	0
2	D	88	Total 88	O 88	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Speckle-type POZ protein




- Molecule 1: Speckle-type POZ protein



- Molecule 1: Speckle-type POZ protein



- Molecule 1: Speckle-type POZ protein



LEU
GLU
THR
SER
GLY

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	36.80Å 88.70Å 88.70Å 90.80° 89.30° 89.90°	Depositor
Resolution (Å)	44.34 – 2.50 44.34 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.9 (44.34-2.50) 98.3 (44.34-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.90 (at 2.42Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.217 , 0.256 0.226 , 0.227	Depositor DCC
R_{free} test set	1916 reflections (5.23%)	wwPDB-VP
Wilson B-factor (Å ²)	42.2	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for h,-l,k 0.013 for h,l,-k 0.470 for h,-k,-l 0.015 for -h,-k,l 0.014 for -h,k,-l 0.477 for -h,l,k 0.470 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5092	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1203	0.58	0/1611
1	B	0.38	0/1203	0.60	0/1612
1	C	0.36	0/1199	0.57	0/1606
1	D	0.36	0/1199	0.59	0/1606
All	All	0.37	0/4804	0.59	0/6435

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1191	0	1164	35	0
1	B	1191	0	1157	50	0
1	C	1187	0	1158	43	0
1	D	1187	0	1158	33	0
2	A	82	0	0	6	0
2	B	93	0	0	13	0
2	C	73	0	0	7	0
2	D	88	0	0	9	0
All	All	5092	0	4637	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:200:THR:HG21	1:C:212:GLN:HB3	1.45	0.99
1:C:190:LEU:HD11	1:D:190:LEU:HD11	1.47	0.96
1:D:205:CYS:HB3	2:D:370:HOH:O	1.68	0.93
1:C:228:MSE:O	1:C:232:GLU:HG2	1.71	0.90
1:A:266:LEU:HD13	1:A:273:LEU:HD23	1.54	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/172 (88%)	143 (95%)	8 (5%)	0	100	100
1	B	151/172 (88%)	147 (97%)	4 (3%)	0	100	100
1	C	148/172 (86%)	140 (95%)	8 (5%)	0	100	100
1	D	148/172 (86%)	140 (95%)	8 (5%)	0	100	100
All	All	598/688 (87%)	570 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/134 (96%)	126 (98%)	2 (2%)	58	80
1	B	128/134 (96%)	126 (98%)	2 (2%)	58	80
1	C	128/134 (96%)	127 (99%)	1 (1%)	79	91
1	D	128/134 (96%)	125 (98%)	3 (2%)	45	72
All	All	512/536 (96%)	504 (98%)	8 (2%)	58	80

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	266	LEU
1	D	245	ASP
1	C	198	ARG
1	B	266	LEU
1	D	221	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	214	HIS
1	C	296	ASN
1	D	177	ASN
1	D	244	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	146/172 (84%)	-1.11	1 (0%) 84 81	24, 44, 71, 93	0
1	B	146/172 (84%)	-1.19	0 100 100	25, 43, 69, 93	0
1	C	146/172 (84%)	-1.20	0 100 100	24, 44, 72, 94	0
1	D	146/172 (84%)	-1.08	0 100 100	24, 44, 68, 92	0
All	All	584/688 (84%)	-1.14	1 (0%) 92 90	24, 44, 71, 94	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	329	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.