



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2024 – 02:05 AM EDT

PDB ID : 4HWB  
Title : Crystal structure of ectodomain 3 of the IL-13 receptor alpha 1 in complex with a human neutralizing monoclonal antibody fragment  
Authors : Xu, Y.  
Deposited on : 2012-11-07  
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4623 (2.64-2.60)
Clashscore	180529	5071 (2.64-2.60)
Ramachandran outliers	177936	5006 (2.64-2.60)
Sidechain outliers	177891	5006 (2.64-2.60)
RSRZ outliers	164620	4622 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	<div> <div>25%</div> <div> <div></div> <div>79%</div> <div>11%</div> <div>9%</div> </div> </div>
2	H	227	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>
3	L	212	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>13%</div> <div></div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-13 receptor subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	111	Total	C	N	O	S	0	0	0
			849	540	144	160	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	311	HIS	-	expression tag	UNP P78552
A	312	HIS	-	expression tag	UNP P78552
A	313	HIS	-	expression tag	UNP P78552
A	314	HIS	-	expression tag	UNP P78552
A	315	HIS	-	expression tag	UNP P78552
A	316	HIS	-	expression tag	UNP P78552
A	317	HIS	-	expression tag	UNP P78552
A	318	HIS	-	expression tag	UNP P78552

- Molecule 2 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	216	Total	C	N	O	S	0	0	0
			1620	1035	263	314	8			

- Molecule 3 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	212	Total	C	N	O	S	0	0	0
			1602	1001	269	328	4			


- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	L	1	Total	O	S	0	0
			5	4	1		
4	L	1	Total	O	S	0	0
			5	4	1		
4	L	1	Total	O	S	0	0
			5	4	1		
4	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (three-letter code:

NHE) (formula: C<sub>8</sub>H<sub>17</sub>NO<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			13	8	1	3	1		
5	A	1	Total	C	N	O	S	0	0
			13	8	1	3	1		
5	A	1	Total	C	N	O	S	0	0
			13	8	1	3	1		
5	H	1	Total	C	N	O	S	0	0
			13	8	1	3	1		
5	H	1	Total	C	N	O	S	0	0
			13	8	1	3	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	10	Total	O	0	0
			10	10		
6	H	50	Total	O	0	0
			50	50		
6	L	17	Total	O	0	0
			17	17		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-13 receptor subunit alpha-1



- Molecule 2: Fab heavy chain



- Molecule 3: Fab light chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.48Å 123.48Å 181.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.17 – 2.61 47.17 – 2.61	Depositor EDS
% Data completeness (in resolution range)	98.5 (47.17-2.61) 98.5 (47.17-2.61)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.29 (at 2.61Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, $R_{free}$	0.197 , 0.224 0.230 , 0.267	Depositor DCC
$R_{free}$ test set	2148 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.1	Xtriage
Anisotropy	0.531	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 74.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4278	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NHE, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/872	0.83	1/1196 (0.1%)
2	H	0.51	0/1666	0.78	0/2271
3	L	0.46	0/1636	0.74	0/2227
All	All	0.51	0/4174	0.77	1/5694 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	242	GLN	C-N-CA	5.83	136.28	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	849	0	740	7	0
2	H	1620	0	1574	13	0
3	L	1602	0	1524	10	0
4	A	10	0	0	0	0
4	H	25	0	0	0	0
4	L	30	0	0	0	0
5	A	39	0	48	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	26	0	32	2	0
6	A	10	0	0	0	0
6	H	50	0	0	0	0
6	L	17	0	0	0	0
All	All	4278	0	3918	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:20:LYS:HG3	2:H:83:GLN:HB3	1.78	0.65
1:A:293:LEU:HD13	2:H:51:VAL:HG21	1.86	0.58
2:H:34:TRP:HB2	2:H:100:MET:HG3	1.85	0.58
3:L:160:VAL:HG22	3:L:172:LEU:HD12	1.85	0.57
2:H:48:TRP:CG	3:L:93:GLU:HA	2.42	0.54
1:A:205:PRO:HB3	5:A:404:NHE:H2'2	1.90	0.54
1:A:210:LEU:HD23	1:A:217:LEU:HD11	1.90	0.53
1:A:295:TYR:O	2:H:60:ARG:NH2	2.41	0.53
2:H:92:THR:HG23	2:H:116:THR:HA	1.90	0.53
2:H:19:LEU:HB2	2:H:87:LEU:HD11	1.93	0.49
1:A:224:PRO:HD2	5:A:404:NHE:HC12	1.95	0.49
1:A:249:PHE:CE1	1:A:274:PRO:HD2	2.48	0.49
2:H:60:ARG:HD2	5:H:307:NHE:O3	2.13	0.48
2:H:48:TRP:CD2	3:L:93:GLU:HA	2.47	0.48
3:L:6:GLN:H	3:L:97:GLN:HE22	1.62	0.47
3:L:178:LEU:HD13	3:L:183:TYR:HB2	1.97	0.46
2:H:37:TRP:O	2:H:49:MET:HB2	2.16	0.46
3:L:40:LYS:HE3	3:L:82:GLU:O	2.16	0.45
3:L:117:PRO:HD3	3:L:129:VAL:HG22	1.98	0.45
5:H:307:NHE:H6'2	5:H:307:NHE:HC12	1.76	0.45
3:L:48:LEU:HA	3:L:59:ILE:HG13	1.98	0.45
3:L:148:ASP:HA	3:L:188:VAL:HB	1.98	0.45
2:H:13:LYS:HD2	2:H:19:LEU:HD13	2.01	0.42
2:H:98:ALA:HB1	2:H:106:PHE:HB3	2.01	0.42
2:H:33:TYR:O	2:H:54:PRO:HD2	2.19	0.42
3:L:142:LYS:HB3	3:L:194:THR:HB	2.01	0.41
1:A:213:HIS:HB3	1:A:218:TYR:HE2	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/122 (89%)	94 (86%)	12 (11%)	3 (3%)	4	6
2	H	212/227 (93%)	205 (97%)	7 (3%)	0	100	100
3	L	210/212 (99%)	203 (97%)	6 (3%)	1 (0%)	25	45
All	All	531/561 (95%)	502 (94%)	25 (5%)	4 (1%)	16	32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	SER
1	A	258	ASN
3	L	135	ASN
1	A	265	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/120 (72%)	85 (98%)	2 (2%)	45	69
2	H	180/193 (93%)	169 (94%)	11 (6%)	15	32
3	L	176/183 (96%)	163 (93%)	13 (7%)	11	23
All	All	443/496 (89%)	417 (94%)	26 (6%)	16	33

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	210	LEU
1	A	272	MET
2	H	14	LYS
2	H	62	SER
2	H	100	MET
2	H	116	THR
2	H	162	SER
2	H	165	LEU
2	H	176	LEU
2	H	184	LEU
2	H	185	SER
2	H	199	THR
2	H	216	LYS
3	L	5	THR
3	L	24	ARG
3	L	30	SER
3	L	34	LEU
3	L	46	ARG
3	L	49	ILE
3	L	91	GLN
3	L	102	GLU
3	L	159	SER
3	L	174	SER
3	L	178	LEU
3	L	180	LYS
3	L	194	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	L	27	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NHE	H	307	-	13,13,13	1.44	1 (7%)	16,17,17	0.86	1 (6%)
4	SO4	A	401	-	4,4,4	0.30	0	6,6,6	0.17	0
4	SO4	L	301	-	4,4,4	0.32	0	6,6,6	0.08	0
4	SO4	H	302	-	4,4,4	0.54	0	6,6,6	0.24	0
5	NHE	A	405	-	13,13,13	1.54	1 (7%)	16,17,17	0.74	1 (6%)
4	SO4	L	304	-	4,4,4	0.28	0	6,6,6	0.18	0
4	SO4	H	303	-	4,4,4	0.33	0	6,6,6	0.19	0
5	NHE	A	404	-	13,13,13	1.58	1 (7%)	16,17,17	0.70	1 (6%)
4	SO4	L	302	-	4,4,4	0.26	0	6,6,6	0.17	0
4	SO4	L	305	-	4,4,4	0.29	0	6,6,6	0.17	0
4	SO4	L	306	-	4,4,4	0.31	0	6,6,6	0.18	0
5	NHE	H	306	-	13,13,13	1.20	1 (7%)	16,17,17	0.52	0
4	SO4	L	303	-	4,4,4	0.33	0	6,6,6	0.18	0
5	NHE	A	403	-	13,13,13	2.01	1 (7%)	16,17,17	0.85	1 (6%)
4	SO4	H	305	-	4,4,4	0.31	0	6,6,6	0.15	0
4	SO4	H	301	-	4,4,4	0.29	0	6,6,6	0.16	0
4	SO4	A	402	-	4,4,4	0.28	0	6,6,6	0.10	0
4	SO4	H	304	-	4,4,4	0.32	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NHE	H	307	-	-	2/7/15/15	0/1/1/1
5	NHE	A	405	-	-	1/7/15/15	0/1/1/1
5	NHE	A	404	-	-	2/7/15/15	0/1/1/1
5	NHE	H	306	-	-	5/7/15/15	0/1/1/1
5	NHE	A	403	-	-	1/7/15/15	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	403	NHE	C2-S	-7.18	1.67	1.77
5	A	404	NHE	C2-S	-5.53	1.69	1.77
5	A	405	NHE	C2-S	-5.40	1.70	1.77
5	H	307	NHE	C2-S	-4.75	1.70	1.77
5	H	306	NHE	C2-S	-3.83	1.72	1.77

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	307	NHE	C1-N-C1'	2.75	119.48	114.18
5	A	403	NHE	C6'-C1'-N	2.71	124.09	110.56
5	A	405	NHE	C6'-C1'-N	2.64	123.72	110.56
5	A	404	NHE	C2'-C1'-N	2.55	123.31	110.56

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	404	NHE	N-C1-C2-S
5	H	306	NHE	C1-C2-S-O1
5	H	307	NHE	C6'-C1'-N-C1
5	H	306	NHE	C1-C2-S-O3
5	A	404	NHE	C2-C1-N-C1'
5	A	405	NHE	C2-C1-N-C1'
5	H	306	NHE	C2'-C1'-N-C1
5	H	306	NHE	C1-C2-S-O2
5	H	306	NHE	C6'-C1'-N-C1
5	A	403	NHE	N-C1-C2-S
5	H	307	NHE	N-C1-C2-S

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	307	NHE	2	0
5	A	404	NHE	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	111/122 (90%)	1.49	30 (27%) 2 2	47, 79, 135, 145	0
2	H	216/227 (95%)	-0.19	7 (3%) 50 45	36, 54, 106, 119	0
3	L	212/212 (100%)	0.10	4 (1%) 66 62	45, 72, 103, 133	0
All	All	539/561 (96%)	0.27	41 (7%) 21 18	36, 66, 115, 145	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	260	GLU	8.1
3	L	212	ALA	7.5
1	A	241	SER	7.4
3	L	211	CYS	6.6
1	A	265	VAL	6.5
1	A	200	VAL	6.4
1	A	264	ASN	6.4
1	A	261	PHE	6.2
1	A	310	ILE	6.1
1	A	259	PRO	6.0
1	A	266	GLU	5.8
1	A	263	ARG	5.7
1	A	257	GLU	5.5
1	A	202	PRO	5.4
1	A	240	ASN	5.2
1	A	244	GLU	4.6
1	A	267	ASN	4.6
1	A	245	THR	4.5
1	A	305	SER	4.3
1	A	243	THR	4.3
1	A	201	LYS	4.2
1	A	242	GLN	3.9
1	A	258	ASN	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	H	133	SER	3.5
2	H	220	LYS	3.5
2	H	1	ALA	3.5
2	H	134	SER	3.4
1	A	221	TRP	3.1
1	A	309	SER	3.0
1	A	308	MET	2.8
1	A	306	GLN	2.7
1	A	246	HIS	2.7
2	H	140	GLY	2.6
2	H	141	THR	2.5
3	L	78	ARG	2.4
1	A	204	PRO	2.4
2	H	139	GLY	2.4
3	L	146	LYS	2.2
1	A	275	GLY	2.2
1	A	272	MET	2.0
1	A	298	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	L	305	5/5	0.52	0.15	167,167,168,168	0
4	SO4	H	304	5/5	0.56	0.18	151,152,153,153	0
4	SO4	L	302	5/5	0.63	0.12	158,158,159,159	0
4	SO4	L	304	5/5	0.64	0.17	172,173,174,174	0
4	SO4	L	303	5/5	0.64	0.12	136,136,137,137	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	L	301	5/5	0.66	0.17	162,162,163,164	0
4	SO4	L	306	5/5	0.66	0.15	161,161,161,162	0
4	SO4	H	301	5/5	0.72	0.10	154,154,155,155	0
4	SO4	H	305	5/5	0.77	0.17	147,148,148,149	0
4	SO4	A	402	5/5	0.80	0.09	139,139,140,140	0
5	NHE	H	307	13/13	0.80	0.23	72,79,92,93	0
4	SO4	H	303	5/5	0.82	0.11	144,144,144,144	0
4	SO4	A	401	5/5	0.83	0.12	139,140,141,142	0
5	NHE	H	306	13/13	0.89	0.17	77,85,102,105	0
4	SO4	H	302	5/5	0.89	0.11	93,95,97,98	0
5	NHE	A	405	13/13	0.90	0.20	93,98,123,123	0
5	NHE	A	404	13/13	0.96	0.11	67,70,72,72	0
5	NHE	A	403	13/13	0.97	0.09	44,48,53,57	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.