



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 22, 2024 – 08:15 AM EDT

PDB ID : 4I3U
Title : Structure of phosphonoacetaldehyde dehydrogenase in complex with phosphonoacetaldehyde
Authors : Nair, S.K.; Agarwal, V.
Deposited on : 2012-11-26
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>20%</div> <div>• •</div> </div> </div>
1	B	488	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>22%</div> <div>• •</div> </div> </div>
1	C	488	<div> <div>2%</div> <div> <div></div> <div>76%</div> <div>18%</div> <div>• •</div> </div> </div>
1	D	488	<div> <div>5%</div> <div> <div></div> <div>75%</div> <div>19%</div> <div>• •</div> </div> </div>
1	E	488	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>20%</div> <div>• •</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	488	<p>2% 73% 20% . .</p>
1	G	488	<p>3% 75% 20% . .</p>
1	H	488	<p>6% 71% 22% . .</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	POA	B	500	-	X	-	-
2	POA	D	500	-	X	-	-
2	POA	E	500	-	X	-	-
2	POA	G	500	-	X	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 30980 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldehyde dehydrogenase (NAD⁺).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			
1	B	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			
1	C	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			
1	D	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			
1	E	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			
1	F	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			
1	G	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			
1	H	473	Total	C	N	O	S	0	0	0
			3626	2294	630	683	19			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q92UV7
A	-1	SER	-	expression tag	UNP Q92UV7
A	0	HIS	-	expression tag	UNP Q92UV7
B	-2	GLY	-	expression tag	UNP Q92UV7
B	-1	SER	-	expression tag	UNP Q92UV7
B	0	HIS	-	expression tag	UNP Q92UV7
C	-2	GLY	-	expression tag	UNP Q92UV7
C	-1	SER	-	expression tag	UNP Q92UV7
C	0	HIS	-	expression tag	UNP Q92UV7
D	-2	GLY	-	expression tag	UNP Q92UV7
D	-1	SER	-	expression tag	UNP Q92UV7
D	0	HIS	-	expression tag	UNP Q92UV7
E	-2	GLY	-	expression tag	UNP Q92UV7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	SER	-	expression tag	UNP Q92UV7
E	0	HIS	-	expression tag	UNP Q92UV7
F	-2	GLY	-	expression tag	UNP Q92UV7
F	-1	SER	-	expression tag	UNP Q92UV7
F	0	HIS	-	expression tag	UNP Q92UV7
G	-2	GLY	-	expression tag	UNP Q92UV7
G	-1	SER	-	expression tag	UNP Q92UV7
G	0	HIS	-	expression tag	UNP Q92UV7
H	-2	GLY	-	expression tag	UNP Q92UV7
H	-1	SER	-	expression tag	UNP Q92UV7
H	0	HIS	-	expression tag	UNP Q92UV7

- Molecule 2 is PHOSPHONOACETALDEHYDE (three-letter code: POA) (formula: $C_2H_5O_4P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			7	2	4	1		
2	B	1	Total	C	O	P	0	0
			7	2	4	1		
2	C	1	Total	C	O	P	0	0
			7	2	4	1		
2	D	1	Total	C	O	P	0	0
			7	2	4	1		
2	E	1	Total	C	O	P	0	0
			7	2	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	F	1	Total	C	O	P	0	0
			7	2	4	1		
2	G	1	Total	C	O	P	0	0
			7	2	4	1		
2	H	1	Total	C	O	P	0	0
			7	2	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	287	Total	O	0	0
			287	287		
3	B	233	Total	O	0	0
			233	233		
3	C	237	Total	O	0	0
			237	237		
3	D	258	Total	O	0	0
			258	258		
3	E	245	Total	O	0	0
			245	245		
3	F	225	Total	O	0	0
			225	225		
3	G	204	Total	O	0	0
			204	204		
3	H	227	Total	O	0	0
			227	227		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Aldehyde dehydrogenase (NAD⁺)




• Molecule 1: Aldehyde dehydrogenase (NAD⁺)




• Molecule 1: Aldehyde dehydrogenase (NAD⁺)









- Molecule 1: Aldehyde dehydrogenase (NAD⁺)



- Molecule 1: Aldehyde dehydrogenase (NAD⁺)





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	93.35Å 172.88Å 138.97Å 90.00° 106.73° 90.00°	Depositor
Resolution (Å)	39.11 – 2.10 39.11 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.6 (39.11-2.10) 96.6 (39.11-2.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.7.1_743, REFMAC	Depositor
R, R_{free}	0.210 , 0.253 0.205 , 0.247	Depositor DCC
R_{free} test set	11837 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	30980	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: POA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	6/3698 (0.2%)	0.84	15/5030 (0.3%)
1	B	0.71	6/3698 (0.2%)	0.79	9/5030 (0.2%)
1	C	0.76	9/3698 (0.2%)	0.83	8/5030 (0.2%)
1	D	0.84	8/3698 (0.2%)	0.85	19/5030 (0.4%)
1	E	0.78	15/3698 (0.4%)	0.86	12/5030 (0.2%)
1	F	0.83	11/3698 (0.3%)	0.82	13/5030 (0.3%)
1	G	0.73	11/3698 (0.3%)	0.81	10/5030 (0.2%)
1	H	0.74	6/3698 (0.2%)	0.84	8/5030 (0.2%)
All	All	0.77	72/29584 (0.2%)	0.83	94/40240 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	E	0	1
1	F	0	1
All	All	0	3

The worst 5 of 72 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	423	TYR	CE1-CZ	-15.98	1.17	1.38
1	D	423	TYR	CE2-CZ	-14.37	1.19	1.38
1	F	423	TYR	CE2-CZ	-14.16	1.20	1.38
1	A	423	TYR	CE1-CZ	-14.14	1.20	1.38
1	F	423	TYR	CG-CD1	-13.85	1.21	1.39

The worst 5 of 94 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	314	ARG	NE-CZ-NH2	16.74	128.67	120.30
1	A	314	ARG	NE-CZ-NH2	14.38	127.49	120.30
1	C	314	ARG	NE-CZ-NH2	13.84	127.22	120.30
1	G	314	ARG	NE-CZ-NH2	13.18	126.89	120.30
1	C	343	GLU	OE1-CD-OE2	-11.86	109.07	123.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	314	ARG	Sidechain
1	E	377	HIS	Sidechain
1	F	314	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3626	0	3654	161	1
1	B	3626	0	3654	138	1
1	C	3626	0	3654	124	1
1	D	3626	0	3654	135	8
1	E	3626	0	3654	133	0
1	F	3626	0	3654	135	0
1	G	3626	0	3654	128	0
1	H	3626	0	3654	149	8
2	A	7	0	2	2	0
2	B	7	0	2	2	0
2	C	7	0	2	3	0
2	D	7	0	2	2	0
2	E	7	0	2	3	0
2	F	7	0	2	2	0
2	G	7	0	2	3	0
2	H	7	0	2	1	0
3	A	287	0	0	31	0
3	B	233	0	0	33	0
3	C	237	0	0	18	0
3	D	258	0	0	19	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	245	0	0	23	0
3	F	225	0	0	31	0
3	G	204	0	0	33	1
3	H	227	0	0	30	1
All	All	30980	0	29248	1023	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 1023 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:HIS:CE1	1:C:85:GLU:HG3	1.47	1.45
1:A:359:HIS:O	1:G:326:ARG:CD	1.76	1.32
1:C:216:ASP:OD2	3:C:824:HOH:O	1.55	1.24
1:H:314:ARG:NH1	3:H:633:HOH:O	1.64	1.23
1:D:314:ARG:NH2	3:D:739:HOH:O	1.72	1.23

The worst 5 of 11 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:348:ARG:NH1	1:H:216:ASP:OD2[1_655]	1.18	1.02
1:D:348:ARG:NH1	1:H:216:ASP:CG[1_655]	1.59	0.61
1:A:269:ASP:OD2	1:C:226:HIS:CE1[2_444]	1.98	0.22
1:D:348:ARG:CG	1:H:216:ASP:OD2[1_655]	1.98	0.22
1:D:348:ARG:NH1	1:H:216:ASP:CB[1_655]	2.01	0.19

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	471/488 (96%)	454 (96%)	15 (3%)	2 (0%)	30	29
1	B	471/488 (96%)	456 (97%)	14 (3%)	1 (0%)	44	45
1	C	471/488 (96%)	456 (97%)	14 (3%)	1 (0%)	44	45
1	D	471/488 (96%)	454 (96%)	15 (3%)	2 (0%)	30	29
1	E	471/488 (96%)	453 (96%)	16 (3%)	2 (0%)	30	29
1	F	471/488 (96%)	455 (97%)	14 (3%)	2 (0%)	30	29
1	G	471/488 (96%)	455 (97%)	15 (3%)	1 (0%)	44	45
1	H	471/488 (96%)	455 (97%)	15 (3%)	1 (0%)	44	45
All	All	3768/3904 (96%)	3638 (96%)	118 (3%)	12 (0%)	37	37

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	392	PRO
1	C	392	PRO
1	E	392	PRO
1	B	392	PRO
1	D	392	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/401 (97%)	375 (96%)	14 (4%)	30	32
1	B	389/401 (97%)	376 (97%)	13 (3%)	33	36
1	C	389/401 (97%)	376 (97%)	13 (3%)	33	36
1	D	389/401 (97%)	375 (96%)	14 (4%)	30	32
1	E	389/401 (97%)	372 (96%)	17 (4%)	24	24
1	F	389/401 (97%)	376 (97%)	13 (3%)	33	36
1	G	389/401 (97%)	376 (97%)	13 (3%)	33	36
1	H	389/401 (97%)	372 (96%)	17 (4%)	24	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	3112/3208 (97%)	2998 (96%)	114 (4%)	29	31

5 of 114 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	51	ARG
1	H	377	HIS
1	F	12	HIS
1	H	359	HIS
1	H	61	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	247	HIS
1	F	176	ASN
1	H	427	GLN
1	G	247	HIS
1	G	377	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	POA	H	500	1	6,6,6	1.84	3 (50%)	7,8,8	2.54	4 (57%)
2	POA	A	500	1	6,6,6	2.25	3 (50%)	7,8,8	2.31	4 (57%)
2	POA	F	500	1	6,6,6	1.75	1 (16%)	7,8,8	2.81	4 (57%)
2	POA	D	500	1	6,6,6	2.26	4 (66%)	7,8,8	2.49	5 (71%)
2	POA	B	500	1	6,6,6	1.88	2 (33%)	7,8,8	2.67	6 (85%)
2	POA	C	500	1	6,6,6	1.75	1 (16%)	7,8,8	2.23	5 (71%)
2	POA	G	500	1	6,6,6	2.28	3 (50%)	7,8,8	2.22	5 (71%)
2	POA	E	500	1	6,6,6	2.21	3 (50%)	7,8,8	2.74	5 (71%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	POA	H	500	1	-	0/3/4/4	-
2	POA	A	500	1	-	0/3/4/4	-
2	POA	F	500	1	-	0/3/4/4	-
2	POA	D	500	1	-	0/3/4/4	-
2	POA	B	500	1	-	0/3/4/4	-
2	POA	C	500	1	-	0/3/4/4	-
2	POA	G	500	1	-	0/3/4/4	-
2	POA	E	500	1	-	0/3/4/4	-

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	500	POA	O2-C2	3.65	1.40	1.20
2	A	500	POA	O2-C2	3.54	1.40	1.20
2	B	500	POA	O2-C2	3.41	1.39	1.20
2	F	500	POA	O2-C2	3.35	1.39	1.20
2	C	500	POA	O2-C2	3.30	1.38	1.20

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	500	POA	O3P-P-C1	3.98	116.63	106.78
2	F	500	POA	O3P-P-O1P	-3.98	102.09	112.39
2	B	500	POA	O3P-P-O1P	-3.82	102.51	112.39
2	E	500	POA	O2P-P-C1	3.72	115.98	106.78
2	H	500	POA	O3P-P-O1P	-3.67	102.91	112.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	500	POA	1	0
2	A	500	POA	2	0
2	F	500	POA	2	0
2	D	500	POA	2	0
2	B	500	POA	2	0
2	C	500	POA	3	0
2	G	500	POA	3	0
2	E	500	POA	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	473/488 (96%)	-0.17	13 (2%) 56 58	13, 20, 38, 56	0
1	B	473/488 (96%)	-0.02	13 (2%) 56 58	14, 22, 40, 57	0
1	C	473/488 (96%)	-0.16	9 (1%) 66 67	12, 21, 39, 58	0
1	D	473/488 (96%)	-0.01	23 (4%) 36 38	13, 21, 39, 55	0
1	E	473/488 (96%)	0.09	17 (3%) 46 48	13, 21, 40, 58	0
1	F	473/488 (96%)	-0.15	10 (2%) 63 65	14, 22, 40, 58	0
1	G	473/488 (96%)	-0.01	16 (3%) 48 50	15, 23, 41, 59	0
1	H	473/488 (96%)	0.11	29 (6%) 28 30	14, 22, 41, 57	0
All	All	3784/3904 (96%)	-0.04	130 (3%) 48 50	12, 22, 40, 59	0

The worst 5 of 130 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	12	HIS	7.9
1	H	216	ASP	6.8
1	E	377	HIS	6.3
1	E	12	HIS	6.3
1	D	344	GLU	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	POA	G	500	7/7	0.96	0.07	17,25,31,31	0
2	POA	B	500	7/7	0.98	0.05	14,17,24,26	0
2	POA	C	500	7/7	0.98	0.05	14,16,22,23	0
2	POA	D	500	7/7	0.98	0.06	13,21,26,34	0
2	POA	E	500	7/7	0.98	0.05	18,21,23,27	0
2	POA	F	500	7/7	0.98	0.05	19,21,28,29	0
2	POA	A	500	7/7	0.98	0.05	6,15,17,19	0
2	POA	H	500	7/7	0.98	0.05	18,24,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.