



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 6, 2025 – 12:58 PM JST

PDB ID : 9IPR / pdb\_00009ipr  
Title : Crystal structure of CTB10-M1  
Authors : Fu, K.; Rao, Y.J.  
Deposited on : 2024-07-11  
Resolution : 1.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1306 (1.94-1.94)
Clashscore	180529	1400 (1.94-1.94)
Ramachandran outliers	177936	1387 (1.94-1.94)
Sidechain outliers	177891	1387 (1.94-1.94)
RSRZ outliers	164620	1306 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	<div><div>%</div><div><div></div><div></div><div></div><div></div></div><div>83%6%11%</div></div>
1	B	141	<div><div>%</div><div><div></div><div></div><div></div><div></div></div><div>82%6%12%</div></div>
1	C	141	<div><div>%</div><div><div></div><div></div><div></div><div></div></div><div>87%5%8%</div></div>
1	D	141	<div><div></div><div><div></div><div></div><div></div><div></div></div><div>78%10%12%</div></div>
1	E	141	<div><div>%</div><div><div></div><div></div><div></div><div></div></div><div>80%8%12%</div></div>
1	F	141	<div><div>2%</div><div><div></div><div></div><div></div><div></div></div><div>81%7%12%</div></div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	141	<div><div>4%</div><div>70%</div><div>17%</div><div>13%</div></div>
1	H	141	<div><div>7%</div><div>77%</div><div>11%</div><div>13%</div></div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8849 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CTB10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	0	0	0
			1041	668	171	195	7			
1	B	124	Total	C	N	O	S	0	0	0
			1014	652	167	188	7			
1	C	130	Total	C	N	O	S	0	0	0
			1050	675	172	197	6			
1	D	124	Total	C	N	O	S	0	0	0
			1017	655	165	190	7			
1	E	124	Total	C	N	O	S	0	0	0
			1016	654	168	187	7			
1	F	124	Total	C	N	O	S	0	0	0
			1012	650	166	189	7			
1	G	123	Total	C	N	O	S	0	0	0
			985	631	162	185	7			
1	H	123	Total	C	N	O	S	0	0	0
			980	632	160	181	7			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
A	133	ALA	-	expression tag	UNP A0A977K7H6
A	134	LEU	-	expression tag	UNP A0A977K7H6
A	135	GLU	-	expression tag	UNP A0A977K7H6
A	136	HIS	-	expression tag	UNP A0A977K7H6
A	137	HIS	-	expression tag	UNP A0A977K7H6
A	138	HIS	-	expression tag	UNP A0A977K7H6
A	139	HIS	-	expression tag	UNP A0A977K7H6
A	140	HIS	-	expression tag	UNP A0A977K7H6
A	141	HIS	-	expression tag	UNP A0A977K7H6
B	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
B	133	ALA	-	expression tag	UNP A0A977K7H6
B	134	LEU	-	expression tag	UNP A0A977K7H6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	135	GLU	-	expression tag	UNP A0A977K7H6
B	136	HIS	-	expression tag	UNP A0A977K7H6
B	137	HIS	-	expression tag	UNP A0A977K7H6
B	138	HIS	-	expression tag	UNP A0A977K7H6
B	139	HIS	-	expression tag	UNP A0A977K7H6
B	140	HIS	-	expression tag	UNP A0A977K7H6
B	141	HIS	-	expression tag	UNP A0A977K7H6
C	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
C	133	ALA	-	expression tag	UNP A0A977K7H6
C	134	LEU	-	expression tag	UNP A0A977K7H6
C	135	GLU	-	expression tag	UNP A0A977K7H6
C	136	HIS	-	expression tag	UNP A0A977K7H6
C	137	HIS	-	expression tag	UNP A0A977K7H6
C	138	HIS	-	expression tag	UNP A0A977K7H6
C	139	HIS	-	expression tag	UNP A0A977K7H6
C	140	HIS	-	expression tag	UNP A0A977K7H6
C	141	HIS	-	expression tag	UNP A0A977K7H6
D	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
D	133	ALA	-	expression tag	UNP A0A977K7H6
D	134	LEU	-	expression tag	UNP A0A977K7H6
D	135	GLU	-	expression tag	UNP A0A977K7H6
D	136	HIS	-	expression tag	UNP A0A977K7H6
D	137	HIS	-	expression tag	UNP A0A977K7H6
D	138	HIS	-	expression tag	UNP A0A977K7H6
D	139	HIS	-	expression tag	UNP A0A977K7H6
D	140	HIS	-	expression tag	UNP A0A977K7H6
D	141	HIS	-	expression tag	UNP A0A977K7H6
E	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
E	133	ALA	-	expression tag	UNP A0A977K7H6
E	134	LEU	-	expression tag	UNP A0A977K7H6
E	135	GLU	-	expression tag	UNP A0A977K7H6
E	136	HIS	-	expression tag	UNP A0A977K7H6
E	137	HIS	-	expression tag	UNP A0A977K7H6
E	138	HIS	-	expression tag	UNP A0A977K7H6
E	139	HIS	-	expression tag	UNP A0A977K7H6
E	140	HIS	-	expression tag	UNP A0A977K7H6
E	141	HIS	-	expression tag	UNP A0A977K7H6
F	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
F	133	ALA	-	expression tag	UNP A0A977K7H6
F	134	LEU	-	expression tag	UNP A0A977K7H6
F	135	GLU	-	expression tag	UNP A0A977K7H6
F	136	HIS	-	expression tag	UNP A0A977K7H6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	137	HIS	-	expression tag	UNP A0A977K7H6
F	138	HIS	-	expression tag	UNP A0A977K7H6
F	139	HIS	-	expression tag	UNP A0A977K7H6
F	140	HIS	-	expression tag	UNP A0A977K7H6
F	141	HIS	-	expression tag	UNP A0A977K7H6
G	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
G	133	ALA	-	expression tag	UNP A0A977K7H6
G	134	LEU	-	expression tag	UNP A0A977K7H6
G	135	GLU	-	expression tag	UNP A0A977K7H6
G	136	HIS	-	expression tag	UNP A0A977K7H6
G	137	HIS	-	expression tag	UNP A0A977K7H6
G	138	HIS	-	expression tag	UNP A0A977K7H6
G	139	HIS	-	expression tag	UNP A0A977K7H6
G	140	HIS	-	expression tag	UNP A0A977K7H6
G	141	HIS	-	expression tag	UNP A0A977K7H6
H	13	PBF	TRP	engineered mutation	UNP A0A977K7H6
H	133	ALA	-	expression tag	UNP A0A977K7H6
H	134	LEU	-	expression tag	UNP A0A977K7H6
H	135	GLU	-	expression tag	UNP A0A977K7H6
H	136	HIS	-	expression tag	UNP A0A977K7H6
H	137	HIS	-	expression tag	UNP A0A977K7H6
H	138	HIS	-	expression tag	UNP A0A977K7H6
H	139	HIS	-	expression tag	UNP A0A977K7H6
H	140	HIS	-	expression tag	UNP A0A977K7H6
H	141	HIS	-	expression tag	UNP A0A977K7H6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	135	Total O 135 135	0	0
2	B	95	Total O 95 95	0	0
2	C	103	Total O 103 103	0	0
2	D	96	Total O 96 96	0	0
2	E	101	Total O 101 101	0	0
2	F	95	Total O 95 95	0	0
2	G	56	Total O 56 56	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	53	Total	O	0	0
			53	53		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CTB10




- Molecule 1: CTB10




- Molecule 1: CTB10



- Molecule 1: CTB10




- Molecule 1: CTB10



- Molecule 1: CTB10






● Molecule 1: CTB10



● Molecule 1: CTB10



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.08Å 88.11Å 87.60Å 90.00° 116.66° 90.00°	Depositor
Resolution (Å)	36.26 – 1.94 36.26 – 1.94	Depositor EDS
% Data completeness (in resolution range)	99.0 (36.26-1.94) 98.9 (36.26-1.94)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 1.94Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.191 , 0.226 0.194 , 0.227	Depositor DCC
$R_{free}$ test set	4599 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.6	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 41.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8849	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PBF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/1046	0.66	0/1411
1	B	0.46	0/1018	0.70	1/1374 (0.1%)
1	C	0.54	0/1055	0.76	0/1426
1	D	0.46	0/1021	0.67	0/1378
1	E	0.37	0/1020	0.63	0/1376
1	F	0.45	0/1016	0.76	1/1372 (0.1%)
1	G	0.55	0/987	0.78	0/1333
1	H	0.54	0/983	0.75	0/1331
All	All	0.48	0/8146	0.72	2/11001 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	21	PRO	N-CA-C	6.52	123.47	113.75
1	B	18	THR	CB-CA-C	5.05	118.45	109.72

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1041	0	1015	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1014	0	983	6	0
1	C	1050	0	1010	7	0
1	D	1017	0	988	10	0
1	E	1016	0	988	7	0
1	F	1012	0	971	7	0
1	G	985	0	946	15	0
1	H	980	0	941	10	0
2	A	135	0	0	0	0
2	B	95	0	0	1	0
2	C	103	0	0	0	0
2	D	96	0	0	0	0
2	E	101	0	0	2	0
2	F	95	0	0	1	0
2	G	56	0	0	1	0
2	H	53	0	0	0	0
All	All	8849	0	7842	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:13:PBF:HI1	1:G:15:ILE:HG12	1.77	0.65
1:D:13:PBF:HI2	1:D:15:ILE:HG12	1.79	0.64
1:A:25:GLU:OE2	1:A:62:LYS:HE3	1.98	0.64
1:B:21:PRO:HD3	1:B:116:ASP:HB2	1.85	0.58
1:F:13:PBF:HI1	1:F:15:ILE:HG13	1.87	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	123/141 (87%)	120 (98%)	3 (2%)	0	100	100
1	B	121/141 (86%)	119 (98%)	2 (2%)	0	100	100
1	C	127/141 (90%)	124 (98%)	3 (2%)	0	100	100
1	D	121/141 (86%)	120 (99%)	1 (1%)	0	100	100
1	E	121/141 (86%)	118 (98%)	3 (2%)	0	100	100
1	F	121/141 (86%)	120 (99%)	1 (1%)	0	100	100
1	G	120/141 (85%)	117 (98%)	2 (2%)	1 (1%)	16	7
1	H	120/141 (85%)	118 (98%)	2 (2%)	0	100	100
All	All	974/1128 (86%)	956 (98%)	17 (2%)	1 (0%)	48	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	114	PHE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/127 (91%)	115 (100%)	0	100	100
1	B	110/127 (87%)	110 (100%)	0	100	100
1	C	113/127 (89%)	112 (99%)	1 (1%)	75	72
1	D	111/127 (87%)	111 (100%)	0	100	100
1	E	110/127 (87%)	110 (100%)	0	100	100
1	F	109/127 (86%)	109 (100%)	0	100	100
1	G	105/127 (83%)	103 (98%)	2 (2%)	52	41
1	H	104/127 (82%)	103 (99%)	1 (1%)	73	68
All	All	877/1016 (86%)	873 (100%)	4 (0%)	86	86

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	78	LEU
1	G	23	GLN
1	G	120	SER
1	H	45	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	95	GLN
1	H	95	GLN
1	B	72	GLN
1	F	57	ASN
1	F	95	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	PBF	E	13	1	19,20,21	0.28	0	23,26,28	0.66	0
1	PBF	A	13	1	19,20,21	0.24	0	23,26,28	0.25	0
1	PBF	B	13	1	19,20,21	0.26	0	23,26,28	0.37	0
1	PBF	G	13	1	19,20,21	0.21	0	23,26,28	0.55	0
1	PBF	H	13	1	19,20,21	0.23	0	23,26,28	0.46	0
1	PBF	D	13	1	19,20,21	0.25	0	23,26,28	0.72	0
1	PBF	F	13	1	19,20,21	0.23	0	23,26,28	0.56	0
1	PBF	C	13	1	19,20,21	0.22	0	23,26,28	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PBF	E	13	1	-	2/13/14/16	0/2/2/2
1	PBF	A	13	1	-	2/13/14/16	0/2/2/2
1	PBF	B	13	1	-	2/13/14/16	0/2/2/2
1	PBF	G	13	1	-	2/13/14/16	0/2/2/2
1	PBF	H	13	1	-	6/13/14/16	0/2/2/2
1	PBF	D	13	1	-	2/13/14/16	0/2/2/2
1	PBF	F	13	1	-	6/13/14/16	0/2/2/2
1	PBF	C	13	1	-	2/13/14/16	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	13	PBF	CA-CB-CG-CD1
1	G	13	PBF	CA-CB-CG-CD1
1	G	13	PBF	CA-CB-CG-CD2
1	C	13	PBF	CA-CB-CG-CD2
1	D	13	PBF	CA-CB-CG-CD1

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	G	13	PBF	1	0
1	H	13	PBF	1	0
1	D	13	PBF	1	0
1	F	13	PBF	1	0
1	C	13	PBF	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	125/141 (88%)	-0.20	1 (0%) 82 87	13, 20, 33, 40	0
1	B	123/141 (87%)	-0.11	1 (0%) 82 87	12, 20, 33, 40	0
1	C	129/141 (91%)	-0.00	2 (1%) 70 75	15, 22, 35, 42	0
1	D	123/141 (87%)	-0.02	0 100 100	15, 23, 36, 42	0
1	E	123/141 (87%)	-0.04	1 (0%) 82 87	16, 23, 32, 36	0
1	F	123/141 (87%)	0.00	3 (2%) 59 66	17, 24, 38, 45	0
1	G	122/141 (86%)	0.47	6 (4%) 36 40	20, 32, 44, 54	0
1	H	122/141 (86%)	0.62	10 (8%) 19 22	19, 35, 47, 56	0
All	All	990/1128 (87%)	0.09	24 (2%) 59 66	12, 24, 41, 56	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	118	THR	3.2
1	A	7	PRO	3.1
1	B	112	PHE	3.0
1	G	112	PHE	2.9
1	H	112	PHE	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PBF	G	13	19/20	0.92	0.11	17,25,39,43	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	PBF	C	13	19/20	0.93	0.09	12,17,33,34	0
1	PBF	E	13	19/20	0.94	0.08	14,20,31,35	0
1	PBF	F	13	19/20	0.94	0.08	13,22,33,35	0
1	PBF	D	13	19/20	0.94	0.08	15,21,30,33	0
1	PBF	H	13	19/20	0.94	0.10	18,23,43,44	0
1	PBF	A	13	19/20	0.95	0.08	10,17,33,36	0
1	PBF	B	13	19/20	0.95	0.08	10,18,34,34	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.