



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 29, 2025 – 04:09 PM JST

PDB ID : 8ISC / pdb_00008isc
Title : Crystal structure of MV in complex with LLP
Authors : Li, Q.; Zhu, Y.M.; Gao, J.; Wei, H.L.; Han, X.; Liu, W.D.; Sun, Y.X.
Deposited on : 2023-03-20
Resolution : 2.27 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	8487 (2.30-2.26)
Clashscore	180529	9437 (2.30-2.26)
Ramachandran outliers	177936	9341 (2.30-2.26)
Sidechain outliers	177891	9342 (2.30-2.26)
RSRZ outliers	164620	8487 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>• 5%</div> </div> </div>
1	B	337	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>• 5%</div> </div> </div>
1	C	337	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>• 6%</div> </div> </div>
1	D	337	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>7%</div> <div>• 5%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Branched chain amino acid: 2-keto-4-methylthiobutyrate aminotransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	321	Total	C	N	O	P	S	0	0	0
			2489	1577	422	481	1	8			
1	B	321	Total	C	N	O	P	S	0	0	0
			2489	1577	422	481	1	8			
1	C	318	Total	C	N	O	P	S	0	0	0
			2463	1559	419	476	1	8			
1	D	320	Total	C	N	O	P	S	0	0	0
			2484	1574	421	480	1	8			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	69	LYS	HIS	engineered mutation	UNP A1TDP1
A	105	PRO	SER	engineered mutation	UNP A1TDP1
A	121	MET	SER	engineered mutation	UNP A1TDP1
A	142	PRO	LYS	engineered mutation	UNP A1TDP1
A	145	ARG	LYS	engineered mutation	UNP A1TDP1
A	152	ASN	HIS	engineered mutation	UNP A1TDP1
A	162	ILE	LEU	engineered mutation	UNP A1TDP1
A	168	GLU	ALA	engineered mutation	UNP A1TDP1
A	215	GLY	ARG	engineered mutation	UNP A1TDP1
B	69	LYS	HIS	engineered mutation	UNP A1TDP1
B	105	PRO	SER	engineered mutation	UNP A1TDP1
B	121	MET	SER	engineered mutation	UNP A1TDP1
B	142	PRO	LYS	engineered mutation	UNP A1TDP1
B	145	ARG	LYS	engineered mutation	UNP A1TDP1
B	152	ASN	HIS	engineered mutation	UNP A1TDP1
B	162	ILE	LEU	engineered mutation	UNP A1TDP1
B	168	GLU	ALA	engineered mutation	UNP A1TDP1
B	215	GLY	ARG	engineered mutation	UNP A1TDP1
C	69	LYS	HIS	engineered mutation	UNP A1TDP1
C	105	PRO	SER	engineered mutation	UNP A1TDP1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	121	MET	SER	engineered mutation	UNP A1TDP1
C	142	PRO	LYS	engineered mutation	UNP A1TDP1
C	145	ARG	LYS	engineered mutation	UNP A1TDP1
C	152	ASN	HIS	engineered mutation	UNP A1TDP1
C	162	ILE	LEU	engineered mutation	UNP A1TDP1
C	168	GLU	ALA	engineered mutation	UNP A1TDP1
C	215	GLY	ARG	engineered mutation	UNP A1TDP1
D	69	LYS	HIS	engineered mutation	UNP A1TDP1
D	105	PRO	SER	engineered mutation	UNP A1TDP1
D	121	MET	SER	engineered mutation	UNP A1TDP1
D	142	PRO	LYS	engineered mutation	UNP A1TDP1
D	145	ARG	LYS	engineered mutation	UNP A1TDP1
D	152	ASN	HIS	engineered mutation	UNP A1TDP1
D	162	ILE	LEU	engineered mutation	UNP A1TDP1
D	168	GLU	ALA	engineered mutation	UNP A1TDP1
D	215	GLY	ARG	engineered mutation	UNP A1TDP1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	307	Total O 307 307	0	0
2	B	299	Total O 299 299	0	0
2	C	286	Total O 286 286	0	0
2	D	264	Total O 264 264	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Branched chain amino acid: 2-keto-4-methylthiobutyrat aminotransferase




- Molecule 1: Branched chain amino acid: 2-keto-4-methylthiobutyrat aminotransferase



- Molecule 1: Branched chain amino acid: 2-keto-4-methylthiobutyrat aminotransferase



- Molecule 1: Branched chain amino acid: 2-keto-4-methylthiobutyrat aminotransferase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.95Å 112.85Å 92.24Å 90.00° 112.37° 90.00°	Depositor
Resolution (Å)	39.93 – 2.27 39.93 – 2.27	Depositor EDS
% Data completeness (in resolution range)	94.1 (39.93-2.27) 94.1 (39.93-2.27)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.190 , 0.248 0.197 , 0.252	Depositor DCC
R_{free} test set	3335 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11081	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	0/2521	1.40	1/3430 (0.0%)
1	B	1.08	0/2521	1.36	3/3430 (0.1%)
1	C	1.05	0/2494	1.41	5/3394 (0.1%)
1	D	1.06	0/2516	1.38	2/3423 (0.1%)
All	All	1.06	0/10052	1.39	11/13677 (0.1%)

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	78	HIS	CB-CA-C	5.79	119.35	109.80
1	C	263	GLY	CA-C-N	5.59	127.77	120.28
1	C	263	GLY	C-N-CA	5.59	127.77	120.28
1	C	85	PHE	CA-C-N	5.32	129.67	122.07
1	C	85	PHE	C-N-CA	5.32	129.67	122.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2489	0	2429	18	0
1	B	2489	0	2429	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2463	0	2404	23	0
1	D	2484	0	2424	12	0
2	A	307	0	0	2	0
2	B	299	0	0	3	0
2	C	286	0	0	9	0
2	D	264	0	0	3	0
All	All	11081	0	9686	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:299:THR:HG22	2:D:503:HOH:O	1.78	0.82
1:B:220:MET:HE2	1:B:276:SER:HB3	1.64	0.80
1:C:299:THR:HG22	2:C:551:HOH:O	1.89	0.72
1:D:18:ILE:N	2:D:401:HOH:O	2.22	0.72
1:B:220:MET:SD	2:B:691:HOH:O	2.47	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/337 (94%)	306 (96%)	12 (4%)	0	100	100
1	B	318/337 (94%)	308 (97%)	10 (3%)	0	100	100
1	C	315/337 (94%)	300 (95%)	15 (5%)	0	100	100
1	D	317/337 (94%)	305 (96%)	12 (4%)	0	100	100
All	All	1268/1348 (94%)	1219 (96%)	49 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/269 (96%)	250 (97%)	7 (3%)	40	54
1	B	257/269 (96%)	249 (97%)	8 (3%)	35	49
1	C	255/269 (95%)	250 (98%)	5 (2%)	50	65
1	D	257/269 (96%)	249 (97%)	8 (3%)	35	49
All	All	1026/1076 (95%)	998 (97%)	28 (3%)	40	54

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	317	ILE
1	D	300	LEU
1	C	78	HIS
1	D	182	VAL
1	C	70	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	298	ASN
1	C	298	ASN
1	D	298	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	195	1	23,24,25	1.26	3 (13%)	25,32,34	1.11	3 (12%)
1	LLP	C	195	1	23,24,25	1.12	2 (8%)	25,32,34	1.10	1 (4%)
1	LLP	B	195	1	23,24,25	1.11	3 (13%)	25,32,34	1.39	4 (16%)
1	LLP	D	195	1	23,24,25	1.40	3 (13%)	25,32,34	1.38	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	195	1	-	5/16/17/19	0/1/1/1
1	LLP	C	195	1	-	3/16/17/19	0/1/1/1
1	LLP	B	195	1	-	3/16/17/19	0/1/1/1
1	LLP	D	195	1	-	5/16/17/19	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	195	LLP	C4-C4'	3.41	1.53	1.46
1	A	195	LLP	CD-CE	3.34	1.63	1.51
1	B	195	LLP	CD-CE	3.16	1.62	1.51
1	D	195	LLP	CD-CE	3.10	1.62	1.51
1	C	195	LLP	CD-CE	3.10	1.62	1.51

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	195	LLP	C3-C4-C5	-3.45	115.61	118.26
1	D	195	LLP	OP4-C5'-C5	3.12	115.29	109.35
1	B	195	LLP	C3-C4-C5	-3.03	115.93	118.26
1	B	195	LLP	C5-C4-C4'	2.70	126.01	121.56
1	A	195	LLP	CD-CG-CB	2.62	122.89	113.62

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	195	LLP	C4-C4'-NZ-CE
1	A	195	LLP	N-CA-CB-CG
1	A	195	LLP	C-CA-CB-CG
1	B	195	LLP	C4-C4'-NZ-CE
1	C	195	LLP	C4-C4'-NZ-CE

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	195	LLP	1	0
1	B	195	LLP	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/337 (94%)	-0.16	6 (1%) 66 67	10, 16, 34, 66	0
1	B	320/337 (94%)	-0.15	2 (0%) 85 86	10, 17, 34, 69	0
1	C	317/337 (94%)	-0.09	5 (1%) 70 71	11, 18, 35, 58	0
1	D	319/337 (94%)	-0.06	2 (0%) 85 86	12, 20, 37, 60	0
All	All	1276/1348 (94%)	-0.12	15 (1%) 76 77	10, 18, 35, 69	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	330	PRO	5.0
1	B	145	ARG	4.7
1	D	18	ILE	4.3
1	A	145	ARG	3.6
1	D	145	ARG	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	195	24/25	0.92	0.11	13,20,25,27	0
1	LLP	B	195	24/25	0.92	0.10	15,19,22,23	0
1	LLP	D	195	24/25	0.92	0.11	14,19,26,26	0
1	LLP	C	195	24/25	0.94	0.10	13,21,25,26	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.