



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2024 – 10:01 pm BST

PDB ID : 2IYC
Title : SENP1 native structure
Authors : Dong, C.; Naismith, J.H.
Deposited on : 2006-07-14
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	<div> <div>5%</div> <div>77%</div> <div>20%</div> <div>.</div> </div>
1	B	226	<div> <div>15%</div> <div>72%</div> <div>23%</div> <div>.</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SENTRIN-SPECIFIC PROTEASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1891	1213	327	337	14			
1	B	226	Total	C	N	O	S	0	0	0
			1891	1213	327	337	14			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	49	Total	O	0	0
			49	49		
2	B	41	Total	O	0	0
			41	41		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SENTRIN-SPECIFIC PROTEASE 1



• Molecule 1: SENTRIN-SPECIFIC PROTEASE 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	71.98Å 71.98Å 200.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	53.00 – 2.45 45.60 – 2.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (53.00-2.45) 100.0 (45.60-2.45)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.217 , 0.267 0.231 , 0.279	Depositor DCC
R_{free} test set	1136 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	52.2	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 63.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3872	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	2/1934 (0.1%)	0.74	1/2599 (0.0%)
1	B	0.84	8/1934 (0.4%)	0.72	3/2599 (0.1%)
All	All	0.79	10/3868 (0.3%)	0.73	4/5198 (0.1%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	571	SER	CB-OG	11.31	1.56	1.42
1	B	430	GLU	CB-CG	10.41	1.72	1.52
1	B	561	ARG	CZ-NH1	8.91	1.44	1.33
1	A	430	GLU	CB-CG	8.17	1.67	1.52
1	B	588	SER	CB-OG	7.46	1.51	1.42
1	B	592	GLN	CD-NE2	6.48	1.49	1.32
1	B	425	GLU	CG-CD	6.44	1.61	1.51
1	A	430	GLU	CD-OE2	6.07	1.32	1.25
1	B	430	GLU	CD-OE1	5.82	1.32	1.25
1	B	577	LYS	CE-NZ	5.03	1.61	1.49

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	430	GLU	CA-CB-CG	6.70	128.15	113.40
1	B	430	GLU	CG-CD-OE1	5.67	129.65	118.30
1	B	587	PHE	CB-CA-C	-5.08	100.25	110.40
1	A	430	GLU	CG-CD-OE2	5.03	128.36	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1891	0	1890	33	2
1	B	1891	0	1890	47	2
2	A	49	0	0	3	0
2	B	41	0	0	9	0
All	All	3872	0	3780	79	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:482:SER:O	1:A:483:LYS:HB3	1.40	1.16
1:A:485:LYS:HE3	1:A:486:GLY:H	1.10	1.09
1:B:482:SER:O	1:B:483:LYS:HB3	1.47	1.08
1:B:449:ARG:HG3	1:B:449:ARG:NH1	1.62	1.03
1:B:449:ARG:HG3	1:B:449:ARG:HH11	0.87	1.03
1:B:449:ARG:HH11	1:B:449:ARG:CG	1.78	0.95
1:B:430:GLU:OE1	1:B:640:HIS:NE2	2.04	0.88
1:A:623:THR:H	1:A:626:HIS:HD2	1.17	0.88
1:B:623:THR:H	1:B:626:HIS:HD2	1.20	0.87
1:A:623:THR:H	1:A:626:HIS:CD2	1.96	0.82
1:A:448:PHE:O	1:A:449:ARG:HB2	1.77	0.82
1:B:482:SER:O	1:B:483:LYS:CB	2.28	0.76
1:A:485:LYS:HE3	1:A:486:GLY:N	1.95	0.76
1:B:466:LEU:HD21	1:B:605:MET:HE2	1.68	0.75
1:A:482:SER:O	1:A:483:LYS:CB	2.24	0.74
1:A:499:THR:O	1:A:503:THR:HG23	1.95	0.67
1:B:566:TYR:HA	1:B:569:GLN:HE21	1.60	0.65
1:A:430:GLU:OE2	1:A:640:HIS:NE2	2.26	0.65
1:B:623:THR:H	1:B:626:HIS:CD2	2.09	0.65
1:A:433:ASN:HA	1:A:436:ARG:HD2	1.80	0.64
1:B:514:LYS:HA	2:B:2021:HOH:O	1.99	0.61
1:A:442:GLU:OE1	1:A:454:ARG:NH1	2.34	0.60
1:B:469:GLU:HG2	2:B:2013:HOH:O	2.00	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:458:GLN:HB2	2:B:2009:HOH:O	2.01	0.59
1:B:472:ASN:OD1	1:B:495:THR:HG23	2.02	0.59
1:B:590:LYS:H	1:B:590:LYS:HE2	1.67	0.58
1:B:458:GLN:OE1	1:B:461:ASN:HB2	2.02	0.58
1:A:598:MET:HB2	1:A:624:GLN:HG2	1.84	0.57
1:B:557:ASN:OD1	1:B:557:ASN:N	2.38	0.57
1:B:539:VAL:HG22	1:B:546:ILE:HG12	1.87	0.56
1:A:552:MET:HE1	1:B:530:LEU:HB3	1.89	0.55
1:B:434:VAL:CG2	1:B:444:LEU:HD13	2.37	0.55
1:B:558:GLU:HB3	2:B:2027:HOH:O	2.06	0.55
1:B:543:LYS:HB3	1:B:545:ASN:ND2	2.22	0.54
1:B:482:SER:HB2	1:B:488:PRO:O	2.09	0.52
1:A:437:ASN:ND2	1:A:437:ASN:H	2.08	0.52
1:A:477:MET:HE2	1:A:638:ILE:HG12	1.92	0.51
1:A:477:MET:CE	1:A:638:ILE:HG12	2.41	0.51
1:B:433:ASN:O	1:B:436:ARG:HG3	2.12	0.51
1:A:623:THR:N	1:A:626:HIS:HD2	1.98	0.50
1:A:483:LYS:HA	2:A:2013:HOH:O	2.11	0.50
1:B:544:LYS:O	1:B:584:TRP:HA	2.12	0.50
1:A:472:ASN:OD1	1:A:495:THR:HG23	2.12	0.49
1:A:564:LEU:HD11	1:A:581:THR:HG22	1.94	0.49
1:A:511:ARG:O	1:A:514:LYS:HG2	2.13	0.48
1:A:419:GLU:HG2	2:A:2001:HOH:O	2.13	0.48
1:A:459:THR:HB	1:A:465:TRP:O	2.14	0.48
1:B:462:HIS:CA	2:B:2010:HOH:O	2.62	0.48
1:A:544:LYS:O	1:A:584:TRP:HA	2.13	0.47
1:A:545:ASN:OD1	1:A:545:ASN:C	2.53	0.47
1:B:480:GLU:HG3	2:B:2017:HOH:O	2.14	0.47
1:B:435:PHE:CZ	1:B:635:VAL:HG21	2.50	0.47
1:B:590:LYS:HG2	1:B:593:GLU:HB2	1.95	0.47
1:B:458:GLN:OE1	1:B:461:ASN:CB	2.63	0.46
1:B:507:GLN:HG2	2:B:2020:HOH:O	2.14	0.46
1:B:589:LYS:HE3	1:B:593:GLU:HG2	1.98	0.46
1:B:566:TYR:HA	1:B:569:GLN:NE2	2.30	0.45
1:B:432:LYS:NZ	2:B:2006:HOH:O	2.50	0.45
1:A:439:ASN:HB3	1:A:442:GLU:HB2	1.99	0.45
1:B:545:ASN:HB2	1:B:585:GLN:O	2.15	0.45
1:B:463:LEU:H	1:B:463:LEU:HG	1.59	0.45
1:B:460:LEU:CD2	1:B:605:MET:HE1	2.47	0.45
1:B:449:ARG:NH1	1:B:449:ARG:CG	2.48	0.44
1:B:599:ASN:OD1	2:B:2030:HOH:O	2.20	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:459:THR:HB	1:B:465:TRP:O	2.16	0.44
1:B:589:LYS:HG3	1:B:593:GLU:HG2	2.00	0.44
1:A:431:ILE:HD11	1:A:636:TRP:HB2	1.99	0.44
1:A:482:SER:HB2	1:A:488:PRO:O	2.17	0.44
1:B:451:THR:O	1:B:451:THR:CG2	2.66	0.43
1:B:518:VAL:HG22	1:B:524:LEU:CD2	2.49	0.43
1:B:430:GLU:OE1	1:B:640:HIS:CE1	2.69	0.42
1:A:481:ARG:O	1:A:484:GLU:HG2	2.19	0.42
1:B:590:LYS:H	1:B:590:LYS:CE	2.30	0.42
1:B:518:VAL:HG22	1:B:524:LEU:HD22	2.02	0.41
1:A:618:ARG:HA	1:A:618:ARG:HD3	1.92	0.41
1:B:590:LYS:HE2	1:B:590:LYS:N	2.33	0.41
1:A:641:ARG:NE	2:A:2046:HOH:O	2.54	0.41
1:A:495:THR:HA	1:A:527:PRO:HG2	2.03	0.41
1:A:575:LYS:O	1:A:577:LYS:HG3	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:430:GLU:OE2	1:B:430:GLU:OE1[3_654]	1.31	0.89
1:A:640:HIS:NE2	1:B:430:GLU:OE1[3_654]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/226 (99%)	213 (95%)	9 (4%)	2 (1%)	17	19
1	B	224/226 (99%)	215 (96%)	7 (3%)	2 (1%)	17	19
All	All	448/452 (99%)	428 (96%)	16 (4%)	4 (1%)	17	19

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	439	ASN
1	B	439	ASN
1	A	483	LYS
1	B	483	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	196 (94%)	13 (6%)	18	23
1	B	209/209 (100%)	193 (92%)	16 (8%)	13	15
All	All	418/418 (100%)	389 (93%)	29 (7%)	15	18

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	432	LYS
1	A	437	ASN
1	A	445	SER
1	A	485	LYS
1	A	494	ASN
1	A	499	THR
1	A	511	ARG
1	A	514	LYS
1	A	518	VAL
1	A	557	ASN
1	A	558	GLU
1	A	585	GLN
1	A	616	LYS
1	B	430	GLU
1	B	433	ASN
1	B	434	VAL
1	B	437	ASN
1	B	449	ARG
1	B	451	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	463	LEU
1	B	494	ASN
1	B	502	LYS
1	B	518	VAL
1	B	555	ILE
1	B	557	ASN
1	B	558	GLU
1	B	590	LYS
1	B	621	ASN
1	B	641	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	437	ASN
1	A	569	GLN
1	A	599	ASN
1	A	626	HIS
1	B	433	ASN
1	B	437	ASN
1	B	569	GLN
1	B	626	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/226 (100%)	0.56	12 (5%) 26 23	19, 29, 33, 48	0
1	B	226/226 (100%)	0.88	35 (15%) 2 1	18, 29, 33, 38	0
All	All	452/452 (100%)	0.72	47 (10%) 6 4	18, 29, 34, 48	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	437	ASN	7.4
1	A	511	ARG	6.2
1	B	579	PHE	6.1
1	A	576	ARG	5.8
1	B	576	ARG	5.6
1	B	436	ARG	5.5
1	B	555	ILE	5.3
1	B	621	ASN	4.9
1	B	583	GLY	4.5
1	B	587	PHE	4.2
1	B	582	ASN	4.1
1	B	541	PHE	3.8
1	B	575	LYS	3.6
1	B	438	GLY	3.5
1	B	462	HIS	3.5
1	A	437	ASN	3.2
1	A	577	LYS	3.2
1	B	580	ASP	3.1
1	B	435	PHE	3.0
1	A	515	LYS	3.0
1	B	483	LYS	3.0
1	A	516	VAL	2.9
1	B	571	SER	2.9
1	B	590	LYS	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	578	GLU	2.8
1	B	439	ASN	2.8
1	B	586	LEU	2.8
1	B	521	VAL	2.7
1	B	547	THR	2.6
1	B	441	ASP	2.6
1	A	514	LYS	2.6
1	B	540	ASP	2.5
1	B	625	GLN	2.4
1	B	572	ILE	2.4
1	B	520	SER	2.4
1	B	519	PHE	2.4
1	A	572	ILE	2.3
1	B	563	LEU	2.3
1	B	577	LYS	2.3
1	B	515	LYS	2.3
1	A	433	ASN	2.2
1	B	584	TRP	2.2
1	A	575	LYS	2.2
1	B	585	GLN	2.2
1	A	535	CYS	2.2
1	B	594	ILE	2.1
1	A	419	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.