



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2025 – 12:09 PM JST

PDB ID : 9J19  
Title : The crystal structure of COVID-19 main protease in complex with an inhibitor minocycline  
Authors : Singh, A.; Jangid, K.; Dhaka, P.; Tomar, S.; Kumar, P.  
Deposited on : 2024-08-04  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.2


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	<div> <div>6%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>..</div> </div> </div>
1	B	296	<div> <div>27%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>..</div> </div> </div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3C-like proteinase nsp5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	1	0
			2280	1444	388	427	21			
1	B	290	Total	C	N	O	S	0	0	0
			2255	1428	384	422	21			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ARG	-	expression tag	UNP P0DTD1
B	4	ARG	-	expression tag	UNP P0DTD1

- Molecule 2 is (4S,4AS,5AR,12AS)-4,7-BIS(DIMETHYLAMINO)-3,10,12,12A-TETRAHYDROXY-1,11-DIOXO-1,4,4A,5,5A,6,11,12A-OCTAHYDROTETRACENE-2-CARBOXAMIDE (three-letter code: MIY) (formula:  $C_{23}H_{27}N_3O_7$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			33	23	3	7		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 3C-like proteinase nsp5



#### • Molecule 1: 3C-like proteinase nsp5



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.94Å 90.76Å 101.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.17 – 2.70 23.17 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.1 (23.17-2.70) 99.1 (23.17-2.70)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.68 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.8.0403	Depositor
R, $R_{free}$	0.250 , 0.314 0.259 , 0.325	Depositor DCC
$R_{free}$ test set	885 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.4	Xtriage
Anisotropy	0.308	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 35.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	4568	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MIY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/2330	0.71	1/3166 (0.0%)
1	B	0.33	0/2304	0.68	0/3132
All	All	0.35	0/4634	0.70	1/6298 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178	GLU	CB-CA-C	-5.04	100.32	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	60	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2280	0	2229	23	0
1	B	2255	0	2210	17	0
2	A	33	0	27	3	0
All	All	4568	0	4466	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:LEU:HD22	1:A:145:CYS:HB2	1.66	0.79
1:B:145:CYS:SG	1:B:164:HIS:O	2.45	0.73
1:B:261:VAL:HG13	1:B:262:LEU:HD22	1.71	0.72
2:A:401:MIY:H712	2:A:401:MIY:H82	1.74	0.68
1:A:32:LEU:HD13	1:A:101:TYR:CE2	2.32	0.65
1:B:225:THR:HG21	1:B:269:LYS:HD3	1.80	0.62
1:B:32:LEU:HD13	1:B:101:TYR:CE2	2.36	0.60
1:B:31:TRP:CE2	1:B:95:ASN:HB2	2.38	0.59
1:A:31:TRP:CE2	1:A:95:ASN:HB2	2.37	0.59
1:A:27:LEU:CD2	1:A:145:CYS:HB2	2.32	0.58
1:A:225:THR:O	1:A:262:LEU:HD13	2.03	0.58
1:B:233:VAL:O	1:B:236:LYS:HB3	2.02	0.58
2:A:401:MIY:H712	2:A:401:MIY:C8	2.33	0.57
1:A:56:ASP:O	1:A:60:ARG:HG3	2.07	0.55
1:B:55:GLU:O	1:B:59:ILE:HG13	2.08	0.54
1:A:69:GLN:HE21	1:A:72:ASN:HA	1.73	0.53
1:A:229:ASP:O	1:A:233:VAL:HG23	2.08	0.53
2:A:401:MIY:H203	2:A:401:MIY:O2	2.11	0.50
1:A:86:VAL:HG13	1:A:179:GLY:HA3	1.95	0.48
1:A:140:PHE:HB3	1:A:144:SER:HB2	1.96	0.47
1:A:72:ASN:N	1:A:72:ASN:OD1	2.47	0.47
1:B:86:VAL:HG13	1:B:179:GLY:CA	2.45	0.47
1:A:190:THR:O	1:A:192:GLN:HG3	2.16	0.47
1:B:10:SER:OG	1:B:14:GLU:OE2	2.29	0.46
1:A:212:VAL:HA	1:A:216:ASP:O	2.16	0.46
1:B:225:THR:HG21	1:B:269:LYS:CD	2.46	0.46
1:B:31:TRP:CD2	1:B:95:ASN:HB2	2.51	0.45
1:B:213:ILE:HD13	1:B:257:THR:HG22	1.98	0.45
1:A:19:GLN:NE2	1:A:119:ASN:HB3	2.31	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:SER:OG	1:A:14:GLU:OE2	2.28	0.45
1:B:237:TYR:O	1:B:238:ASN:HB3	2.17	0.45
1:A:233:VAL:HG21	1:A:269:LYS:HE3	1.99	0.44
1:B:49:MET:HB3	1:B:189:GLN:HB2	1.99	0.43
1:B:190:THR:O	1:B:192:GLN:HG3	2.17	0.43
1:A:31:TRP:CD2	1:A:95:ASN:HB2	2.54	0.43
1:A:86:VAL:CG1	1:A:179:GLY:CA	2.97	0.43
1:A:86:VAL:HG13	1:A:179:GLY:CA	2.49	0.42
1:B:51:ASN:OD1	1:B:51:ASN:N	2.54	0.41
1:B:27:LEU:HB2	1:B:145:CYS:O	2.21	0.41
1:A:108:PRO:HA	1:A:130:MET:CG	2.51	0.40
1:A:263:ASP:O	1:A:266:ALA:HB3	2.21	0.40
1:A:145:CYS:HA	1:A:163:HIS:HD2	1.86	0.40
1:A:230:PHE:CD1	1:A:265:CYS:HB3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/296 (98%)	273 (94%)	16 (6%)	1 (0%)	37	61
1	B	286/296 (97%)	265 (93%)	19 (7%)	2 (1%)	19	42
All	All	576/592 (97%)	538 (93%)	35 (6%)	3 (0%)	25	49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	TYR
1	B	71	GLY
1	B	154	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/255 (99%)	244 (96%)	9 (4%)	30	59
1	B	251/255 (98%)	237 (94%)	14 (6%)	17	41
All	All	504/510 (99%)	481 (95%)	23 (5%)	23	49

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	THR
1	A	59	ILE
1	A	144	SER
1	A	153	ASP
1	A	214	ASN
1	A	222	ARG
1	A	236	LYS
1	A	267	SER
1	A	291	PHE
1	B	24	THR
1	B	27	LEU
1	B	49	MET
1	B	72	ASN
1	B	90	LYS
1	B	154	TYR
1	B	216	ASP
1	B	227	LEU
1	B	228	ASN
1	B	229	ASP
1	B	254	SER
1	B	256	GLN
1	B	267	SER
1	B	299	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	69	GLN
1	A	107	GLN
1	A	163	HIS
1	A	180	ASN
1	A	273	GLN
1	A	274	ASN
1	A	277	ASN
1	B	19	GLN
1	B	69	GLN
1	B	163	HIS
1	B	172	HIS
1	B	180	ASN
1	B	256	GLN
1	B	277	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MIY	A	401	-	35,36,36	1.24	2 (5%)	41,58,58	1.73	9 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MIY	A	401	-	-	1/12/70/70	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	MIY	C21-N2	5.52	1.48	1.33
2	A	401	MIY	O5-C15	2.35	1.28	1.23

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	MIY	C7-C6-C5	3.94	117.42	110.49
2	A	401	MIY	C15-C16-C17	3.60	121.65	118.80
2	A	401	MIY	C1-C18-C17	3.32	113.77	109.88
2	A	401	MIY	C18-C17-C16	-3.26	119.76	123.06
2	A	401	MIY	O7-C18-C17	-3.02	105.31	110.14
2	A	401	MIY	O6-C17-C18	2.90	117.57	113.37
2	A	401	MIY	C18-C1-C2	2.79	120.19	115.75
2	A	401	MIY	C71-N7-CN7	-2.40	108.38	116.12
2	A	401	MIY	C8-C9-C10	-2.11	118.43	123.52

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	MIY	C3-C2-C21-N2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	MIY	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	293/296 (98%)	0.56	17 (5%) 30 28	13, 26, 53, 82	1 (0%)
1	B	290/296 (97%)	1.30	81 (27%) 2 2	14, 40, 95, 111	0
All	All	583/592 (98%)	0.93	98 (16%) 5 5	13, 30, 88, 111	1 (0%)

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	140	PHE	6.4
1	B	224	THR	5.8
1	B	242	LEU	5.7
1	B	233	VAL	5.4
1	B	257	THR	5.2
1	B	236	LYS	5.2
1	B	51	ASN	5.0
1	B	231	ASN	5.0
1	B	232	LEU	4.8
1	B	249	ILE	4.7
1	B	246	HIS	4.4
1	B	256	GLN	4.3
1	B	230	PHE	4.3
1	B	223	PHE	4.1
1	B	243	THR	4.0
1	A	139	SER	4.0
1	B	227	LEU	3.9
1	B	118	TYR	3.8
1	B	294	PHE	3.8
1	B	263	ASP	3.8
1	B	258	GLY	3.7
1	B	165	MET	3.7
1	B	213	ILE	3.7
1	B	154	TYR	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	283	GLY	3.6
1	B	64	HIS	3.6
1	B	219	PHE	3.6
1	B	279	ARG	3.5
1	B	244	GLN	3.5
1	B	255	ALA	3.5
1	B	259	ILE	3.5
1	B	171	VAL	3.5
1	A	118	TYR	3.4
1	B	235	MET	3.4
1	B	194	ALA	3.4
1	B	234	ALA	3.3
1	B	247	VAL	3.3
1	B	261	VAL	3.3
1	B	298	ARG	3.3
1	B	218	TRP	3.3
1	A	195	GLY	3.3
1	B	237	TYR	3.3
1	A	67	LEU	3.3
1	B	42	VAL	3.1
1	B	241	PRO	3.1
1	B	297	VAL	3.1
1	A	154	TYR	3.1
1	B	50	LEU	3.1
1	B	273	GLN	3.0
1	A	217[A]	ARG	3.0
1	A	120	GLY	3.0
1	B	226	THR	2.9
1	B	46	SER	2.9
1	B	271	LEU	2.9
1	B	217	ARG	2.8
1	B	26	THR	2.8
1	A	194	ALA	2.8
1	B	299	GLN	2.8
1	B	60	ARG	2.8
1	B	220	LEU	2.8
1	B	59	ILE	2.7
1	B	266	ALA	2.7
1	B	193	ALA	2.7
1	B	206	ALA	2.7
1	B	196	THR	2.7
1	B	252	PRO	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	262	LEU	2.7
1	B	49	MET	2.6
1	B	168	PRO	2.5
1	B	222	ARG	2.5
1	B	144	SER	2.5
1	B	253	LEU	2.5
1	B	265	CYS	2.5
1	B	65	ASN	2.5
1	A	46	SER	2.4
1	B	296	VAL	2.4
1	B	191	ALA	2.3
1	A	45	THR	2.3
1	B	208	LEU	2.3
1	A	247	VAL	2.3
1	B	47	GLU	2.3
1	B	268	LEU	2.2
1	B	225	THR	2.2
1	A	119	ASN	2.2
1	B	172	HIS	2.2
1	B	260	ALA	2.2
1	A	214	ASN	2.2
1	B	285	ALA	2.2
1	B	281	ILE	2.2
1	B	250	LEU	2.2
1	B	216	ASP	2.1
1	A	110	GLN	2.1
1	A	170	GLY	2.1
1	A	298	ARG	2.1
1	B	245	ASP	2.1
1	B	282	LEU	2.0
1	B	52	PRO	2.0
1	B	280	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.




## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MIY	A	401	33/33	0.59	0.33	49,67,79,81	33

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.