



wwPDB EM Validation Summary Report ⓘ

Nov 5, 2024 – 10:30 AM JST

PDB ID : 8JHO
EMDB ID : EMD-36283
Title : Cryo-EM structure of the histone deacetylase complex Rpd3S in complex with di-nucleosome
Authors : Wang, H.
Deposited on : 2023-05-25
Resolution : 7.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis	: FAILED
Mogul	: 1.8.5 (274361), CSD as541be (2020)
MolProbity	: 4.02b-467
Percentile statistics	: 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	: FAILED
Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 7.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	135	73% 24%
1	E	135	81% 13%
1	a	135	76% 24%
1	e	135	70% 27%
2	B	102	77% 21%
2	F	102	73% 5% 22%
2	b	102	80% 20%
2	f	102	78% 22%
3	C	129	79% 5% 16%
3	G	129	81% 18%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	c	129	 84% 16%
3	g	129	 82% 18%
4	D	122	 75% 21%
4	H	122	 78% 22%
4	d	122	 78% 21%
4	h	122	 76% 22%
5	I	350	 60% 33% . .
6	J	350	 61% 32% . .
7	K	1536	 36% 64%
8	L	433	 88% 11%
9	M	401	 72% 27%
9	O	401	 66% 33%
10	N	684	 54% 46%
10	P	684	 22% 78%

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 42813 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	102	Total	C	N	O	S	0	0
			837	529	162	143	3		
1	E	117	Total	C	N	O	S	0	0
			954	597	191	164	2		
1	a	102	Total	C	N	O	S	0	0
			837	529	162	143	3		
1	e	98	Total	C	N	O	S	0	0
			810	512	157	139	2		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	ALA	CYS	engineered mutation	UNP A0A310TTQ1
E	110	ALA	CYS	engineered mutation	UNP A0A310TTQ1
a	110	ALA	CYS	engineered mutation	UNP A0A310TTQ1
e	110	ALA	CYS	engineered mutation	UNP A0A310TTQ1

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	81	Total	C	N	O	S	0	0
			646	407	126	112	1		
2	F	80	Total	C	N	O	S	0	0
			638	401	125	111	1		
2	b	82	Total	C	N	O	S	0	0
			653	412	127	113	1		
2	f	80	Total	C	N	O	S	0	0
			638	401	125	111	1		

- Molecule 3 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	109	Total	C	N	O	0	0
			843	531	167	145		
3	G	106	Total	C	N	O	0	0
			818	516	160	142		
3	c	109	Total	C	N	O	0	0
			843	531	167	145		
3	g	106	Total	C	N	O	0	0
			818	516	160	142		

- Molecule 4 is a protein called Histone H2B.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	96	Total	C	N	O	S	0	0
			757	475	140	140	2		
4	H	95	Total	C	N	O	S	0	0
			745	469	134	140	2		
4	d	96	Total	C	N	O	S	0	0
			757	475	140	140	2		
4	h	95	Total	C	N	O	S	0	0
			745	469	134	140	2		

- Molecule 5 is a DNA chain called Di-nucleosome template forward.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	340	Total	C	N	O	P	0	0
			6937	3299	1252	2046	340		

- Molecule 6 is a DNA chain called Di-nucleosome template reverse.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	340	Total	C	N	O	P	0	0
			7003	3319	1310	2034	340		

- Molecule 7 is a protein called Transcriptional regulatory protein SIN3.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	K	549	Total	C	N	O	S	0	0
			4597	2954	774	854	15		

- Molecule 8 is a protein called Histone deacetylase RPD3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	L	384	Total	C	N	O	S	0	0
			3048	1941	512	569	26		

- Molecule 9 is a protein called Chromatin modification-related protein EAF3.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	M	294	Total	C	N	O	S	0	0
			2398	1541	394	449	14		
9	O	267	Total	C	N	O	S	0	0
			2190	1414	359	404	13		

- Molecule 10 is a protein called RCO1 isoform 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	N	372	Total	C	N	O	S	0	0
			3045	1935	526	566	18		
10	P	151	Total	C	N	O	S	0	0
			1249	802	206	231	10		

- Molecule 11 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
11	L	1	Total	Zn	0
			1	1	
11	N	4	Total	Zn	0
			4	4	
11	P	2	Total	Zn	0
			2	2	

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3

Chain A:  73% 24%




- Molecule 1: Histone H3

Chain E:  81% 13%




- Molecule 1: Histone H3

Chain a:  76% 24%





- Molecule 1: Histone H3

Chain e:  70% 27%




- Molecule 2: Histone H4

Chain B:  77% 21%




- Molecule 2: Histone H4

Chain F:  73% 5% 22%




- Molecule 2: Histone H4

Chain b: 80% 20%



- Molecule 2: Histone H4

Chain f: 78% 22%



- Molecule 3: Histone H2A

Chain C: 79% 5% 16%




- Molecule 3: Histone H2A

Chain G: 81% 0% 18%




- Molecule 3: Histone H2A

Chain c: 84% 16%



- Molecule 3: Histone H2A

Chain g: 82% 18%



- Molecule 4: Histone H2B

Chain D: 75% 0% 21%

- Molecule 9: Chromatin modification-related protein EAF3

- Molecule 9: Chromatin modification-related protein EAF3

- Molecule 10: RCO1 isoform 1



GLN
SER
GLU

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	31310	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	44	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ML3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/836	1.03	3/1120 (0.3%)
1	E	0.60	0/966	1.13	7/1291 (0.5%)
1	a	0.34	0/836	0.80	0/1120
1	e	0.43	0/822	0.86	3/1103 (0.3%)
2	B	0.54	0/653	0.95	3/873 (0.3%)
2	F	0.61	0/645	1.33	8/862 (0.9%)
2	b	0.34	0/660	0.80	0/883
2	f	0.40	0/645	0.83	0/862
3	C	0.66	0/853	1.25	5/1149 (0.4%)
3	G	0.44	0/828	0.77	0/1117
3	c	0.37	0/853	0.76	0/1149
3	g	0.33	0/828	0.68	0/1117
4	D	0.68	0/768	1.25	3/1032 (0.3%)
4	H	0.44	0/756	0.74	0/1015
4	d	0.44	0/768	0.78	0/1032
4	h	0.36	0/756	0.69	2/1015 (0.2%)
5	I	1.14	1/7773 (0.0%)	1.66	231/11987 (1.9%)
6	J	1.12	3/7865 (0.0%)	1.64	232/12145 (1.9%)
7	K	0.32	0/4699	0.54	1/6334 (0.0%)
8	L	0.34	0/3127	0.51	0/4231
9	M	0.32	0/2446	0.57	3/3292 (0.1%)
9	O	0.30	0/2235	0.58	2/3008 (0.1%)
10	N	0.32	0/3115	0.56	3/4195 (0.1%)
10	P	0.31	0/1278	0.60	2/1716 (0.1%)
All	All	0.74	4/45011 (0.0%)	1.17	508/63648 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
2	B	0	1
2	F	0	1
3	C	0	2
3	G	0	1
4	D	0	4
4	d	0	1
5	I	0	23
6	J	0	18
All	All	0	53

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	J	42	DG	C2-N2	-5.38	1.29	1.34
6	J	67	DC	C4-N4	-5.34	1.29	1.33
5	I	22	DG	C2-N2	-5.11	1.29	1.34
6	J	60	DG	C2-N2	-5.11	1.29	1.34

The worst 5 of 508 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	78	ARG	NE-CZ-NH2	10.19	125.39	120.30
1	A	72	ARG	NE-CZ-NH2	9.84	125.22	120.30
5	I	5	DA	N1-C6-N6	-9.59	112.85	118.60
6	J	31	DA	N1-C6-N6	-9.36	112.99	118.60
6	J	14	DA	N1-C6-N6	-9.28	113.03	118.60

There are no chirality outliers.

5 of 53 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	53	ARG	Sidechain
2	B	92	ARG	Sidechain
3	C	32	ARG	Sidechain
3	C	50	TYR	Sidechain
4	D	30	ARG	Sidechain

5.2 Too-close contacts

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/135 (73%)	92 (93%)	7 (7%)	0	100	100
1	E	113/135 (84%)	110 (97%)	3 (3%)	0	100	100
1	a	99/135 (73%)	96 (97%)	3 (3%)	0	100	100
1	e	96/135 (71%)	94 (98%)	2 (2%)	0	100	100
2	B	79/102 (78%)	77 (98%)	2 (2%)	0	100	100
2	F	78/102 (76%)	76 (97%)	2 (3%)	0	100	100
2	b	80/102 (78%)	78 (98%)	2 (2%)	0	100	100
2	f	78/102 (76%)	76 (97%)	2 (3%)	0	100	100
3	C	107/129 (83%)	104 (97%)	3 (3%)	0	100	100
3	G	104/129 (81%)	102 (98%)	2 (2%)	0	100	100
3	c	107/129 (83%)	105 (98%)	2 (2%)	0	100	100
3	g	104/129 (81%)	101 (97%)	3 (3%)	0	100	100
4	D	94/122 (77%)	90 (96%)	4 (4%)	0	100	100
4	H	93/122 (76%)	84 (90%)	9 (10%)	0	100	100
4	d	94/122 (77%)	92 (98%)	2 (2%)	0	100	100
4	h	93/122 (76%)	84 (90%)	9 (10%)	0	100	100
7	K	543/1536 (35%)	526 (97%)	17 (3%)	0	100	100
8	L	382/433 (88%)	369 (97%)	13 (3%)	0	100	100
9	M	288/401 (72%)	284 (99%)	4 (1%)	0	100	100
9	O	261/401 (65%)	255 (98%)	6 (2%)	0	100	100
10	N	362/684 (53%)	347 (96%)	15 (4%)	0	100	100
10	P	147/684 (22%)	140 (95%)	7 (5%)	0	100	100
All	All	3501/6091 (58%)	3382 (97%)	119 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/108 (80%)	86 (100%)	0	100	100
1	E	99/108 (92%)	98 (99%)	1 (1%)	73	82
1	a	86/108 (80%)	86 (100%)	0	100	100
1	e	85/108 (79%)	85 (100%)	0	100	100
2	B	66/78 (85%)	66 (100%)	0	100	100
2	F	65/78 (83%)	65 (100%)	0	100	100
2	b	67/78 (86%)	67 (100%)	0	100	100
2	f	65/78 (83%)	65 (100%)	0	100	100
3	C	86/101 (85%)	86 (100%)	0	100	100
3	G	84/101 (83%)	84 (100%)	0	100	100
3	c	86/101 (85%)	86 (100%)	0	100	100
3	g	84/101 (83%)	84 (100%)	0	100	100
4	D	82/102 (80%)	82 (100%)	0	100	100
4	H	81/102 (79%)	81 (100%)	0	100	100
4	d	82/102 (80%)	82 (100%)	0	100	100
4	h	81/102 (79%)	81 (100%)	0	100	100
7	K	510/1391 (37%)	510 (100%)	0	100	100
8	L	326/367 (89%)	324 (99%)	2 (1%)	84	88
9	M	268/359 (75%)	267 (100%)	1 (0%)	89	91
9	O	245/359 (68%)	245 (100%)	0	100	100
10	N	353/653 (54%)	353 (100%)	0	100	100
10	P	146/653 (22%)	146 (100%)	0	100	100
All	All	3133/5338 (59%)	3129 (100%)	4 (0%)	92	95

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	63	ARG
8	L	150	HIS
8	L	259	MET
9	M	292	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
9	M	389	ASN
3	c	38	ASN
10	N	304	ASN
1	e	108	ASN
9	O	275	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	ML3	A	36	1	10,11,12	0.77	0	10,14,16	0.83	0
1	ML3	a	36	1	10,11,12	0.76	0	10,14,16	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ML3	A	36	1	-	5/8/10/12	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ML3	a	36	1	-	5/8/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	36	ML3	SG-CD-CE-NZ
1	a	36	ML3	SG-CD-CE-NZ
1	A	36	ML3	CD-CE-NZ-CM1
1	A	36	ML3	CD-CE-NZ-CM2
1	a	36	ML3	CD-CE-NZ-CM1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.