



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 2, 2025 – 02:00 PM JST

PDB ID : 9KA2 / pdb_00009ka2
Title : Crystal structure of beta-ketoacyl-ACP synthase FabH from E. coli
Authors : Zhang, L.; Zhang, L.; Cai, C.
Deposited on : 2024-10-28
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3717 (1.50-1.50)
Clashscore	180529	4048 (1.50-1.50)
Ramachandran outliers	177936	3970 (1.50-1.50)
Sidechain outliers	177891	3967 (1.50-1.50)
RSRZ outliers	164620	3718 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>6% • 6%</div> </div> </div>
1	B	337	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>7% 6%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5316 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-ketoacyl-[acyl-carrier-protein] synthase III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	3	0
			2364	1485	407	458	14			
1	B	317	Total	C	N	O	S	0	0	0
			2350	1472	407	458	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP P0A6R0
A	-18	GLY	-	expression tag	UNP P0A6R0
A	-17	SER	-	expression tag	UNP P0A6R0
A	-16	SER	-	expression tag	UNP P0A6R0
A	-15	HIS	-	expression tag	UNP P0A6R0
A	-14	HIS	-	expression tag	UNP P0A6R0
A	-13	HIS	-	expression tag	UNP P0A6R0
A	-12	HIS	-	expression tag	UNP P0A6R0
A	-11	HIS	-	expression tag	UNP P0A6R0
A	-10	HIS	-	expression tag	UNP P0A6R0
A	-9	SER	-	expression tag	UNP P0A6R0
A	-8	SER	-	expression tag	UNP P0A6R0
A	-7	GLY	-	expression tag	UNP P0A6R0
A	-6	LEU	-	expression tag	UNP P0A6R0
A	-5	VAL	-	expression tag	UNP P0A6R0
A	-4	PRO	-	expression tag	UNP P0A6R0
A	-3	ARG	-	expression tag	UNP P0A6R0
A	-2	GLY	-	expression tag	UNP P0A6R0
A	-1	SER	-	expression tag	UNP P0A6R0
A	0	HIS	-	expression tag	UNP P0A6R0
B	-19	MET	-	initiating methionine	UNP P0A6R0
B	-18	GLY	-	expression tag	UNP P0A6R0
B	-17	SER	-	expression tag	UNP P0A6R0
B	-16	SER	-	expression tag	UNP P0A6R0
B	-15	HIS	-	expression tag	UNP P0A6R0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP P0A6R0
B	-13	HIS	-	expression tag	UNP P0A6R0
B	-12	HIS	-	expression tag	UNP P0A6R0
B	-11	HIS	-	expression tag	UNP P0A6R0
B	-10	HIS	-	expression tag	UNP P0A6R0
B	-9	SER	-	expression tag	UNP P0A6R0
B	-8	SER	-	expression tag	UNP P0A6R0
B	-7	GLY	-	expression tag	UNP P0A6R0
B	-6	LEU	-	expression tag	UNP P0A6R0
B	-5	VAL	-	expression tag	UNP P0A6R0
B	-4	PRO	-	expression tag	UNP P0A6R0
B	-3	ARG	-	expression tag	UNP P0A6R0
B	-2	GLY	-	expression tag	UNP P0A6R0
B	-1	SER	-	expression tag	UNP P0A6R0
B	0	HIS	-	expression tag	UNP P0A6R0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	312	Total O 312 312	0	0
2	B	290	Total O 290 290	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-ketoacyl-[acyl-carrier-protein] synthase III



- Molecule 1: Beta-ketoacyl-[acyl-carrier-protein] synthase III



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.27Å 64.54Å 164.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.57 – 1.50 30.57 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.57-1.50) 99.7 (30.57-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 1.50Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.185 , 0.198 0.186 , 0.200	Depositor DCC
R_{free} test set	5535 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.719	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5316	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2412	0.71	0/3282
1	B	0.40	0/2389	0.64	0/3251
All	All	0.44	0/4801	0.68	0/6533

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	151	ARG	Sidechain
1	A	249	ARG	Sidechain
1	B	235	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2364	0	2372	14	0
1	B	2350	0	2343	15	0
2	A	312	0	0	1	0
2	B	290	0	0	4	0
All	All	5316	0	4715	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:249:ARG:HG3	2:B:425:HOH:O	1.55	1.06
1:A:257:LYS:HE3	2:A:500:HOH:O	1.73	0.87
1:A:151:ARG:O	1:A:155:ILE:HG12	1.92	0.70
1:B:245:GLN:HB3	1:B:269:LEU:HB3	1.77	0.65
1:B:70:LYS:HD2	2:B:596:HOH:O	1.99	0.63
1:A:198:ASN:HB3	1:A:200:GLU:OE2	2.07	0.55
1:B:132:LYS:HD3	1:B:133:TYR:CE2	2.42	0.54
1:B:110:ALA:HB3	1:B:114:GLY:HA2	1.91	0.52
1:B:244:HIS:HD2	1:B:279:SER:HB3	1.76	0.50
1:A:260[B]:MET:CG	1:A:264:ASN:HB2	2.42	0.49
1:B:256:LYS:NZ	2:B:403:HOH:O	2.40	0.48
1:B:112:CYS:SG	2:B:593:HOH:O	2.59	0.48
1:B:87:PHE:HA	1:B:88:PRO:C	2.39	0.47
1:A:242:VAL:HB	1:A:301:LEU:HD23	1.97	0.47
1:B:223:ILE:HG12	1:B:312:SER:HB3	1.96	0.46
1:A:242:VAL:HB	1:A:301:LEU:CD2	2.47	0.45
1:A:87:PHE:HA	1:A:88:PRO:C	2.42	0.45
1:A:50:THR:O	1:A:54:MET:HG3	2.16	0.45
1:A:245:GLN:HB3	1:A:269[A]:LEU:HB3	2.00	0.44
1:B:19:ASN:O	1:B:23:GLU:HG3	2.18	0.43
1:A:22:LEU:HD13	1:A:155:ILE:HG22	2.01	0.43
1:A:260[B]:MET:HG2	1:A:264:ASN:HB2	2.01	0.43
1:A:247:ASN:OD1	1:A:249:ARG:HB2	2.19	0.42
1:A:172:PRO:HA	1:A:173:GLY:HA3	1.76	0.42
1:B:151:ARG:HA	1:B:154:ILE:HG22	2.02	0.42
1:B:214:LYS:N	1:B:214:LYS:HD2	2.34	0.42
1:A:76:ILE:HD11	1:A:100:ILE:HD11	2.02	0.41
1:B:247:ASN:HD21	1:B:250:ILE:HG13	1.85	0.41
1:B:225:ASP:OD1	1:B:235:ARG:NH1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/337 (94%)	311 (98%)	7 (2%)	0	100	100
1	B	315/337 (94%)	308 (98%)	7 (2%)	0	100	100
All	All	633/674 (94%)	619 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/264 (95%)	247 (99%)	3 (1%)	67	44
1	B	247/264 (94%)	245 (99%)	2 (1%)	79	62
All	All	497/528 (94%)	492 (99%)	5 (1%)	79	53

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	225	ASP
1	A	269[A]	LEU
1	A	269[B]	LEU
1	B	70	LYS
1	B	235	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/337 (94%)	0.52	10 (3%) 50 53	12, 21, 35, 48	3 (0%)
1	B	317/337 (94%)	0.54	9 (2%) 55 58	17, 25, 38, 47	0
All	All	634/674 (94%)	0.53	19 (2%) 52 56	12, 23, 36, 48	3 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	151	ARG	3.8
1	A	27	ASP	3.3
1	B	32	TRP	3.0
1	A	148	PRO	2.9
1	B	235	ARG	2.9
1	B	26	VAL	2.8
1	B	71	ASP	2.7
1	B	1	MET	2.6
1	A	26	VAL	2.5
1	A	31	GLU	2.5
1	A	1	MET	2.4
1	A	155	ILE	2.4
1	A	32	TRP	2.3
1	A	149	THR	2.3
1	B	317	PHE	2.3
1	B	234	ASP	2.2
1	B	100	ILE	2.1
1	A	269[A]	LEU	2.0
1	B	72	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.