



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2024 – 09:52 AM EST

PDB ID : 1KEL  
Title : CATALYTIC ANTIBODY 28B4 FAB FRAGMENT COMPLEXED WITH  
HAPTEN (1-[N-4'-NITROBENZYL-N-4'-CARBOXYBUTYLAMINO]  
METHYLPHOSPHONIC ACID)  
Authors : Hsieh-Wilson, L.C.; Schultz, P.G.; Stevens, R.C.  
Deposited on : 1996-04-16  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	L	217	<div> <div>5%</div> <div>83%</div> <div>16%</div> <div>.</div> </div>
2	H	218	<div> <div>11%</div> <div>74%</div> <div>23%</div> <div>.</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4154 atoms, of which 796 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 28B4 FAB.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	217	Total	C	H	N	O	S	0	0	0
			2091	1055	403	288	339	6			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	25	PHE	SER	conflict	PIR PC4203
L	32	SER	THR	conflict	PIR PC4203
L	45	SER	PRO	conflict	PIR PC4203

- Molecule 2 is a protein called 28B4 FAB.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	218	Total	C	H	N	O	S	0	0	0
			2022	1043	375	275	322	7			

- Molecule 3 is 1-[N-4'-NITROBENZYL-N-4'-CARBOXYBUTYLAMINO]METHYLPHOSPHONIC ACID (three-letter code: AAH) (formula: C<sub>13</sub>H<sub>19</sub>N<sub>2</sub>O<sub>7</sub>P).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
3	H	1	41	13	18	2	7	1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 28B4 FAB



#### • Molecule 2: 28B4 FAB



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.70Å 58.20Å 43.20Å 94.30° 113.80° 78.90°	Depositor
Resolution (Å)	20.00 – 1.90 20.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.0 (20.00-1.90) 90.4 (20.00-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.25 (at 1.90Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.199 , 0.258 0.262 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.6	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 39.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	4154	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.59	0/1726	0.79	0/2338
2	H	0.61	0/1692	0.81	0/2309
All	All	0.60	0/3418	0.80	0/4647

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1688	403	1631	24	1
2	H	1647	375	1604	38	0
3	H	23	18	17	0	0
All	All	3358	796	3252	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:66:ARG:HG2	1:L:82:ARG:NH1	1.79	0.98
2:H:162:SER:H	2:H:202:ASN:HD21	1.28	0.81
2:H:190:PRO:O	2:H:193:PRO:HD2	1.85	0.76
2:H:125:PRO:HB3	2:H:151:TYR:HB3	1.68	0.74
2:H:192:SER:OG	2:H:193:PRO:HD3	1.89	0.72
2:H:162:SER:H	2:H:202:ASN:ND2	1.90	0.68
1:L:66:ARG:HG2	1:L:82:ARG:HH11	1.56	0.66
1:L:146:PRO:O	1:L:203:HIS:HE1	1.80	0.65
2:H:153:PRO:O	2:H:205:HIS:HE1	1.85	0.60
2:H:161:ASN:O	2:H:164:SER:HB2	2.02	0.60
1:L:66:ARG:HG2	1:L:82:ARG:HH12	1.62	0.59
1:L:2:VAL:HG13	1:L:27:GLN:HG2	1.85	0.57
2:H:198:THR:HG23	2:H:215:LYS:HD2	1.86	0.57
1:L:33:ASN:ND2	1:L:35:ASN:H	2.03	0.56
2:H:40:PRO:HB2	2:H:43:LYS:HB2	1.87	0.56
2:H:190:PRO:HB2	2:H:193:PRO:HD2	1.87	0.55
1:L:6:GLN:HE21	1:L:104:GLY:HA3	1.71	0.55
2:H:176:LEU:HD11	2:H:179:ASP:HA	1.89	0.54
1:L:33:ASN:HD22	1:L:35:ASN:H	1.54	0.54
2:H:2:VAL:HG13	2:H:27:PHE:CD1	2.43	0.54
1:L:53:ILE:HD13	1:L:59:ARG:HA	1.89	0.53
2:H:190:PRO:HB2	2:H:193:PRO:CD	2.37	0.53
2:H:161:ASN:ND2	2:H:200:THR:H	2.05	0.53
1:L:33:ASN:HD22	1:L:33:ASN:C	2.12	0.53
1:L:126:SER:OG	1:L:128:GLU:HG2	2.09	0.52
2:H:58:TYR:CD1	2:H:74:ARG:HD3	2.44	0.52
2:H:124:PRO:HG3	2:H:208:SER:HB2	1.92	0.52
2:H:34:PHE:HB3	2:H:81:LEU:HD22	1.93	0.51
2:H:96:TYR:O	2:H:112:GLY:HA2	2.11	0.50
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.48	0.49
1:L:143:ASN:OD1	2:H:170:HIS:HE1	1.95	0.49
2:H:194:ARG:HD2	2:H:195:PRO:HA	1.93	0.49
1:L:24:ARG:HG3	1:L:24:ARG:HH11	1.79	0.48
2:H:141:MET:SD	2:H:190:PRO:HA	2.54	0.47
2:H:198:THR:HG23	2:H:215:LYS:CD	2.44	0.47
2:H:53:ASN:HD21	2:H:56:LYS:HG2	1.79	0.47
1:L:203:HIS:HD2	1:L:205:THR:OG1	1.98	0.47
1:L:40:TRP:CZ3	1:L:93:CYS:HB3	2.50	0.47
1:L:66:ARG:HD2	1:L:81:SER:O	2.15	0.47
1:L:165:LEU:CD1	2:H:177:GLN:HG3	2.45	0.46
2:H:100:ARG:O	2:H:106:MET:HA	2.16	0.46
1:L:120:VAL:HA	1:L:140:PHE:O	2.15	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:190:PRO:C	2:H:193:PRO:HD2	2.36	0.46
2:H:11:LEU:HD12	2:H:116:THR:O	2.14	0.46
1:L:141:LEU:HD23	1:L:201:ALA:HB2	1.99	0.45
1:L:42:LEU:HB2	1:L:52:LEU:HD11	1.98	0.45
2:H:29:PHE:HZ	2:H:74:ARG:HB2	1.81	0.45
1:L:71:GLY:HA3	1:L:76:PHE:HA	2.00	0.44
2:H:205:HIS:HD2	2:H:208:SER:OG	2.01	0.44
1:L:24:ARG:HH11	1:L:24:ARG:CG	2.30	0.44
2:H:160:TRP:CZ3	2:H:201:CYS:HB3	2.53	0.44
2:H:35:ASN:OD1	2:H:50:PHE:HB3	2.16	0.44
2:H:71:THR:HB	2:H:84:GLN:HB3	2.00	0.43
1:L:165:LEU:HD11	2:H:177:GLN:HG3	2.00	0.43
2:H:161:ASN:HD21	2:H:200:THR:H	1.66	0.43
2:H:40:PRO:HB2	2:H:43:LYS:HD3	2.00	0.42
1:L:165:LEU:HD23	1:L:165:LEU:HA	1.85	0.42
2:H:62:TYR:HE1	2:H:72:ILE:HG13	1.84	0.41
2:H:33:TYR:CE2	2:H:52:ARG:HG2	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:57:SER:HG	1:L:216:ARG:HH22[1_566]	1.29	0.31

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	215/217 (99%)	212 (99%)	3 (1%)	0	100	100
2	H	216/218 (99%)	210 (97%)	5 (2%)	1 (0%)	25	17
All	All	431/435 (99%)	422 (98%)	8 (2%)	1 (0%)	44	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	155	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	194/194 (100%)	186 (96%)	8 (4%)	26	19
2	H	182/182 (100%)	172 (94%)	10 (6%)	18	10
All	All	376/376 (100%)	358 (95%)	18 (5%)	21	14

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	9	LEU
1	L	27	GLN
1	L	33	ASN
1	L	45	SER
1	L	50	LYS
1	L	84	GLU
1	L	204	LYS
1	L	208	SER
2	H	13	GLN
2	H	50	PHE
2	H	53	ASN
2	H	137	GLN
2	H	138	THR
2	H	164	SER
2	H	176	LEU
2	H	183	LEU
2	H	193	PRO
2	H	194	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	6	GLN
1	L	18	GLN
1	L	27	GLN
1	L	33	ASN
1	L	142	ASN
1	L	203	HIS
2	H	53	ASN
2	H	161	ASN
2	H	170	HIS
2	H	202	ASN
2	H	205	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	AAH	H	219	-	23,23,23	1.98	8 (34%)	29,31,31	1.29	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AAH	H	219	-	-	3/18/20/20	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	219	AAH	P13-O16	-4.31	1.45	1.55
3	H	219	AAH	C17-N11	3.38	1.55	1.47
3	H	219	AAH	P13-O14	3.28	1.56	1.50
3	H	219	AAH	P13-O15	3.17	1.62	1.55
3	H	219	AAH	O8-N7	-2.69	1.18	1.22
3	H	219	AAH	O9-N7	-2.37	1.19	1.35
3	H	219	AAH	O22-C21	2.33	1.38	1.30
3	H	219	AAH	C6-N7	2.13	1.50	1.45

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	219	AAH	O15-P13-C12	3.12	113.66	106.78
3	H	219	AAH	C10-N11-C17	2.66	116.81	111.28
3	H	219	AAH	O15-P13-O14	-2.21	106.68	112.39
3	H	219	AAH	O22-C21-C20	2.17	120.87	114.00

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	H	219	AAH	N11-C17-C18-C19
3	H	219	AAH	C19-C20-C21-O23
3	H	219	AAH	C19-C20-C21-O22

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

**Warning:** The R factor obtained from EDS is 0.2697, which does not match the depositor's R factor of 0.199. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	217/217 (100%)	0.50	11 (5%) 34 36	5, 11, 21, 29	0
2	H	218/218 (100%)	0.79	23 (10%) 13 13	6, 13, 25, 49	0
All	All	435/435 (100%)	0.64	34 (7%) 20 21	5, 12, 22, 49	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	135	ALA	7.3
2	H	138	THR	6.8
2	H	134	SER	5.4
2	H	137	GLN	5.2
1	L	72	SER	5.1
2	H	136	ALA	5.0
2	H	42	GLY	3.6
1	L	217	ASN	3.6
1	L	71	GLY	3.5
2	H	139	ASN	3.2
2	H	13	GLN	3.1
2	H	140	SER	3.1
2	H	120	ALA	3.1
1	L	214	PHE	3.0
2	H	111	GLN	2.7
2	H	123	THR	2.7
2	H	178	SER	2.6
2	H	121	LYS	2.6
1	L	46	GLY	2.6
2	H	193	PRO	2.6
1	L	82	ARG	2.5
1	L	150	ASN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	H	122	THR	2.3
2	H	176	LEU	2.3
2	H	167	SER	2.2
2	H	211	LYS	2.2
1	L	18	GLN	2.1
2	H	11	LEU	2.1
2	H	165	LEU	2.1
2	H	43	LYS	2.1
1	L	2	VAL	2.1
1	L	9	LEU	2.1
1	L	174	LYS	2.0
2	H	163	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	AAH	H	219	23/23	0.93	0.10	0,11,34,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers ⓘ

There are no such residues in this entry.