



wwPDB EM Validation Summary Report ⓘ

May 18, 2024 – 09:22 AM EDT

PDB ID : 7KGB
EMDB ID : EMD-22865
Title : CryoEM structure of A2296-methylated Mycobacterium tuberculosis ribosome bound with SEQ-9
Authors : Cui, Z.; Zhang, J.
Deposited on : 2020-10-16
Resolution : 2.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev92
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	0	57	95% 5%
2	1	55	87% 13%
3	2	47	89% 11%
4	3	64	97% .
5	4	37	100%
6	6	80	9% 71% 29%
7	A	3138	8% 85% 15% .
8	B	115	89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	C	280	
10	D	217	
11	E	223	
12	F	187	
13	G	179	
14	H	152	
15	J	195	
16	K	122	
17	L	146	
18	M	138	
19	N	180	
20	O	122	
21	P	113	
22	Q	129	
23	R	104	
24	S	197	
25	T	100	
26	U	105	
27	V	215	
28	W	86	
29	X	64	
30	Y	77	
31	Z	65	
32	a	1537	
33	c	274	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
34	d	201	
35	e	220	
36	f	96	
37	g	156	
38	h	132	
39	i	151	
40	j	101	
41	k	139	
42	l	124	
43	m	124	
44	n	61	
45	o	89	
46	p	162	
47	q	135	
48	r	84	
49	s	93	
50	t	86	
51	v	24	
52	y	77	

2 Entry composition

There are 55 unique types of molecules in this entry. The entry contains 145319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	0	54	Total	C	N	O	0	0
			429	266	94	69		

- Molecule 2 is a protein called 50S ribosomal protein L33 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	1	48	Total	C	N	O	S	0	0
			400	245	84	67	4		

- Molecule 3 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	2	42	Total	C	N	O	S	0	0
			358	212	94	51	1		

- Molecule 4 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues	Atoms				AltConf	Trace
4	3	62	Total	C	N	O	0	0
			494	298	112	84		

- Molecule 5 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	4	37	Total	C	N	O	S	0	0
			299	182	66	47	4		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
4	1	VAL	MET	conflict	UNP P9WH89

- Molecule 6 is a protein called 50S ribosomal protein L31.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	6	57	Total	C	N	O	S	0	0
			446	277	82	82	5		

- Molecule 7 is a RNA chain called 23S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	A	3118	Total	C	N	O	P	0	0
			66962	29851	12340	21653	3118		

- Molecule 8 is a RNA chain called 5S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	B	115	Total	C	N	O	P	0	0
			2458	1097	456	790	115		

- Molecule 9 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	C	272	Total	C	N	O	S	0	0
			2088	1277	437	369	5		

- Molecule 10 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	D	213	Total	C	N	O	S	0	0
			1590	985	307	292	6		

- Molecule 11 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	E	207	Total	C	N	O	S	0	0
			1552	958	303	289	2		

- Molecule 12 is a protein called 50S ribosomal protein L5.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	F	178	Total	C	N	O	S	0	0
			1408	885	267	251	5		

- Molecule 13 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	G	174	Total	C	N	O	S	0	0
			1330	836	249	244	1		

- Molecule 14 is a protein called 50S ribosomal protein L9.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	H	47	Total	C	N	O	S	0	0
			350	220	64	65	1		

- Molecule 15 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	J	146	Total	C	N	O	S	0	0
			1143	724	217	199	3		

- Molecule 16 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	K	121	Total	C	N	O	S	0	0
			934	585	179	168	2		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	1	VAL	MET	conflict	UNP A0A045HTP7

- Molecule 17 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	L	143	Total	C	N	O	S	0	0
			1068	662	216	188	2		

- Molecule 18 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	M	134	Total	C	N	O	S	0	0
			1072	679	215	177	1		

- Molecule 19 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	N	116	Total	C	N	O	S	0	0
			908	574	175	158	1		

- Molecule 20 is a protein called 50S ribosomal protein L18.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	O	116	Total	C	N	O		0	0
			886	541	188	157			

- Molecule 21 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	P	112	Total	C	N	O	S	0	0
			907	573	174	159	1		

- Molecule 22 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	Q	122	Total	C	N	O		0	0
			980	608	205	167			

- Molecule 23 is a protein called 50S ribosomal protein L21.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	R	100	Total	C	N	O		0	0
			757	482	138	137			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	1	MET	-	insertion	UNP P9WHC3

- Molecule 24 is a protein called 50S ribosomal protein L22.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	S	113	Total	C	N	O		0	0
			860	533	178	149			

- Molecule 25 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues	Atoms				AltConf	Trace
25	T	98	Total	C	N	O	0	0
			759	480	141	138		

- Molecule 26 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues	Atoms				AltConf	Trace
26	U	90	Total	C	N	O	S	0
			699	430	138	129	2	0

- Molecule 27 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms				AltConf	Trace
27	V	95	Total	C	N	O		0
			735	456	152	127		0

- Molecule 28 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues	Atoms				AltConf	Trace
28	W	71	Total	C	N	O	0	0
			526	325	108	93		

- Molecule 29 is a protein called 50S ribosomal protein L28.

Mol	Chain	Residues	Atoms				AltConf	Trace
29	X	63	Total	C	N	O	S	0
			476	289	101	81	5	0

- Molecule 30 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms				AltConf	Trace
30	Y	65	Total	C	N	O	S	0
			541	331	106	103	1	0

- Molecule 31 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues	Atoms				AltConf	Trace
31	Z	59	Total	C	N	O	0	0
			476	293	101	82		

- Molecule 32 is a RNA chain called 16S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	a	1519	Total	C	N	O	P	0	0
			32621	14536	5961	10605	1519		

- Molecule 33 is a protein called 30S ribosomal protein S3.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	c	207	Total	C	N	O	S	0	0
			1654	1030	322	298	4		

- Molecule 34 is a protein called 30S ribosomal protein S4.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	d	200	Total	C	N	O	S	0	0
			1650	1036	316	296	2		

- Molecule 35 is a protein called 30S ribosomal protein S5.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	e	160	Total	C	N	O	S	0	0
			1149	726	214	206	3		

- Molecule 36 is a protein called 30S ribosomal protein S6.

Mol	Chain	Residues	Atoms					AltConf	Trace
36	f	95	Total	C	N	O	S	0	0
			757	480	133	141	3		

- Molecule 37 is a protein called 30S ribosomal protein S7.

Mol	Chain	Residues	Atoms					AltConf	Trace
37	g	152	Total	C	N	O	S	0	0
			1193	742	234	215	2		

- Molecule 38 is a protein called 30S ribosomal protein S8.

Mol	Chain	Residues	Atoms					AltConf	Trace
38	h	130	Total	C	N	O	S	0	0
			999	627	187	184	1		

- Molecule 39 is a protein called 30S ribosomal protein S9.

Mol	Chain	Residues	Atoms				AltConf	Trace
39	i	127	Total	C	N	O	0	0
			993	628	195	170		

- Molecule 40 is a protein called 30S ribosomal protein S10.

Mol	Chain	Residues	Atoms					AltConf	Trace
40	j	99	Total	C	N	O	S	0	0
			789	496	146	144	3		

- Molecule 41 is a protein called 30S ribosomal protein S11.

Mol	Chain	Residues	Atoms				AltConf	Trace
41	k	117	Total	C	N	O	0	0
			873	540	175	158		

- Molecule 42 is a protein called 30S ribosomal protein S12.

Mol	Chain	Residues	Atoms					AltConf	Trace
42	l	122	Total	C	N	O	S	0	0
			959	594	197	166	2		

- Molecule 43 is a protein called 30S ribosomal protein S13.

Mol	Chain	Residues	Atoms					AltConf	Trace
43	m	116	Total	C	N	O	S	0	0
			945	578	196	168	3		

- Molecule 44 is a protein called 30S ribosomal protein S14 type Z.

Mol	Chain	Residues	Atoms					AltConf	Trace
44	n	60	Total	C	N	O	S	0	0
			468	294	96	73	5		

- Molecule 45 is a protein called 30S ribosomal protein S15.

Mol	Chain	Residues	Atoms				AltConf	Trace
45	o	87	Total	C	N	O	0	0
			718	449	144	125		

- Molecule 46 is a protein called 30S ribosomal protein S16.

Mol	Chain	Residues	Atoms				AltConf	Trace
46	p	91	Total	C	N	O	0	0
			728	462	140	126		

- Molecule 47 is a protein called 30S ribosomal protein S17.

Mol	Chain	Residues	Atoms					AltConf	Trace
47	q	94	Total	C	N	O	S	0	0
			763	477	151	132	3		

- Molecule 48 is a protein called 30S ribosomal protein S18 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
48	r	63	Total	C	N	O	S	0	0
			497	309	96	89	3		

- Molecule 49 is a protein called 30S ribosomal protein S19.

Mol	Chain	Residues	Atoms					AltConf	Trace
49	s	83	Total	C	N	O	S	0	0
			672	432	125	114	1		

- Molecule 50 is a protein called 30S ribosomal protein S20.

Mol	Chain	Residues	Atoms				AltConf	Trace
50	t	82	Total	C	N	O	0	0
			631	381	137	113		

- Molecule 51 is a protein called 50S ribosomal protein L37.

Mol	Chain	Residues	Atoms				AltConf	Trace
51	v	22	Total	C	N	O	0	0
			186	111	47	28		


- Molecule 52 is a RNA chain called tRNA-initiator Met(CAU).

Mol	Chain	Residues	Atoms					AltConf	Trace
52	y	77	Total	C	N	O	P	0	0
			1644	732	297	538	77		

- Molecule 53 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
53	1	1	Total	Zn	0
			1	1	
53	4	1	Total	Zn	0
			1	1	
53	6	1	Total	Zn	0
			1	1	
53	X	1	Total	Zn	0
			1	1	
53	n	1	Total	Zn	0
			1	1	
53	r	1	Total	Zn	0
			1	1	

- Molecule 54 is Sequanamycin 9 (three-letter code: WDP) (formula: $C_{61}H_{102}N_4O_{20}S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
54	A	1	Total	C	N	O	S	0
			86	61	4	20	1	

- Molecule 55 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
55	A	306	Total	Mg	0
			306	306	
55	B	7	Total	Mg	0
			7	7	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		AltConf
55	C	3	Total 3	Mg 3	0
55	D	1	Total 1	Mg 1	0
55	L	1	Total 1	Mg 1	0
55	M	1	Total 1	Mg 1	0
55	a	126	Total 126	Mg 126	0
55	t	1	Total 1	Mg 1	0
55	y	1	Total 1	Mg 1	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 50S ribosomal protein L32

Chain 0:  95% 5%





- Molecule 2: 50S ribosomal protein L33 2

Chain 1:  87% 13%



- Molecule 3: 50S ribosomal protein L34

Chain 2:  89% 11%



- Molecule 4: 50S ribosomal protein L35

Chain 3:  97% .




- Molecule 5: 50S ribosomal protein L36



Chain 4:  100%




- Molecule 6: 50S ribosomal protein L31





• Molecule 7: 23S rRNA






- Molecule 8: 5S rRNA




- Molecule 9: 50S ribosomal protein L2



- Molecule 10: 50S ribosomal protein L3




- Molecule 11: 50S ribosomal protein L4




- Molecule 12: 50S ribosomal protein L5

Chain F:  94% .. 5%




- Molecule 13: 50S ribosomal protein L6

Chain G:  7% 97% ..




- Molecule 14: 50S ribosomal protein L9

Chain H:  27% 31% 69%




- Molecule 15: 50S ribosomal protein L13

Chain J:  75% 25%




- Molecule 16: 50S ribosomal protein L14

Chain K:  99%




- Molecule 17: 50S ribosomal protein L15

Chain L:  97% ..




- Molecule 18: 50S ribosomal protein L16

Chain M:  97%




- Molecule 19: 50S ribosomal protein L17

Chain N:  64% 36%




- Molecule 20: 50S ribosomal protein L18

Chain O:  95% 5%




- Molecule 21: 50S ribosomal protein L19

Chain P:  99%




- Molecule 22: 50S ribosomal protein L20

Chain Q:  95% 5%




- Molecule 23: 50S ribosomal protein L21

Chain R:  94%




- Molecule 24: 50S ribosomal protein L22



Chain S:  57% 43%






- Molecule 25: 50S ribosomal protein L23





- Molecule 26: 50S ribosomal protein L24



- Molecule 27: 50S ribosomal protein L25



- Molecule 28: 50S ribosomal protein L27



- Molecule 29: 50S ribosomal protein L28





- Molecule 30: 50S ribosomal protein L29


MET
ALA
VAL
G4
L68
ALA
THR
GLY
PRO
ASP
GLY
LYS
GLU
SER

- Chain Z:  91% 9%


- Chain a: 84% 14%




- Chain c:  9% 75% 24%




- Molecule 40: 30S ribosomal protein S10




- Molecule 41: 30S ribosomal protein S11




- Molecule 42: 30S ribosomal protein S12




- Molecule 43: 30S ribosomal protein S13



- Molecule 44: 30S ribosomal protein S14 type Z




- Molecule 45: 30S ribosomal protein S15




- Molecule 46: 30S ribosomal protein S16








- Molecule 47: 30S ribosomal protein S17




- Molecule 48: 30S ribosomal protein S18 1




- Molecule 49: 30S ribosomal protein S19




- Molecule 50: 30S ribosomal protein S20




- Molecule 51: 50S ribosomal protein L37



- Molecule 52: tRNA-initiator Met(CAU)





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	1065544	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.173	Depositor
Minimum map value	-0.059	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.02	Depositor
Map size (\AA)	339.19998, 339.19998, 339.19998	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4OC, G7M, 2MG, 5MU, 5MC, MG, OMG, ZN, WDP, OMC, UR3, MA6, 6MZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	0	0.39	0/435	0.59	0/581
2	1	0.36	0/407	0.58	0/543
3	2	0.39	0/361	0.74	0/473
4	3	0.34	0/499	0.63	0/664
5	4	0.36	0/303	0.62	0/402
6	6	0.32	0/455	0.57	0/611
7	A	0.79	1/74825 (0.0%)	0.82	31/116743 (0.0%)
8	B	0.62	0/2749	0.79	0/4284
9	C	0.41	0/2129	0.60	0/2861
10	D	0.40	0/1613	0.59	0/2174
11	E	0.37	0/1575	0.57	0/2129
12	F	0.35	0/1429	0.60	0/1921
13	G	0.31	0/1351	0.56	0/1824
14	H	0.26	0/353	0.53	0/474
15	J	0.40	0/1170	0.55	0/1584
16	K	0.40	0/944	0.60	0/1268
17	L	0.37	0/1081	0.60	0/1443
18	M	0.40	0/1098	0.61	0/1481
19	N	0.39	0/925	0.55	0/1242
20	O	0.37	0/895	0.64	0/1202
21	P	0.40	0/922	0.60	0/1236
22	Q	0.42	0/992	0.60	0/1329
23	R	0.39	0/766	0.53	0/1030
24	S	0.38	0/874	0.59	0/1186
25	T	0.38	0/770	0.58	0/1038
26	U	0.35	0/705	0.68	1/941 (0.1%)
27	V	0.31	0/747	0.59	0/1010
28	W	0.42	0/531	0.62	0/707
29	X	0.38	0/484	0.58	0/648
30	Y	0.36	0/544	0.65	0/727
31	Z	0.35	0/480	0.63	0/645
32	a	0.66	0/36305	0.80	9/56645 (0.0%)

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
33	c	0.32	0/1678	0.58	0/2254
34	d	0.33	0/1683	0.60	0/2269
35	e	0.35	0/1165	0.60	1/1578 (0.1%)
36	f	0.36	0/767	0.58	0/1036
37	g	0.31	0/1210	0.56	0/1631
38	h	0.37	0/1014	0.56	0/1369
39	i	0.31	0/1011	0.61	0/1356
40	j	0.32	0/803	0.60	0/1086
41	k	0.33	0/891	0.54	0/1204
42	l	0.36	0/970	0.65	0/1295
43	m	0.31	0/953	0.65	0/1274
44	n	0.38	0/477	0.57	0/634
45	o	0.34	0/727	0.62	0/973
46	p	0.34	0/742	0.66	0/1000
47	q	0.33	0/774	0.69	1/1032 (0.1%)
48	r	0.37	0/502	0.63	0/674
49	s	0.30	0/690	0.53	0/928
50	t	0.28	0/633	0.52	0/838
51	v	0.35	0/188	0.64	0/243
52	y	0.60	1/1836 (0.1%)	0.79	0/2859
All	All	0.67	2/157431 (0.0%)	0.77	43/236579 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
12	F	0	1
23	R	0	1
26	U	0	1
36	f	0	1
39	i	0	1
42	l	0	2
46	p	0	1
All	All	0	8

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
52	y	1	C	OP3-P	-10.83	1.48	1.61
7	A	616	A	N9-C4	-5.14	1.34	1.37

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	600	G	O4'-C1'-N9	8.21	114.76	108.20
7	A	1955	G	N3-C2-N2	-6.79	115.15	119.90
7	A	2309	C	N3-C2-O2	-6.54	117.33	121.90
7	A	946	C	N3-C2-O2	-6.28	117.50	121.90
7	A	3060	C	C2-N1-C1'	6.27	125.70	118.80

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
12	F	143	ALA	Peptide
23	R	50	THR	Peptide
26	U	90	GLU	Peptide
36	f	17	ARG	Sidechain
39	i	78	ARG	Peptide

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	0	52/57 (91%)	52 (100%)	0	0	100	100
2	1	46/55 (84%)	46 (100%)	0	0	100	100
3	2	40/47 (85%)	39 (98%)	1 (2%)	0	100	100
4	3	60/64 (94%)	60 (100%)	0	0	100	100
5	4	35/37 (95%)	34 (97%)	1 (3%)	0	100	100
6	6	53/80 (66%)	46 (87%)	7 (13%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
9	C	270/280 (96%)	259 (96%)	11 (4%)	0	100	100
10	D	211/217 (97%)	198 (94%)	13 (6%)	0	100	100
11	E	205/223 (92%)	201 (98%)	4 (2%)	0	100	100
12	F	176/187 (94%)	162 (92%)	12 (7%)	2 (1%)	14	34
13	G	172/179 (96%)	162 (94%)	10 (6%)	0	100	100
14	H	45/152 (30%)	41 (91%)	4 (9%)	0	100	100
15	J	144/195 (74%)	141 (98%)	3 (2%)	0	100	100
16	K	119/122 (98%)	113 (95%)	6 (5%)	0	100	100
17	L	141/146 (97%)	127 (90%)	13 (9%)	1 (1%)	22	46
18	M	132/138 (96%)	122 (92%)	10 (8%)	0	100	100
19	N	114/180 (63%)	113 (99%)	1 (1%)	0	100	100
20	O	114/122 (93%)	107 (94%)	7 (6%)	0	100	100
21	P	110/113 (97%)	103 (94%)	7 (6%)	0	100	100
22	Q	120/129 (93%)	117 (98%)	3 (2%)	0	100	100
23	R	98/104 (94%)	87 (89%)	10 (10%)	1 (1%)	15	37
24	S	111/197 (56%)	108 (97%)	3 (3%)	0	100	100
25	T	96/100 (96%)	89 (93%)	7 (7%)	0	100	100
26	U	86/105 (82%)	74 (86%)	11 (13%)	1 (1%)	13	32
27	V	93/215 (43%)	85 (91%)	8 (9%)	0	100	100
28	W	69/86 (80%)	65 (94%)	4 (6%)	0	100	100
29	X	61/64 (95%)	56 (92%)	5 (8%)	0	100	100
30	Y	63/77 (82%)	60 (95%)	3 (5%)	0	100	100
31	Z	57/65 (88%)	55 (96%)	2 (4%)	0	100	100
33	c	205/274 (75%)	187 (91%)	18 (9%)	0	100	100
34	d	198/201 (98%)	174 (88%)	24 (12%)	0	100	100
35	e	158/220 (72%)	140 (89%)	18 (11%)	0	100	100
36	f	93/96 (97%)	86 (92%)	7 (8%)	0	100	100
37	g	150/156 (96%)	144 (96%)	6 (4%)	0	100	100
38	h	128/132 (97%)	124 (97%)	4 (3%)	0	100	100
39	i	125/151 (83%)	107 (86%)	18 (14%)	0	100	100
40	j	97/101 (96%)	85 (88%)	12 (12%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
41	k	115/139 (83%)	110 (96%)	5 (4%)	0	100	100
42	l	120/124 (97%)	90 (75%)	29 (24%)	1 (1%)	19	43
43	m	114/124 (92%)	102 (90%)	12 (10%)	0	100	100
44	n	58/61 (95%)	54 (93%)	4 (7%)	0	100	100
45	o	85/89 (96%)	77 (91%)	7 (8%)	1 (1%)	13	32
46	p	89/162 (55%)	78 (88%)	10 (11%)	1 (1%)	14	34
47	q	91/135 (67%)	75 (82%)	16 (18%)	0	100	100
48	r	61/84 (73%)	56 (92%)	5 (8%)	0	100	100
49	s	81/93 (87%)	72 (89%)	9 (11%)	0	100	100
50	t	80/86 (93%)	77 (96%)	3 (4%)	0	100	100
51	v	20/24 (83%)	20 (100%)	0	0	100	100
All	All	5161/6188 (83%)	4780 (93%)	373 (7%)	8 (0%)	50	73

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
12	F	144	VAL
23	R	51	THR
42	l	114	ARG
12	F	143	ALA
17	L	5	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	0	44/47 (94%)	44 (100%)	0	100	100
2	1	45/51 (88%)	45 (100%)	0	100	100
3	2	36/40 (90%)	36 (100%)	0	100	100
4	3	53/54 (98%)	53 (100%)	0	100	100
5	4	35/35 (100%)	35 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	6	49/66 (74%)	49 (100%)	0	100	100
9	C	212/219 (97%)	212 (100%)	0	100	100
10	D	163/166 (98%)	163 (100%)	0	100	100
11	E	159/172 (92%)	158 (99%)	1 (1%)	86	95
12	F	147/155 (95%)	147 (100%)	0	100	100
13	G	143/147 (97%)	142 (99%)	1 (1%)	84	94
14	H	36/121 (30%)	36 (100%)	0	100	100
15	J	120/161 (74%)	120 (100%)	0	100	100
16	K	100/101 (99%)	100 (100%)	0	100	100
17	L	107/110 (97%)	107 (100%)	0	100	100
18	M	110/114 (96%)	110 (100%)	0	100	100
19	N	94/139 (68%)	93 (99%)	1 (1%)	73	90
20	O	88/93 (95%)	88 (100%)	0	100	100
21	P	98/99 (99%)	98 (100%)	0	100	100
22	Q	95/99 (96%)	95 (100%)	0	100	100
23	R	81/83 (98%)	81 (100%)	0	100	100
24	S	87/140 (62%)	87 (100%)	0	100	100
25	T	82/83 (99%)	81 (99%)	1 (1%)	71	88
26	U	77/88 (88%)	77 (100%)	0	100	100
27	V	75/164 (46%)	75 (100%)	0	100	100
28	W	51/62 (82%)	51 (100%)	0	100	100
29	X	52/52 (100%)	52 (100%)	0	100	100
30	Y	58/66 (88%)	58 (100%)	0	100	100
31	Z	51/55 (93%)	51 (100%)	0	100	100
33	c	170/210 (81%)	169 (99%)	1 (1%)	86	95
34	d	176/177 (99%)	175 (99%)	1 (1%)	86	95
35	e	114/159 (72%)	114 (100%)	0	100	100
36	f	84/85 (99%)	84 (100%)	0	100	100
37	g	127/131 (97%)	127 (100%)	0	100	100
38	h	106/108 (98%)	106 (100%)	0	100	100
39	i	102/120 (85%)	100 (98%)	2 (2%)	55	81

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
40	j	89/90 (99%)	87 (98%)	2 (2%)	52	79
41	k	90/107 (84%)	90 (100%)	0	100	100
42	l	104/105 (99%)	104 (100%)	0	100	100
43	m	99/104 (95%)	99 (100%)	0	100	100
44	n	46/47 (98%)	46 (100%)	0	100	100
45	o	77/79 (98%)	77 (100%)	0	100	100
46	p	75/125 (60%)	74 (99%)	1 (1%)	69	87
47	q	84/105 (80%)	83 (99%)	1 (1%)	71	88
48	r	53/72 (74%)	53 (100%)	0	100	100
49	s	75/85 (88%)	75 (100%)	0	100	100
50	t	62/65 (95%)	62 (100%)	0	100	100
51	v	18/20 (90%)	18 (100%)	0	100	100
All	All	4299/4976 (86%)	4287 (100%)	12 (0%)	92	98

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
39	i	127	ARG
40	j	23	ARG
47	q	53	LYS
40	j	82	LYS
25	T	70	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
34	d	126	ASN
46	p	41	HIS
46	p	65	GLN
20	O	118	ASN
10	D	34	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
32	a	1515/1537 (98%)	210 (13%)	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
52	y	76/77 (98%)	15 (19%)	0
7	A	3114/3138 (99%)	440 (14%)	2 (0%)
8	B	114/115 (99%)	13 (11%)	0
All	All	4819/4867 (99%)	678 (14%)	2 (0%)

5 of 678 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
7	A	13	A
7	A	35	G
7	A	44	A
7	A	51	G
7	A	63	A

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
7	A	913	G
7	A	3127	A

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

14 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	6MZ	A	2296	7	18,25,26	2.31	3 (16%)	16,36,39	2.11	4 (25%)
32	5MC	a	960	32	18,22,23	1.76	5 (27%)	26,32,35	1.19	3 (11%)
32	4OC	a	1395	32	20,23,24	2.75	6 (30%)	26,32,35	1.13	3 (11%)
32	MA6	a	1512	32	18,26,27	0.80	0	19,38,41	1.46	2 (10%)
32	2MG	a	959	32	18,26,27	2.82	6 (33%)	16,38,41	1.39	3 (18%)
7	6MZ	A	2268	7	18,25,26	2.22	5 (27%)	16,36,39	3.86	4 (25%)
32	G7M	a	518	32	20,26,27	3.00	3 (15%)	17,39,42	0.87	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
32	UR3	a	1491	32	19,22,23	2.67	6 (31%)	26,32,35	1.31	3 (11%)
32	MA6	a	1511	32	18,26,27	0.80	0	19,38,41	1.46	2 (10%)
32	2MG	a	1509	32	18,26,27	2.83	6 (33%)	16,38,41	1.31	3 (18%)
7	5MU	A	2177	55,7	19,22,23	2.36	8 (42%)	28,32,35	2.37	6 (21%)
7	OMC	A	2736	55,7	19,22,23	1.95	6 (31%)	26,31,34	0.93	1 (3%)
7	OMG	A	2489	52,7	18,26,27	3.10	5 (27%)	19,38,41	1.33	4 (21%)
7	OMG	A	2791	7	18,26,27	3.08	5 (27%)	19,38,41	1.29	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	6MZ	A	2296	7	-	2/5/27/28	0/3/3/3
32	5MC	a	960	32	-	0/7/25/26	0/2/2/2
32	4OC	a	1395	32	-	2/9/29/30	0/2/2/2
32	MA6	a	1512	32	-	2/7/29/30	0/3/3/3
32	2MG	a	959	32	-	2/5/27/28	0/3/3/3
7	6MZ	A	2268	7	-	2/5/27/28	0/3/3/3
32	G7M	a	518	32	-	1/3/25/26	0/3/3/3
32	UR3	a	1491	32	-	0/7/25/26	0/2/2/2
32	MA6	a	1511	32	-	1/7/29/30	0/3/3/3
32	2MG	a	1509	32	-	0/5/27/28	0/3/3/3
7	5MU	A	2177	55,7	-	0/7/25/26	0/2/2/2
7	OMC	A	2736	55,7	-	0/9/27/28	0/2/2/2
7	OMG	A	2489	52,7	-	3/5/27/28	0/3/3/3
7	OMG	A	2791	7	-	0/5/27/28	0/3/3/3

The worst 5 of 64 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
32	a	518	G7M	O6-C6	11.23	1.46	1.23
7	A	2791	OMG	O6-C6	10.17	1.44	1.23
7	A	2489	OMG	O6-C6	9.97	1.43	1.23
32	a	1395	4OC	O2-C2	9.57	1.41	1.23
7	A	2296	6MZ	C6-N6	8.48	1.48	1.35

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	2268	6MZ	C9-N6-C6	-13.16	111.53	122.87
7	A	2177	5MU	C4-N3-C2	-5.77	119.88	127.35
7	A	2177	5MU	C5-C6-N1	-5.64	117.54	123.34
7	A	2177	5MU	N3-C2-N1	5.63	122.36	114.89
7	A	2296	6MZ	C2-N1-C6	5.63	121.42	116.59

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	2296	6MZ	C5-C6-N6-C9
7	A	2296	6MZ	N1-C6-N6-C9
7	A	2489	OMG	O4'-C4'-C5'-O5'
7	A	2489	OMG	C1'-C2'-O2'-CM2
32	a	959	2MG	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 454 ligands modelled in this entry, 453 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
54	WDP	A	3201	-	83,90,90	5.19	30 (36%)	102,132,132	2.54	24 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
54	WDP	A	3201	-	-	46/103/158/158	0/4/5/5

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
54	A	3201	WDP	C04-N03	28.11	1.53	1.27
54	A	3201	WDP	O28-S26	20.37	1.66	1.43
54	A	3201	WDP	O27-S26	19.10	1.65	1.43
54	A	3201	WDP	S26-N25	9.98	1.77	1.61
54	A	3201	WDP	C18-N20	9.92	1.52	1.35

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
54	A	3201	WDP	C05-C04-N03	-14.95	109.49	126.93
54	A	3201	WDP	O17-C18-N20	8.05	120.65	109.92
54	A	3201	WDP	O28-S26-O27	-7.90	109.83	119.55
54	A	3201	WDP	C24-N25-S26	-5.55	108.87	119.59
54	A	3201	WDP	O50-C51-C53	4.66	121.78	111.56

There are no chirality outliers.

5 of 46 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
54	A	3201	WDP	C12-C14-C15-C16
54	A	3201	WDP	C12-C14-C15-O17
54	A	3201	WDP	O17-C15-C35-O36
54	A	3201	WDP	C22-C21-N20-C18
54	A	3201	WDP	C24-C21-N20-C18

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
47	q	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	q	43:LYS	C	44:GLY	N	6.14


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-22865. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 163




Y Index: 178




Z Index: 144


6.3.2 Raw map



X Index: 163



Y Index: 178




Z Index: 144


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 914 nm³; this corresponds to an approximate mass of 826 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.370 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.370 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.70	-	-
Author-provided FSC curve	2.69	2.99	2.73
Unmasked-calculated*	2.92	3.26	2.97

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-22865 and PDB model 7KGB. Per-residue inclusion information can be found in section [3](#) on page [15](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).







































































9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ





































The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8880	 0.5760
0	 0.9020	 0.6260
1	 0.9100	 0.6200
2	 0.9540	 0.6530
3	 0.9360	 0.6400
4	 0.8960	 0.6060
6	 0.6700	 0.5070
A	 0.8950	 0.5760
B	 0.9520	 0.5860
C	 0.9250	 0.6380
D	 0.9110	 0.6170
E	 0.8920	 0.6040
F	 0.8040	 0.5420
G	 0.7370	 0.5270
H	 0.1110	 0.3090
J	 0.9070	 0.6130
K	 0.8880	 0.6180
L	 0.8930	 0.6100
M	 0.8870	 0.6100
N	 0.9330	 0.6290
O	 0.8480	 0.5660
P	 0.8530	 0.6040
Q	 0.9320	 0.6250
R	 0.8600	 0.6040
S	 0.9430	 0.6270
T	 0.8230	 0.5900
U	 0.7830	 0.5560
V	 0.7740	 0.5430
W	 0.9410	 0.6300
X	 0.9110	 0.6180
Y	 0.8340	 0.5660
Z	 0.8770	 0.6200
a	 0.9480	 0.5780
c	 0.6960	 0.5350
d	 0.6470	 0.4890



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
e	 0.8030	 0.5720
f	 0.7650	 0.5460
g	 0.8040	 0.5560
h	 0.8660	 0.5750
i	 0.7490	 0.5280
j	 0.5680	 0.4690
k	 0.8600	 0.5820
l	 0.7630	 0.5600
m	 0.7460	 0.5240
n	 0.8710	 0.5920
o	 0.8390	 0.5680
p	 0.8050	 0.5520
q	 0.7540	 0.5430
r	 0.8300	 0.5710
s	 0.7790	 0.5420
t	 0.7980	 0.5550
v	 0.8800	 0.6410
y	 0.8890	 0.5630