



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 19, 2024 – 09:11 PM EDT

PDB ID : 3KXF
Title : Crystal Structure of SB27 TCR in complex with the 'restriction triad' mutant HLA-B*3508-13mer
Authors : Archbold, J.K.; Tynan, F.E.; Gras, S.; Rossjohn, J.
Deposited on : 2009-12-03
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1351 (3.10-3.10)
Clashscore	180529	1454 (3.10-3.10)
Ramachandran outliers	177936	1391 (3.10-3.10)
Sidechain outliers	177891	1391 (3.10-3.10)
RSRZ outliers	164620	1351 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 80% 18% .
1	C	276	 82% 16% .
1	I	276	 81% 18% .
1	K	276	 80% 18% .
2	B	99	 87% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	99	 89% 11%
2	J	99	 76% 24%
2	L	99	 86% 13% •
3	D	204	 77% 20% •
3	G	204	 75% 22% •
3	M	204	 81% 17% •
3	N	204	 77% 20% •
4	E	241	 78% 21%
4	H	241	 76% 22% •
4	O	241	 76% 22% •
4	P	241	 71% 25% •
5	Q	13	 85% 15%
5	R	13	 92% 8%
5	S	13	 85% 15%
5	T	13	 69% 31%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	IOD	I	280	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 26991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, B-35 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	0	0
			2246	1400	412	427	7			
1	C	276	Total	C	N	O	S	0	0	0
			2246	1400	412	427	7			
1	K	276	Total	C	N	O	S	0	0	0
			2246	1400	412	427	7			
1	I	276	Total	C	N	O	S	0	1	0
			2254	1405	415	427	7			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	65	ALA	GLN	engineered mutation	UNP P30685
A	69	ALA	THR	engineered mutation	UNP P30685
A	155	ALA	GLN	engineered mutation	UNP P30685
A	156	ARG	LEU	SEE REMARK 999	UNP P30685
C	65	ALA	GLN	engineered mutation	UNP P30685
C	69	ALA	THR	engineered mutation	UNP P30685
C	155	ALA	GLN	engineered mutation	UNP P30685
C	156	ARG	LEU	SEE REMARK 999	UNP P30685
K	65	ALA	GLN	engineered mutation	UNP P30685
K	69	ALA	THR	engineered mutation	UNP P30685
K	155	ALA	GLN	engineered mutation	UNP P30685
K	156	ARG	LEU	SEE REMARK 999	UNP P30685
I	65	ALA	GLN	engineered mutation	UNP P30685
I	69	ALA	THR	engineered mutation	UNP P30685
I	155	ALA	GLN	engineered mutation	UNP P30685
I	156	ARG	LEU	SEE REMARK 999	UNP P30685

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			
2	F	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			
2	L	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			
2	J	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			

- Molecule 3 is a protein called SB27 T cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	204	Total	C	N	O	S	0	0	0
			1637	1033	263	334	7			
3	G	204	Total	C	N	O	S	0	1	0
			1640	1035	263	335	7			
3	N	204	Total	C	N	O	S	0	0	0
			1637	1033	263	334	7			
3	M	204	Total	C	N	O	S	0	0	0
			1637	1033	263	334	7			

- Molecule 4 is a protein called SB27 T cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	241	Total	C	N	O	S	0	0	0
			1915	1203	331	372	9			
4	H	241	Total	C	N	O	S	0	0	0
			1915	1203	331	372	9			
4	P	241	Total	C	N	O	S	0	1	0
			1919	1206	331	373	9			
4	O	241	Total	C	N	O	S	0	0	0
			1914	1203	331	371	9			

- Molecule 5 is a protein called peptide from Trans-activator protein BZLF1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	Q	13	Total	C	N	O	0	0	0
			101	66	15	20			
5	R	13	Total	C	N	O	0	0	0
			101	66	15	20			
5	T	13	Total	C	N	O	0	0	0
			101	66	15	20			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	S	13	Total	C	N	O	0	0	0
			101	66	15	20			

- Molecule 6 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	I	0	0
			3	3		
6	B	1	Total	I	0	0
			1	1		
6	C	4	Total	I	0	0
			4	4		
6	D	1	Total	I	0	0
			1	1		
6	E	2	Total	I	0	0
			2	2		
6	F	1	Total	I	0	0
			1	1		
6	G	1	Total	I	0	0
			1	1		
6	H	1	Total	I	0	0
			1	1		
6	K	3	Total	I	0	0
			3	3		
6	L	1	Total	I	0	0
			1	1		
6	P	1	Total	I	0	0
			1	1		
6	I	4	Total	I	0	0
			4	4		
6	J	1	Total	I	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	O	0	0
			2	2		
7	B	2	Total	O	0	0
			2	2		
7	D	3	Total	O	0	0
			3	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	E	3	Total 3	O 3	0	0
7	G	3	Total 3	O 3	0	0
7	H	7	Total 7	O 7	0	0
7	K	3	Total 3	O 3	0	0
7	L	1	Total 1	O 1	0	0
7	N	2	Total 2	O 2	0	0
7	P	6	Total 6	O 6	0	0
7	I	1	Total 1	O 1	0	0
7	J	3	Total 3	O 3	0	0
7	M	2	Total 2	O 2	0	0
7	O	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, B-35 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, B-35 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, B-35 alpha chain





- Molecule 1: HLA class I histocompatibility antigen, B-35 alpha chain








- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin





- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



- Molecule 3: SB27 T cell receptor alpha chain




- Molecule 3: SB27 T cell receptor alpha chain





- Molecule 3: SB27 T cell receptor alpha chain

Chain N: 77% 20% .




- Molecule 3: SB27 T cell receptor alpha chain

Chain M: 81% 17% .




- Molecule 4: SB27 T cell receptor beta chain

Chain E: 78% 21% .




- Molecule 4: SB27 T cell receptor beta chain

Chain H: 76% 22% .




- Molecule 4: SB27 T cell receptor beta chain

Chain P: 71% 25% .



- Molecule 4: SB27 T cell receptor beta chain

Chain O: 76% 22%



- Molecule 5: peptide from Trans-activator protein BZLF1

Chain Q: 85% 15%



- Molecule 5: peptide from Trans-activator protein BZLF1

Chain R: 92% 8%



- Molecule 5: peptide from Trans-activator protein BZLF1

Chain T: 69% 31%



- Molecule 5: peptide from Trans-activator protein BZLF1

Chain S: 85% 15%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.69Å 207.08Å 123.51Å 90.00° 90.19° 90.00°	Depositor
Resolution (Å)	80.00 – 3.10 80.00 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (80.00-3.10) 98.6 (80.00-3.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 3.13Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.213 , 0.291 0.184 , 0.214	Depositor DCC
R_{free} test set	3612 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtriage
Anisotropy	0.384	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 25.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.265 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	26991	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2309	0.54	0/3140
1	C	0.35	0/2309	0.55	0/3140
1	I	0.34	0/2320	0.53	0/3154
1	K	0.34	0/2309	0.54	0/3140
2	B	0.36	0/852	0.49	0/1152
2	F	0.35	0/852	0.52	0/1152
2	J	0.34	0/852	0.48	0/1152
2	L	0.35	0/852	0.48	0/1152
3	D	0.37	0/1675	0.56	0/2276
3	G	0.39	0/1681	0.56	1/2284 (0.0%)
3	M	0.38	0/1675	0.52	0/2276
3	N	0.38	0/1675	0.53	0/2276
4	E	0.36	0/1967	0.53	0/2676
4	H	0.36	0/1967	0.54	0/2676
4	O	0.38	0/1966	0.54	0/2676
4	P	0.36	0/1974	0.53	0/2687
5	Q	0.39	0/104	0.52	0/142
5	R	0.36	0/104	0.51	0/142
5	S	0.31	0/104	0.49	0/142
5	T	0.37	0/104	0.48	0/142
All	All	0.36	0/27651	0.53	1/37577 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	164	LEU	CA-CB-CG	5.71	128.42	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2246	0	2111	48	0
1	C	2246	0	2111	43	0
1	I	2254	0	2124	33	0
1	K	2246	0	2111	32	0
2	B	829	0	794	8	0
2	F	829	0	794	4	0
2	J	829	0	794	13	0
2	L	829	0	794	5	0
3	D	1637	0	1532	33	0
3	G	1640	0	1537	28	0
3	M	1637	0	1532	21	0
3	N	1637	0	1532	24	0
4	E	1915	0	1809	27	0
4	H	1915	0	1809	33	0
4	O	1914	0	1809	37	0
4	P	1919	0	1813	40	0
5	Q	101	0	102	2	0
5	R	101	0	102	1	0
5	S	101	0	102	1	0
5	T	101	0	102	3	0
6	A	3	0	0	1	0
6	B	1	0	0	0	0
6	C	4	0	0	1	0
6	D	1	0	0	0	0
6	E	2	0	0	1	0
6	F	1	0	0	0	0
6	G	1	0	0	1	0
6	H	1	0	0	0	0
6	I	4	0	0	3	0
6	J	1	0	0	0	0
6	K	3	0	0	1	0
6	L	1	0	0	1	0
6	P	1	0	0	0	0
7	A	2	0	0	0	0
7	B	2	0	0	0	0
7	D	3	0	0	0	0
7	E	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	G	3	0	0	0	0
7	H	7	0	0	0	0
7	I	1	0	0	1	0
7	J	3	0	0	0	0
7	K	3	0	0	0	0
7	L	1	0	0	0	0
7	M	2	0	0	0	0
7	N	2	0	0	0	0
7	O	3	0	0	0	0
7	P	6	0	0	0	0
All	All	26991	0	25414	404	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 404 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:THR:CG2	1:A:202:ARG:HB3	1.55	1.36
1:C:195:SER:HB3	1:C:196:ASP:CB	1.65	1.26
1:A:252:GLY:CA	1:A:253:GLU:HB2	1.64	1.25
1:A:252:GLY:HA3	1:A:253:GLU:CB	1.79	1.11
3:M:202:PHE:H	3:M:203:PRO:HD2	1.07	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	274/276 (99%)	244 (89%)	27 (10%)	3 (1%)	12 39
1	C	274/276 (99%)	248 (90%)	22 (8%)	4 (2%)	8 33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	275/276 (100%)	252 (92%)	23 (8%)	0	100	100
1	K	274/276 (99%)	255 (93%)	18 (7%)	1 (0%)	30	63
2	B	97/99 (98%)	93 (96%)	4 (4%)	0	100	100
2	F	97/99 (98%)	91 (94%)	6 (6%)	0	100	100
2	J	97/99 (98%)	85 (88%)	11 (11%)	1 (1%)	13	42
2	L	97/99 (98%)	93 (96%)	2 (2%)	2 (2%)	5	25
3	D	202/204 (99%)	179 (89%)	18 (9%)	5 (2%)	4	22
3	G	203/204 (100%)	185 (91%)	16 (8%)	2 (1%)	13	42
3	M	202/204 (99%)	181 (90%)	17 (8%)	4 (2%)	6	26
3	N	202/204 (99%)	182 (90%)	17 (8%)	3 (2%)	8	33
4	E	239/241 (99%)	225 (94%)	14 (6%)	0	100	100
4	H	239/241 (99%)	218 (91%)	18 (8%)	3 (1%)	10	36
4	O	239/241 (99%)	219 (92%)	16 (7%)	4 (2%)	7	30
4	P	240/241 (100%)	224 (93%)	13 (5%)	3 (1%)	10	36
5	Q	11/13 (85%)	9 (82%)	2 (18%)	0	100	100
5	R	11/13 (85%)	9 (82%)	2 (18%)	0	100	100
5	S	11/13 (85%)	10 (91%)	0	1 (9%)	0	3
5	T	11/13 (85%)	11 (100%)	0	0	100	100
All	All	3295/3332 (99%)	3013 (91%)	246 (8%)	36 (1%)	12	39

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	253	GLU
1	C	119	ASP
3	D	128	ASP
3	G	131	SER
4	H	69	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	223 (96%)	8 (4%)	31	61
1	C	231/231 (100%)	222 (96%)	9 (4%)	27	58
1	I	232/231 (100%)	222 (96%)	10 (4%)	25	55
1	K	231/231 (100%)	220 (95%)	11 (5%)	21	51
2	B	94/94 (100%)	92 (98%)	2 (2%)	48	72
2	F	94/94 (100%)	90 (96%)	4 (4%)	25	55
2	J	94/94 (100%)	91 (97%)	3 (3%)	34	63
2	L	94/94 (100%)	89 (95%)	5 (5%)	19	48
3	D	189/189 (100%)	176 (93%)	13 (7%)	13	39
3	G	190/189 (100%)	170 (90%)	20 (10%)	5	22
3	M	189/189 (100%)	175 (93%)	14 (7%)	11	36
3	N	189/189 (100%)	170 (90%)	19 (10%)	6	23
4	E	208/208 (100%)	195 (94%)	13 (6%)	15	42
4	H	208/208 (100%)	191 (92%)	17 (8%)	9	33
4	O	208/208 (100%)	192 (92%)	16 (8%)	10	35
4	P	209/208 (100%)	187 (90%)	22 (10%)	5	22
5	Q	11/11 (100%)	11 (100%)	0	100	100
5	R	11/11 (100%)	11 (100%)	0	100	100
5	S	11/11 (100%)	11 (100%)	0	100	100
5	T	11/11 (100%)	11 (100%)	0	100	100
All	All	2935/2932 (100%)	2749 (94%)	186 (6%)	15	42

5 of 186 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	N	150	GLN
1	I	29	ASP
3	N	198	GLU
4	P	109	ARG
1	I	194	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	255	GLN
4	P	85	GLN
4	O	118	ASN
1	K	260	HIS
4	P	11	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 24 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	276/276 (100%)	-1.76	0	100	100	12, 26, 36, 43	6 (2%)
1	C	276/276 (100%)	-1.74	0	100	100	11, 24, 35, 42	7 (2%)
1	I	276/276 (100%)	-1.71	0	100	100	10, 26, 37, 43	9 (3%)
1	K	276/276 (100%)	-1.80	0	100	100	11, 24, 35, 42	7 (2%)
2	B	99/99 (100%)	-1.71	0	100	100	11, 23, 31, 41	8 (8%)
2	F	99/99 (100%)	-1.78	0	100	100	12, 23, 37, 42	5 (5%)
2	J	99/99 (100%)	-1.70	0	100	100	15, 27, 38, 45	5 (5%)
2	L	99/99 (100%)	-1.79	0	100	100	13, 24, 35, 37	7 (7%)
3	D	204/204 (100%)	-1.74	0	100	100	15, 27, 38, 44	4 (1%)
3	G	204/204 (100%)	-1.74	0	100	100	9, 24, 38, 45	5 (2%)
3	M	204/204 (100%)	-1.58	1 (0%)	87	75	11, 27, 38, 43	6 (2%)
3	N	204/204 (100%)	-1.73	0	100	100	11, 24, 37, 45	8 (3%)
4	E	241/241 (100%)	-1.81	0	100	100	11, 24, 34, 41	4 (1%)
4	H	241/241 (100%)	-1.80	0	100	100	11, 22, 35, 43	2 (0%)
4	O	241/241 (100%)	-1.74	0	100	100	12, 24, 33, 38	2 (0%)
4	P	241/241 (100%)	-1.78	0	100	100	11, 22, 33, 43	5 (2%)
5	Q	13/13 (100%)	-1.98	0	100	100	2, 4, 7, 11	0
5	R	13/13 (100%)	-1.86	0	100	100	2, 10, 24, 25	0
5	S	13/13 (100%)	-1.86	0	100	100	12, 17, 22, 23	0
5	T	13/13 (100%)	-1.88	0	100	100	5, 8, 17, 19	0
All	All	3332/3332 (100%)	-1.75	1 (0%)	100	100	2, 24, 36, 45	90 (2%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	M	203	PRO	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	IOD	C	279	1/1	0.99	0.11	154,154,154,154	0
6	IOD	E	245	1/1	0.99	0.07	149,149,149,149	0
6	IOD	H	244	1/1	0.99	0.09	151,151,151,151	0
6	IOD	I	280	1/1	0.99	0.07	114,114,114,114	0
6	IOD	J	100	1/1	0.99	0.03	115,115,115,115	0
6	IOD	C	278	1/1	1.00	0.02	39,39,39,39	0
6	IOD	A	277	1/1	1.00	0.01	31,31,31,31	0
6	IOD	C	280	1/1	1.00	0.01	15,15,15,15	0
6	IOD	D	205	1/1	1.00	0.02	89,89,89,89	0
6	IOD	E	244	1/1	1.00	0.03	50,50,50,50	0
6	IOD	A	278	1/1	1.00	0.01	14,14,14,14	0
6	IOD	F	100	1/1	1.00	0.03	93,93,93,93	0
6	IOD	G	205	1/1	1.00	0.04	73,73,73,73	0
6	IOD	A	279	1/1	1.00	0.01	21,21,21,21	0
6	IOD	K	277	1/1	1.00	0.01	18,18,18,18	0
6	IOD	K	278	1/1	1.00	0.01	10,10,10,10	0
6	IOD	K	279	1/1	1.00	0.01	32,32,32,32	0
6	IOD	L	100	1/1	1.00	0.03	69,69,69,69	0
6	IOD	P	1	1/1	1.00	0.08	86,86,86,86	0
6	IOD	I	277	1/1	1.00	0.02	37,37,37,37	0
6	IOD	I	278	1/1	1.00	0.02	40,40,40,40	0
6	IOD	I	279	1/1	1.00	0.01	28,28,28,28	0
6	IOD	B	100	1/1	1.00	0.02	74,74,74,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	IOD	C	277	1/1	1.00	0.01	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.