



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2025 – 12:05 AM EDT

PDB ID : 7KYL / pdb_00007kyl
Title : Powassan virus Envelope protein DIII in complex with neutralizing Fab POWV-80
Authors : Errico, J.M.; Nelson, C.A.; Fremont, D.H.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2020-12-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	228	<div> <div>5%</div> <div>96%</div> <div>.</div> </div>
1	X	228	<div> <div>5%</div> <div>93%</div> <div>7%</div> <div>.</div> </div>
2	L	213	<div> <div>2%</div> <div>97%</div> <div>.</div> </div>
2	Y	213	<div> <div>2%</div> <div>99%</div> <div>.</div> </div>
3	E	103	<div> <div>4%</div> <div>87%</div> <div>10%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Z	103	<div><div></div><div>6%</div><div>85%</div><div>5%</div><div>10%</div></div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POWV-80 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	228	Total	C	N	O	S	0	0	0
			1746	1115	279	344	8			
1	X	228	Total	C	N	O	S	0	0	0
			1747	1115	279	345	8			

- Molecule 2 is a protein called POWV-80 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	213	Total	C	N	O	S	0	0	0
			1633	1010	280	335	8			
2	Y	213	Total	C	N	O	S	0	0	0
			1634	1010	280	336	8			

- Molecule 3 is a protein called Envelope protein domain III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	93	Total	C	N	O	S	0	0	0
			721	460	121	134	6			
3	Z	93	Total	C	N	O	S	0	0	0
			721	460	121	134	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	297	MET	-	initiating methionine	UNP Q91LY1
Z	297	MET	-	initiating methionine	UNP Q91LY1

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	1	Total	Na	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	Y	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	211	Total	O	0	0
			211	211		
6	L	273	Total	O	0	0
			273	273		
6	E	67	Total	O	0	0
			67	67		
6	X	244	Total	O	0	0
			244	244		
6	Y	229	Total	O	0	0
			229	229		
6	Z	64	Total	O	0	0
			64	64		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: POWV-80 Fab heavy chain




- Molecule 1: POWV-80 Fab heavy chain




- Molecule 2: POWV-80 Fab light chain





- Molecule 2: POWV-80 Fab light chain



- Molecule 3: Envelope protein domain III



- Molecule 3: Envelope protein domain III



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	79.15Å 62.56Å 141.55Å 90.00° 97.97° 90.00°	Depositor
Resolution (Å)	46.67 – 2.00 46.67 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.67-2.00) 99.9 (46.67-2.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.18.2-3874	Depositor
R, R_{free}	0.190 , 0.224 0.190 , 0.224	Depositor DCC
R_{free} test set	91950 reflections (1.16%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.516	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 67.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9292	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.7443e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	H	0.27	0/1795	0.51	0/2458
1	X	0.27	0/1796	0.50	0/2458
2	L	0.27	0/1663	0.48	0/2260
2	Y	0.27	0/1664	0.48	0/2260
3	E	0.23	0/740	0.44	0/1007
3	Z	0.23	0/740	0.45	0/1007
All	All	0.26	0/8398	0.48	0/11450

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1746	0	1690	4	0
1	X	1747	0	1690	9	0
2	L	1633	0	1563	2	0
2	Y	1634	0	1563	0	0
3	E	721	0	714	2	0
3	Z	721	0	714	2	0
4	H	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	Y	1	0	0	0	0
6	E	67	0	0	1	0
6	H	211	0	0	1	0
6	L	273	0	0	0	0
6	X	244	0	0	0	0
6	Y	229	0	0	0	0
6	Z	64	0	0	0	0
All	All	9292	0	7934	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:85:MET:HB3	1:X:88:LEU:HD21	1.80	0.64
1:H:132:PRO:HB3	1:H:158:TYR:HB3	1.84	0.59
1:X:229:ILE:HG13	1:X:230:THR:HG23	1.87	0.57
3:E:391:GLN:NE2	6:E:401:HOH:O	2.34	0.54
2:L:209:ASN:HB2	2:L:212:GLU:HG3	1.90	0.53
1:X:132:PRO:HB3	1:X:158:TYR:HB3	1.92	0.51
2:L:201:THR:HG23	1:X:218:LYS:HB3	1.95	0.47
1:X:41:GLN:HB2	1:X:47:LEU:HD23	1.99	0.45
1:X:201:TRP:CG	1:X:202:PRO:HA	2.53	0.44
1:H:4:VAL:HG13	1:H:29:PHE:CD1	2.53	0.43
1:X:69:ARG:NH1	1:X:92:ASP:OD2	2.43	0.43
1:X:13:LEU:HB2	1:X:160:PRO:HG3	2.01	0.42
1:H:69:ARG:NH1	6:H:402:HOH:O	2.44	0.42
1:X:8:GLU:CD	1:X:119:GLY:H	2.23	0.42
3:Z:318:PRO:HD2	3:Z:391:GLN:HE21	1.85	0.41
1:H:53:ILE:HD11	1:H:57:GLY:HA2	2.03	0.41
3:Z:379:PRO:HB3	3:Z:396:LYS:HE2	2.02	0.40
3:E:379:PRO:HB3	3:E:396:LYS:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	226/228 (99%)	222 (98%)	4 (2%)	0	100	100
1	X	226/228 (99%)	220 (97%)	6 (3%)	0	100	100
2	L	211/213 (99%)	206 (98%)	5 (2%)	0	100	100
2	Y	211/213 (99%)	205 (97%)	6 (3%)	0	100	100
3	E	91/103 (88%)	90 (99%)	1 (1%)	0	100	100
3	Z	91/103 (88%)	90 (99%)	1 (1%)	0	100	100
All	All	1056/1088 (97%)	1033 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	198/198 (100%)	196 (99%)	2 (1%)	73	78
1	X	198/198 (100%)	198 (100%)	0	100	100
2	L	186/186 (100%)	184 (99%)	2 (1%)	70	76
2	Y	186/186 (100%)	185 (100%)	1 (0%)	86	90
3	E	82/90 (91%)	82 (100%)	0	100	100
3	Z	82/90 (91%)	81 (99%)	1 (1%)	67	73
All	All	932/948 (98%)	926 (99%)	6 (1%)	84	88

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	14	VAL
1	H	230	THR
2	L	90	ARG
2	L	190	SER
2	Y	90	ARG
3	Z	392	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	Y	1	2	7,8,9	1.91	1 (14%)	9,10,12	2.26	5 (55%)
2	PCA	L	1	2	7,8,9	1.89	1 (14%)	9,10,12	2.09	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	Y	1	2	-	0/0/11/13	0/1/1/1
2	PCA	L	1	2	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Y	1	PCA	CD-N	4.93	1.46	1.34
2	L	1	PCA	CD-N	4.87	1.46	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Y	1	PCA	CA-N-CD	-3.15	102.79	113.58
2	Y	1	PCA	OE-CD-CG	-3.03	121.31	126.72
2	L	1	PCA	OE-CD-CG	-2.88	121.58	126.72
2	L	1	PCA	CA-N-CD	-2.77	104.11	113.58
2	Y	1	PCA	CB-CA-N	2.75	110.81	103.24
2	L	1	PCA	CB-CA-N	2.68	110.60	103.24
2	Y	1	PCA	CG-CD-N	2.58	114.71	108.39
2	L	1	PCA	CG-CD-N	2.56	114.66	108.39
2	Y	1	PCA	CB-CA-C	-2.54	109.17	112.66
2	L	1	PCA	CB-CA-C	-2.30	109.50	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	228/228 (100%)	0.15	11 (4%) 36 35	19, 28, 55, 107	0
1	X	228/228 (100%)	0.16	11 (4%) 36 35	18, 27, 57, 90	0
2	L	212/213 (99%)	0.04	5 (2%) 59 58	17, 26, 54, 115	0
2	Y	212/213 (99%)	0.05	4 (1%) 66 65	19, 30, 49, 71	0
3	E	93/103 (90%)	0.47	4 (4%) 40 38	24, 40, 63, 99	0
3	Z	93/103 (90%)	0.66	6 (6%) 26 24	27, 42, 62, 94	0
All	All	1066/1088 (97%)	0.18	41 (3%) 44 42	17, 29, 56, 115	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	213	CYS	5.2
1	X	43	PRO	4.2
1	H	145	THR	4.1
1	X	230	THR	3.9
1	H	141	CYS	3.5
1	X	229	ILE	3.5
1	X	216	SER	3.3
1	H	144	THR	3.2
1	H	143	GLY	3.0
3	E	304	TYR	2.9
1	H	142	GLY	2.9
1	H	147	SER	2.9
2	L	156	ASN	2.9
3	Z	304	TYR	2.8
1	H	43	PRO	2.7
1	H	146	GLY	2.7
2	Y	213	CYS	2.6
1	H	230	THR	2.5
1	X	42	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	Z	362	ASN	2.5
1	X	147	SER	2.4
3	Z	324	ASP	2.4
3	E	338	PRO	2.4
1	X	203	SER	2.3
2	Y	203	PRO	2.3
3	Z	392	GLN	2.3
1	X	45	LYS	2.2
1	H	44	GLU	2.2
1	X	145	THR	2.2
3	E	396	LYS	2.2
3	Z	323	HIS	2.2
3	Z	340	ARG	2.2
2	Y	156	ASN	2.2
1	X	10	GLU	2.1
2	L	154	ARG	2.1
3	E	316	ARG	2.1
2	L	193	CYS	2.0
2	Y	193	CYS	2.0
1	H	173	SER	2.0
2	L	212	GLU	2.0
1	X	213	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	Y	1	8/9	0.94	0.08	26,28,34,45	0
2	PCA	L	1	8/9	0.96	0.06	18,22,23,28	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NA	H	301	1/1	0.93	0.15	40,40,40,40	0
5	CL	Y	301	1/1	0.94	0.15	60,60,60,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.