



Full wwPDB NMR Structure Validation Report ⓘ

Mar 19, 2025 – 09:06 PM EDT

PDB ID : 2L2X
Title : Thiostrepton, oxidized at CA-CB bond of residue 9
Authors : Jonker, H.R.A.; Baumann, S.; Wolf, A.; Schoof, S.; Hiller, F.; Schulte, K.W.;
Kirschner, K.N.; Schwalbe, H.; Arndt, H.-D.
Deposited on : 2010-08-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

| | | |
|--------------------------------|---|--|
| MolProbity | : | 4.02b-467 |
| Mogul | : | 2022.3.0, CSD as543be (2022) |
| Percentile statistics | : | 20231227.v01 (using entries in the PDB archive December 27th 2023) |
| wwPDB-RCI | : | v_1n_11_5_13_A (Berjanski et al., 2005) |
| PANAV | : | Wang et al. (2010) |
| wwPDB-ShiftChecker | : | v1.2 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.41.4 |


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | NMR archive (#Entries) |
|-----------------------|-----------------------------|---------------------------|
| Clashscore | 210492 | 14027 |
| Ramachandran outliers | 207382 | 12486 |
| Sidechain outliers | 206894 | 12463 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 18 | |

2 Ensemble composition and analysis ⓘ

This entry contains 20 models.

Cyrange was unable to find well-defined residues.

Error message: The number of core atoms (6) was below the domain threshold value (8).

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 189 atoms, of which 80 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Thiostrepton.

| Mol | Chain | Residues | Atoms | | | | | | Trace |
|-----|-------|----------|-------|----|----|----|----|---|-------|
| 1 | A | 18 | Total | C | H | N | O | S | 1 |
| | | | 189 | 69 | 80 | 18 | 17 | 5 | |

There are 2 discrepancies between the modelled and reference sequences:


| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| A | 0 | QUA | - | SEE REMARK 999 | UNP P0C8P8 |
| A | 17 | NH2 | - | amidation | UNP P0C8P8 |

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Thiostrepton




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Thiostrepton




4.2.2 Score per residue for model 2

- Molecule 1: Thiostrepton




4.2.3 Score per residue for model 3

- Molecule 1: Thiostrepton




4.2.4 Score per residue for model 4

- Molecule 1: Thiostrepton




4.2.5 Score per residue for model 5

- Molecule 1: Thiostrepton




4.2.6 Score per residue for model 6

- Molecule 1: Thiostrepton




4.2.7 Score per residue for model 7

- Molecule 1: Thiostrepton




4.2.8 Score per residue for model 8

- Molecule 1: Thiostrepton




4.2.9 Score per residue for model 9

- Molecule 1: Thiostrepton




4.2.10 Score per residue for model 10

- Molecule 1: Thiostrepton




4.2.11 Score per residue for model 11

- Molecule 1: Thiostrepton




4.2.12 Score per residue for model 12

- Molecule 1: Thiostrepton




4.2.13 Score per residue for model 13

- Molecule 1: Thiostrepton




4.2.14 Score per residue for model 14

- Molecule 1: Thiostrepton




4.2.15 Score per residue for model 15

- Molecule 1: Thiostrepton




4.2.16 Score per residue for model 16

- Molecule 1: Thiostrepton




4.2.17 Score per residue for model 17

- Molecule 1: Thiostrepton




4.2.18 Score per residue for model 18

- Molecule 1: Thiostrepton




4.2.19 Score per residue for model 19

- Molecule 1: Thiostrepton



4.2.20 Score per residue for model 20

- Molecule 1: Thiostrepton



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, torsion angle dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---------------|--------------------|---------|
| ARIA | refinement | 1.2 |
| CNS | structure solution | 1.1 |

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MH6, DHA, QUA, NH2, BB9, DBU, TS9

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | Chirality | Planarity |
|-----|-------|-----------|-----------|
| 1 | A | 0.0±0.0 | 0.1±0.2 |
| All | All | 0 | 1 |

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

| Mol | Chain | Res | Type | Group | Models (Total) |
|-----|-------|-----|------|---------|----------------|
| 1 | A | 13 | BB9 | Peptide | 1 |

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|-----|-------|-------|----------|----------|---------|
| 1 | A | 109 | 80 | 77 | 1±0 |
| All | All | 2180 | 1600 | 1540 | 20 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All unique clashes are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|-------------|----------------|----------|-------------|--------|-------|
| | | | | Worst | Total |
| 1:A:5:SER:H | 1:A:13:BB9:HN1 | 0.55 | 1.43 | 13 | 20 |

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-------------|-------------|------------|-------------|-----|
| 1 | A | 5/18 (28%) | 4±0 (78±6%) | 1±0 (22±6%) | 0±0 (0±0%) | 100 | 100 |
| All | All | 100/360 (28%) | 78 (78%) | 22 (22%) | 0 (0%) | 100 | 100 |

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-------------|--------------|------------|-------------|-----|
| 1 | A | 3/4 (75%) | 3±0 (100±0%) | 0±0 (0±0%) | 100 | 100 |
| All | All | 60/80 (75%) | 60 (100%) | 0 (0%) | 100 | 100 |

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

| Mol | Type | Chain | Res | Link | Bond lengths | | |
|-----|------|-------|-----|------|--------------|-----------|-------------|
| | | | | | Counts | RMSZ | #Z>2 |
| 1 | BB9 | A | 11 | 1 | 2,5,6 | 0.52±0.02 | 0±0 (0±0%) |
| 1 | BB9 | A | 9 | 1 | 2,5,6 | 0.53±0.02 | 0±0 (0±0%) |
| 1 | BB9 | A | 6 | 1 | 2,5,6 | 0.52±0.02 | 0±0 (0±0%) |
| 1 | DBU | A | 8 | 1 | 3,4,6 | 3.41±0.02 | 1±0 (33±0%) |
| 1 | DHA | A | 3 | 1 | 3,4,5 | 0.54±0.01 | 0±0 (0±0%) |
| 1 | MH6 | A | 14 | 1 | 3,3,6 | 1.24±0.03 | 0±0 (0±0%) |
| 1 | DHA | A | 16 | 1 | 3,4,5 | 0.58±0.05 | 0±0 (0±0%) |
| 1 | TS9 | A | 10 | 1 | 7,8,10 | 0.83±0.01 | 0±0 (0±0%) |
| 1 | BB9 | A | 13 | 1 | 2,4,6 | 0.67±0.02 | 0±0 (0±0%) |
| 1 | BB9 | A | 15 | 1 | 2,5,6 | 0.47±0.02 | 0±0 (0±0%) |

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

| Mol | Type | Chain | Res | Link | Bond angles | | |
|-----|------|-------|-----|------|-------------|-----------|--------------|
| | | | | | Counts | RMSZ | #Z>2 |
| 1 | BB9 | A | 11 | 1 | 1,5,7 | 0.83±0.06 | 0±0 (0±0%) |
| 1 | BB9 | A | 9 | 1 | 1,5,7 | 0.77±0.06 | 0±0 (0±0%) |
| 1 | BB9 | A | 6 | 1 | 1,5,7 | 0.69±0.04 | 0±0 (0±0%) |
| 1 | DBU | A | 8 | 1 | 3,4,7 | 1.40±0.05 | 1±0 (31±7%) |
| 1 | DHA | A | 3 | 1 | 2,4,6 | 3.16±0.02 | 1±0 (50±0%) |
| 1 | MH6 | A | 14 | 1 | 1,3,7 | 0.38±0.04 | 0±0 (0±0%) |
| 1 | DHA | A | 16 | 1 | 2,4,6 | 3.12±0.10 | 1±0 (50±0%) |
| 1 | TS9 | A | 10 | 1 | 6,12,15 | 0.69±0.01 | 0±0 (0±0%) |
| 1 | BB9 | A | 13 | 1 | 3,4,7 | 2.68±0.02 | 3±0 (100±0%) |
| 1 | BB9 | A | 15 | 1 | 1,5,7 | 0.93±0.09 | 0±0 (0±0%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-------------|-------|
| 1 | BB9 | A | 6 | 1 | - | 0±0,0,4,6 | - |
| 1 | DBU | A | 8 | 1 | - | 0±0,1,2,6 | - |
| 1 | DHA | A | 16 | 1 | - | 0±0,0,2,4 | - |
| 1 | BB9 | A | 11 | 1 | - | 0±0,0,4,6 | - |
| 1 | BB9 | A | 15 | 1 | - | 0±0,0,4,6 | - |
| 1 | BB9 | A | 9 | 1 | - | 0±0,0,4,6 | - |
| 1 | DHA | A | 3 | 1 | - | 0±0,0,2,4 | - |
| 1 | BB9 | A | 13 | 1 | - | 0±0,0,2,6 | - |
| 1 | TS9 | A | 10 | 1 | - | 0±0,9,12,16 | - |

All unique bond outliers are listed below.

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) | Models | |
|-----|-------|-----|------|-------|------|-------------|----------|--------|-------|
| | | | | | | | | Worst | Total |
| 1 | A | 8 | DBU | CA-N | 5.85 | 1.47 | 1.33 | 8 | 20 |

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) | Models | |
|-----|-------|-----|------|---------|------|-------------|----------|--------|-------|
| | | | | | | | | Worst | Total |
| 1 | A | 16 | DHA | O-C-CA | 4.53 | 117.03 | 125.53 | 2 | 20 |
| 1 | A | 3 | DHA | O-C-CA | 4.51 | 117.08 | 125.53 | 14 | 20 |
| 1 | A | 13 | BB9 | C-CA-CB | 2.79 | 126.59 | 121.45 | 19 | 20 |
| 1 | A | 13 | BB9 | C-CA-N | 2.72 | 119.42 | 116.44 | 3 | 20 |
| 1 | A | 13 | BB9 | CB-CA-N | 2.72 | 114.16 | 122.53 | 15 | 20 |
| 1 | A | 8 | DBU | CB-CA-N | 2.33 | 124.24 | 122.88 | 7 | 19 |

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided