



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2024 – 10:26 AM EDT

PDB ID : 4L3O
Title : Crystal Structure of SIRT2 in complex with the macrocyclic peptide S2iL5
Authors : Yamagata, K.; Nishimasu, H.; Ishitani, R.; Nureki, O.
Deposited on : 2013-06-06
Resolution : 2.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	<div> <div>93%</div> <div>5%</div> <div>.</div> </div>
1	B	302	<div> <div>87%</div> <div>11%</div> <div>.</div> </div>
1	C	302	<div> <div>4%</div> <div>87%</div> <div>10%</div> <div>..</div> </div>
1	D	302	<div> <div>85%</div> <div>11%</div> <div>.</div> </div>
2	E	16	<div> <div>88%</div> <div>12%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	16	 81% 19%
2	G	16	 69% 31%
2	H	16	 75% 19% 6%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10482 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	1	0
			2368	1520	393	436	19			
1	B	295	Total	C	N	O	S	0	0	0
			2329	1497	388	425	19			
1	C	294	Total	C	N	O	S	0	0	0
			2320	1491	387	423	19			
1	D	293	Total	C	N	O	S	0	0	0
			2314	1485	383	427	19			


- Molecule 2 is a protein called cyclic peptide S2iL5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	E	16	Total 144	C 90	F 3	N 27	O 23	S 1	0	0	1
2	F	16	Total 144	C 90	F 3	N 27	O 23	S 1	0	0	1
2	G	16	Total 144	C 90	F 3	N 27	O 23	S 1	0	0	1
2	H	16	Total 144	C 90	F 3	N 27	O 23	S 1	0	0	1

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	G	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	132	Total	O	0	0
			132	132		
6	B	103	Total	O	0	0
			103	103		
6	C	122	Total	O	0	0
			122	122		
6	D	96	Total	O	0	0
			96	96		
6	E	17	Total	O	0	0
			17	17		
6	F	15	Total	O	0	0
			15	15		
6	G	12	Total	O	0	0
			12	12		
6	H	10	Total	O	0	0
			10	10		



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: NAD-dependent protein deacetylase sirtuin-2





- Molecule 1: NAD-dependent protein deacetylase sirtuin-2




- Molecule 1: NAD-dependent protein deacetylase sirtuin-2




- Molecule 1: NAD-dependent protein deacetylase sirtuin-2





- Molecule 2: cyclic peptide S2iL5

Chain E:  88% 12%




- Molecule 2: cyclic peptide S2iL5

Chain F:  81% 19%




- Molecule 2: cyclic peptide S2iL5

Chain G:  69% 31%



- Molecule 2: cyclic peptide S2iL5

Chain H:  75% 19% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.18Å 135.61Å 148.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.59 – 2.52 46.59 – 2.52	Depositor EDS
% Data completeness (in resolution range)	97.1 (46.59-2.52) 97.1 (46.59-2.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.7.2 _869	Depositor
R, R_{free}	0.211 , 0.263 0.207 , 0.258	Depositor DCC
R_{free} test set	6136 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	10482	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2, EDO, ZN, FAK, ACY, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/2425	0.37	0/3269
1	B	0.22	0/2383	0.37	0/3215
1	C	0.22	0/2374	0.38	0/3204
1	D	0.22	0/2367	0.37	0/3195
2	E	0.22	0/129	0.42	0/173
2	F	0.22	0/129	0.38	0/173
2	G	0.23	0/129	0.42	0/173
2	H	0.25	0/129	0.42	0/173
All	All	0.22	0/10065	0.37	0/13575

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2368	0	2353	6	0
1	B	2329	0	2314	14	0
1	C	2320	0	2303	14	0
1	D	2314	0	2286	15	0
2	E	144	0	121	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	144	0	121	2	0
2	G	144	0	121	2	0
2	H	144	0	120	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	12	0	12	0	0
4	B	12	0	12	1	0
4	C	12	0	12	0	0
4	D	12	0	12	1	0
5	B	4	0	6	1	0
5	C	8	0	12	0	0
5	G	4	0	6	1	0
6	A	132	0	0	0	0
6	B	103	0	0	0	0
6	C	122	0	0	1	0
6	D	96	0	0	0	0
6	E	17	0	0	0	0
6	F	15	0	0	0	0
6	G	12	0	0	0	0
6	H	10	0	0	0	0
All	All	10482	0	9811	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:343:ASP:OD1	1:D:346:ARG:NH2	2.27	0.68
2:G:1010:ARG:H	5:G:1101:EDO:H22	1.59	0.67
1:C:138:LEU:HD21	1:C:169:ILE:HG13	1.81	0.62
1:D:257:LEU:HB2	1:D:282:ARG:HG2	1.83	0.60
1:B:71:MET:HE3	1:B:331:LEU:HD11	1.84	0.59

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/302 (97%)	288 (98%)	6 (2%)	0	100	100
1	B	291/302 (96%)	286 (98%)	4 (1%)	1 (0%)	41	59
1	C	290/302 (96%)	280 (97%)	10 (3%)	0	100	100
1	D	289/302 (96%)	279 (96%)	9 (3%)	1 (0%)	41	59
2	E	13/16 (81%)	10 (77%)	3 (23%)	0	100	100
2	F	13/16 (81%)	11 (85%)	2 (15%)	0	100	100
2	G	13/16 (81%)	11 (85%)	2 (15%)	0	100	100
2	H	13/16 (81%)	11 (85%)	1 (8%)	1 (8%)	1	1
All	All	1216/1272 (96%)	1176 (97%)	37 (3%)	3 (0%)	47	67

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	1002	TYR
1	D	219	PRO
1	B	219	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/263 (98%)	253 (98%)	6 (2%)	50	74
1	B	253/263 (96%)	244 (96%)	9 (4%)	35	59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	252/263 (96%)	243 (96%)	9 (4%)	35	59
1	D	252/263 (96%)	245 (97%)	7 (3%)	43	68
2	E	13/13 (100%)	12 (92%)	1 (8%)	13	23
2	F	13/13 (100%)	13 (100%)	0	100	100
2	G	13/13 (100%)	12 (92%)	1 (8%)	13	23
2	H	13/13 (100%)	12 (92%)	1 (8%)	13	23
All	All	1068/1104 (97%)	1034 (97%)	34 (3%)	39	63

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	180	GLN
1	D	183	LEU
2	G	1002	TYR
1	B	211	GLU
1	B	183	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	167	GLN
1	D	180	GLN
2	G	1006	HIS
1	D	326	GLN
2	F	1006	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAK	E	1008	2	13,14,15	1.52	2 (15%)	13,18,20	2.01	1 (7%)
2	FAK	H	1008	2	13,14,15	1.55	2 (15%)	13,18,20	1.59	1 (7%)
2	FAK	F	1008	2	13,14,15	1.52	2 (15%)	13,18,20	1.59	1 (7%)
2	FAK	G	1008	2	13,14,15	1.55	2 (15%)	13,18,20	1.86	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAK	E	1008	2	-	0/15/16/18	-
2	FAK	H	1008	2	-	3/15/16/18	-
2	FAK	F	1008	2	-	2/15/16/18	-
2	FAK	G	1008	2	-	0/15/16/18	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1008	FAK	CH-NZ	4.07	1.43	1.33
2	F	1008	FAK	CH-NZ	4.00	1.43	1.33
2	G	1008	FAK	CH-NZ	3.98	1.43	1.33
2	H	1008	FAK	CH-NZ	3.95	1.43	1.33
2	H	1008	FAK	CB-CA	-2.83	1.49	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1008	FAK	CT-CH-NZ	6.35	120.63	115.20
2	G	1008	FAK	CT-CH-NZ	5.79	120.15	115.20
2	F	1008	FAK	CT-CH-NZ	4.87	119.36	115.20
2	H	1008	FAK	CT-CH-NZ	4.80	119.30	115.20

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	1008	FAK	NZ-CH-CT-FI3
2	H	1008	FAK	NZ-CH-CT-FI3
2	F	1008	FAK	NZ-CH-CT-FI2
2	H	1008	FAK	OH-CH-CT-FI3
2	H	1008	FAK	NZ-CH-CT-FI1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MES	C	502	-	12,12,12	2.34	1 (8%)	15,16,16	2.16	7 (46%)
4	MES	A	502	-	12,12,12	2.35	1 (8%)	15,16,16	2.02	6 (40%)
4	MES	B	502	-	12,12,12	2.28	1 (8%)	15,16,16	1.86	4 (26%)
5	EDO	B	503	-	3,3,3	0.43	0	2,2,2	0.33	0
5	EDO	G	1101	-	3,3,3	0.42	0	2,2,2	0.39	0
4	MES	D	502	-	12,12,12	2.31	1 (8%)	15,16,16	1.94	4 (26%)
5	EDO	C	504	-	3,3,3	0.45	0	2,2,2	0.34	0
5	EDO	C	503	-	3,3,3	0.43	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	C	502	-	-	1/6/14/14	0/1/1/1
4	MES	A	502	-	-	1/6/14/14	0/1/1/1
4	MES	B	502	-	-	0/6/14/14	0/1/1/1
5	EDO	B	503	-	-	0/1/1/1	-
5	EDO	G	1101	-	-	0/1/1/1	-
4	MES	D	502	-	-	1/6/14/14	0/1/1/1
5	EDO	C	504	-	-	0/1/1/1	-
5	EDO	C	503	-	-	0/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	MES	C8-S	-7.86	1.66	1.77
4	C	502	MES	C8-S	-7.81	1.66	1.77
4	D	502	MES	C8-S	-7.73	1.66	1.77
4	B	502	MES	C8-S	-7.64	1.66	1.77

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	MES	C5-N4-C3	4.74	119.05	108.84
4	C	502	MES	C5-N4-C3	4.41	118.34	108.84
4	D	502	MES	C5-N4-C3	4.37	118.26	108.84
4	B	502	MES	C5-N4-C3	4.18	117.84	108.84
4	C	502	MES	C6-C5-N4	-3.71	104.49	110.12

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	MES	C8-C7-N4-C3
4	C	502	MES	C8-C7-N4-C3
4	D	502	MES	C8-C7-N4-C3

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	502	MES	1	0
5	B	503	EDO	1	0
5	G	1101	EDO	1	0
4	D	502	MES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/302 (98%)	-0.38	2 (0%) 87 89	13, 25, 58, 71	0
1	B	295/302 (97%)	-0.11	4 (1%) 75 78	15, 38, 70, 101	0
1	C	294/302 (97%)	-0.11	12 (4%) 37 41	12, 29, 73, 91	0
1	D	293/302 (97%)	-0.14	2 (0%) 87 89	17, 42, 65, 82	0
2	E	13/16 (81%)	-0.40	0 100 100	15, 20, 26, 27	0
2	F	13/16 (81%)	-0.11	0 100 100	16, 19, 26, 26	0
2	G	13/16 (81%)	-0.24	0 100 100	15, 23, 29, 29	0
2	H	13/16 (81%)	-0.10	0 100 100	17, 24, 32, 37	0
All	All	1231/1272 (96%)	-0.19	20 (1%) 72 74	12, 32, 67, 101	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	110	TYR	4.2
1	C	214	PHE	3.7
1	B	99	PRO	3.6
1	C	210	LYS	3.2
1	C	107	LEU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FAK	G	1008	15/16	0.95	0.15	11,20,27,29	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FAK	F	1008	15/16	0.96	0.14	15,18,32,39	0
2	FAK	E	1008	15/16	0.97	0.12	10,15,23,30	0
2	FAK	H	1008	15/16	0.97	0.11	16,23,36,37	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ZN	C	501	1/1	0.89	0.07	70,70,70,70	0
4	MES	A	502	12/12	0.93	0.21	35,59,64,67	0
5	EDO	B	503	4/4	0.93	0.14	35,36,42,42	0
5	EDO	C	503	4/4	0.93	0.17	24,28,29,38	0
4	MES	D	502	12/12	0.94	0.22	47,55,67,74	0
3	ZN	D	501	1/1	0.95	0.04	58,58,58,58	0
4	MES	B	502	12/12	0.95	0.19	38,58,68,72	0
4	MES	C	502	12/12	0.95	0.14	40,52,63,68	0
5	EDO	C	504	4/4	0.96	0.39	30,31,40,42	0
5	EDO	G	1101	4/4	0.96	0.14	27,33,34,37	0
3	ZN	A	501	1/1	0.98	0.04	66,66,66,66	0
3	ZN	B	501	1/1	0.99	0.07	58,58,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.