



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2024 – 11:23 AM EST

PDB ID : 3L4W  
Title : Crystal complex of N-terminal Human Maltase-Glucoamylase with miglitol  
Authors : Sim, L.; Rose, D.R.  
Deposited on : 2009-12-21  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	875	
2	B	2	
2	C	2	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltase-glucoamylase, intestinal.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	863	Total	C	N	O	S	0	5	0
			6949	4432	1176	1314	27			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	772	ASP	ASN	variant	UNP O43451
A	869	ALA	-	expression tag	UNP O43451
A	870	HIS	-	expression tag	UNP O43451
A	871	HIS	-	expression tag	UNP O43451
A	872	HIS	-	expression tag	UNP O43451
A	873	HIS	-	expression tag	UNP O43451
A	874	HIS	-	expression tag	UNP O43451
A	875	HIS	-	expression tag	UNP O43451

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is (2R,3R,4R,5S)-1-(2-hydroxyethyl)-2-(hydroxymethyl)piperidine-3,4,5-triol (three-letter code: MIG) (formula: C<sub>8</sub>H<sub>17</sub>NO<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	434	Total	O	0	0
			434	434		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltase-glucoamylase, intestinal



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.71Å 107.85Å 111.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.92 – 2.00 19.92 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.9 (19.92-2.00) 96.8 (19.92-2.00)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.02 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.198 , 0.232 0.197 , 0.229	Depositor DCC
$R_{free}$ test set	3759 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.0	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 51.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7505	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, MIG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/7158	0.71	2/9764 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	624	ARG	NE-CZ-NH2	-6.11	117.25	120.30
1	A	624	ARG	NE-CZ-NH1	5.90	123.25	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6949	0	6614	81	0
2	B	28	0	24	0	0
2	C	28	0	25	0	0
3	A	14	0	17	0	0
4	A	28	0	26	0	0
5	A	24	0	32	3	0
6	A	434	0	0	6	0
All	All	7505	0	6738	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:HIS:HE1	1:A:261:ASP:OD1	1.40	1.03
1:A:98:HIS:CE1	1:A:261:ASP:OD1	2.28	0.86
1:A:204:THR:HG21	1:A:473:LEU:HD23	1.60	0.84
1:A:204:THR:HG21	1:A:473:LEU:CD2	2.12	0.79
1:A:204:THR:CG2	1:A:473:LEU:HD23	2.14	0.77
1:A:202:ARG:HG3	1:A:204:THR:HG23	1.68	0.76
1:A:292:LEU:HD23	1:A:567:MET:CE	2.17	0.74
1:A:54:VAL:HB	1:A:135:ARG:HB3	1.70	0.74
1:A:300:GLU:HG3	1:A:603:GLN:HE21	1.52	0.73
1:A:183:HIS:HE1	1:A:198:PRO:O	1.73	0.70
1:A:201:ASN:O	1:A:497:HIS:HE1	1.74	0.70
1:A:204:THR:CG2	1:A:473:LEU:CD2	2.70	0.69
1:A:453:GLY:HA3	1:A:458:CYS:SG	2.35	0.67
1:A:464:ASN:ND2	1:A:484:MET:H	1.96	0.64
1:A:464:ASN:HD21	1:A:484:MET:H	1.46	0.62
1:A:493:GLN:HE21	1:A:497:HIS:HD2	1.46	0.61
1:A:576:ALA:O	1:A:577:LEU:HB2	2.00	0.60
1:A:493:GLN:NE2	1:A:497:HIS:HD2	1.98	0.60
1:A:841:THR:HB	1:A:854:THR:HG22	1.83	0.60
1:A:335:ARG:HG2	1:A:407:PRO:HB3	1.85	0.59
1:A:544:THR:CG2	1:A:577:LEU:HD22	2.33	0.58
1:A:712:ARG:H	1:A:714:GLN:HE21	1.53	0.57
1:A:164:GLN:HE22	1:A:462:ASN:HB2	1.70	0.56
1:A:292:LEU:HD23	1:A:567:MET:HE1	1.86	0.56
1:A:794:ASN:HD21	1:A:870:HIS:CD2	2.23	0.56
1:A:167:GLN:NE2	1:A:254:ARG:HE	2.05	0.55
1:A:794:ASN:HB3	1:A:868:TRP:O	2.05	0.55
1:A:179:GLY:O	1:A:190:HIS:HE1	1.89	0.55
1:A:360:LYS:HE2	6:A:4367:HOH:O	2.07	0.54
1:A:201:ASN:H	1:A:498:ASN:ND2	2.06	0.54
1:A:243:VAL:HG11	1:A:251:ILE:HD11	1.89	0.54
1:A:53:HIS:HD2	1:A:72:LEU:HD22	1.72	0.53
1:A:56:GLY:O	1:A:135:ARG:NH2	2.40	0.53
1:A:204:THR:HG21	1:A:473:LEU:HD22	1.90	0.53
1:A:300:GLU:HG3	1:A:603:GLN:HG3	1.91	0.53
1:A:591:GLY:HA2	1:A:594:TYR:CD2	2.45	0.52
1:A:9:GLU:HB2	1:A:41:VAL:HG11	1.90	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:ASN:H	1:A:498:ASN:HD21	1.57	0.52
1:A:243:VAL:CG1	1:A:251:ILE:HD11	2.41	0.51
1:A:682:PRO:HA	1:A:725:ILE:HG13	1.91	0.51
1:A:90:GLU:OE1	1:A:98:HIS:HD2	1.94	0.51
1:A:291:ALA:O	1:A:295:HIS:HE1	1.95	0.49
1:A:476[A]:TYR:CE2	1:A:479:CYS:HB2	2.47	0.49
1:A:618:LEU:HD11	1:A:723:ASP:HB3	1.93	0.49
1:A:488:GLN:HB2	1:A:496:ILE:HD11	1.94	0.49
1:A:673:LEU:HA	1:A:678:LEU:O	2.13	0.49
1:A:493:GLN:NE2	1:A:497:HIS:CD2	2.79	0.48
1:A:402:ILE:HD11	6:A:4324:HOH:O	2.12	0.48
1:A:300:GLU:HG3	1:A:603:GLN:NE2	2.25	0.48
1:A:460:THR:HA	1:A:465:ASN:ND2	2.29	0.48
1:A:310:VAL:HG23	5:A:3003:GOL:H12	1.96	0.48
1:A:201:ASN:O	1:A:497:HIS:CE1	2.61	0.47
1:A:446:GLU:N	1:A:447:VAL:HA	2.29	0.47
1:A:681:THR:HG22	1:A:691:VAL:HG11	1.98	0.46
1:A:424:TRP:CE2	1:A:428:PHE:HE2	2.34	0.46
1:A:827:VAL:HG22	1:A:868:TRP:HB3	1.95	0.46
1:A:493:GLN:HE21	1:A:497:HIS:CD2	2.31	0.45
1:A:131:VAL:O	5:A:3004:GOL:H32	2.16	0.45
1:A:204:THR:HG22	1:A:473:LEU:CD2	2.47	0.45
1:A:42:PRO:HD3	1:A:194:TRP:CH2	2.52	0.45
1:A:60:ASN:ND2	6:A:4032:HOH:O	2.49	0.45
1:A:631:TYR:HA	1:A:634:LEU:HG	1.99	0.45
1:A:143:THR:HG23	6:A:4184:HOH:O	2.17	0.44
1:A:550:LEU:HD13	1:A:587:TRP:CG	2.52	0.44
1:A:202:ARG:CG	1:A:204:THR:HG23	2.43	0.44
1:A:295:HIS:HD2	1:A:325:HIS:NE2	2.16	0.44
1:A:606:LYS:HE3	6:A:4422:HOH:O	2.18	0.43
1:A:53:HIS:HD2	1:A:72:LEU:CD2	2.30	0.43
1:A:707:SER:HB3	1:A:848:LEU:HG	2.01	0.43
1:A:417:ASN:HA	1:A:418:PRO:HD2	1.90	0.43
1:A:317:ALA:HB1	1:A:620:LEU:HD21	2.01	0.42
1:A:711:TRP:CG	1:A:716:VAL:HG11	2.54	0.42
1:A:794:ASN:ND2	1:A:870:HIS:CD2	2.87	0.42
1:A:791:VAL:HA	1:A:795:ARG:O	2.20	0.42
1:A:794:ASN:HD21	1:A:870:HIS:HD2	1.66	0.42
1:A:239:ASN:ND2	6:A:4139:HOH:O	2.53	0.41
1:A:536:ALA:HA	5:A:3001:GOL:H11	2.02	0.41
1:A:476[A]:TYR:CD2	1:A:479:CYS:HB2	2.55	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:412:PHE:HA	1:A:413:PRO:HD3	1.96	0.41
1:A:792:THR:O	1:A:792:THR:OG1	2.37	0.41
1:A:526:ARG:O	1:A:538:HIS:CE1	2.74	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	864/875 (99%)	825 (96%)	36 (4%)	3 (0%)	37 35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	180	LEU
1	A	405	VAL
1	A	73	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	761/767 (99%)	747 (98%)	14 (2%)	54 59

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	GLU
1	A	135	ARG
1	A	161	PHE
1	A	187[A]	GLN
1	A	187[B]	GLN
1	A	193	ASN
1	A	281	ILE
1	A	322	ASP
1	A	349	GLU
1	A	616	ASP
1	A	788	GLU
1	A	801	SER
1	A	848	LEU
1	A	854	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	A	60	ASN
1	A	62	ASN
1	A	98	HIS
1	A	107	ASN
1	A	117	GLN
1	A	164	GLN
1	A	167	GLN
1	A	183	HIS
1	A	190	HIS
1	A	193	ASN
1	A	239	ASN
1	A	295	HIS
1	A	357	ASN
1	A	434	GLN
1	A	464	ASN
1	A	465	ASN
1	A	493	GLN
1	A	497	HIS
1	A	498	ASN
1	A	603	GLN
1	A	625	HIS
1	A	669	HIS
1	A	714	GLN
1	A	794	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	802	GLN
1	A	870	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	B	1	1,2	14,14,15	0.85	0	17,19,21	2.29	6 (35%)
2	NAG	B	2	2	14,14,15	1.10	1 (7%)	17,19,21	3.05	8 (47%)
2	NAG	C	1	1,2	14,14,15	0.59	0	17,19,21	1.44	2 (11%)
2	NAG	C	2	2	14,14,15	0.62	0	17,19,21	1.18	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	B	2	2	-	3/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	C1-C2	3.15	1.56	1.52

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C1-O5-C5	8.93	124.15	112.19
2	B	2	NAG	C4-C3-C2	-5.59	102.83	111.02
2	B	1	NAG	C4-C3-C2	4.47	117.58	111.02
2	C	1	NAG	O5-C1-C2	-4.25	104.71	111.29
2	B	1	NAG	O5-C1-C2	-4.18	104.83	111.29
2	B	2	NAG	C2-N2-C7	3.92	128.16	122.90
2	B	1	NAG	O4-C4-C5	-3.58	100.50	109.32
2	B	2	NAG	O5-C5-C4	2.90	117.89	110.83
2	B	1	NAG	C2-N2-C7	2.56	126.33	122.90
2	B	1	NAG	O3-C3-C2	-2.55	104.11	109.40
2	B	1	NAG	C1-O5-C5	2.35	115.34	112.19
2	C	2	NAG	C3-C4-C5	-2.33	106.00	110.23
2	B	2	NAG	C1-C2-N2	2.22	113.93	110.43
2	B	2	NAG	O5-C5-C6	2.08	111.71	107.66
2	C	1	NAG	C1-O5-C5	2.04	114.92	112.19
2	B	2	NAG	C6-C5-C4	-2.03	108.03	113.02
2	B	2	NAG	O3-C3-C2	2.01	113.58	109.40

There are no chirality outliers.


All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C3-C2-N2-C7
2	B	2	NAG	C4-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	B	2	NAG	O5-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	3001	-	5,5,5	0.48	0	5,5,5	0.46	0
5	GOL	A	3004	-	5,5,5	0.43	0	5,5,5	0.43	0
4	NAG	A	2006	1	14,14,15	0.55	0	17,19,21	0.89	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	3003	-	5,5,5	0.26	0	5,5,5	0.47	0
5	GOL	A	3002	-	5,5,5	0.28	0	5,5,5	0.44	0
3	MIG	A	1001	-	14,14,14	0.89	1 (7%)	18,19,19	1.21	1 (5%)
4	NAG	A	2005	1	14,14,15	0.67	0	17,19,21	1.16	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	3001	-	-	1/4/4/4	-
5	GOL	A	3004	-	-	4/4/4/4	-
4	NAG	A	2006	1	-	2/6/23/26	0/1/1/1
5	GOL	A	3003	-	-	2/4/4/4	-
5	GOL	A	3002	-	-	1/4/4/4	-
3	MIG	A	1001	-	-	1/5/25/25	0/1/1/1
4	NAG	A	2005	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	MIG	CAI-CAJ	2.01	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2005	NAG	C1-O5-C5	2.90	116.08	112.19
3	A	1001	MIG	CAJ-CAI-NAN	-2.00	107.00	110.72

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	3003	GOL	C1-C2-C3-O3
5	A	3004	GOL	C1-C2-C3-O3
3	A	1001	MIG	OAA-CAF-CAH-NAN
5	A	3002	GOL	C1-C2-C3-O3
5	A	3004	GOL	O1-C1-C2-C3
5	A	3003	GOL	O2-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
5	A	3004	GOL	O1-C1-C2-O2
5	A	3004	GOL	O2-C2-C3-O3
4	A	2006	NAG	C8-C7-N2-C2
4	A	2006	NAG	O7-C7-N2-C2
5	A	3001	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	3001	GOL	1	0
5	A	3004	GOL	1	0
5	A	3003	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	863/875 (98%)	-0.04	29 (3%)	48 46	12, 29, 46, 67	5 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	72	LEU	4.9
1	A	73	PRO	4.7
1	A	135	ARG	4.3
1	A	836	SER	4.3
1	A	76	PRO	3.4
1	A	793	GLN	3.4
1	A	376	SER	3.3
1	A	204	THR	3.1
1	A	191	ASP	3.1
1	A	7	VAL	3.1
1	A	794	ASN	2.9
1	A	838	THR	2.9
1	A	77	VAL	2.9
1	A	719	GLU	2.8
1	A	835	PRO	2.8
1	A	36	GLN	2.8
1	A	83	ASP	2.7
1	A	518	ASN	2.6
1	A	50	HIS	2.5
1	A	462	ASN	2.4
1	A	26	CYS	2.3
1	A	792	THR	2.3
1	A	834	VAL	2.3
1	A	74	SER	2.3
1	A	692	MET	2.2
1	A	476[A]	TYR	2.1
1	A	81	ASN	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	847	ASN	2.0
1	A	616	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	B	2	14/15	0.68	0.18	57,62,64,64	0
2	NAG	C	2	14/15	0.71	0.15	55,57,60,61	0
2	NAG	B	1	14/15	0.83	0.12	32,36,45,46	0
2	NAG	C	1	14/15	0.92	0.09	43,46,49,51	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain B:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	2006	14/15	0.69	0.14	56,60,61,63	0
4	NAG	A	2005	14/15	0.79	0.11	50,53,56,56	0
5	GOL	A	3003	6/6	0.86	0.11	38,41,41,45	0
5	GOL	A	3001	6/6	0.88	0.11	35,40,41,44	0
5	GOL	A	3004	6/6	0.90	0.10	45,50,52,52	0
5	GOL	A	3002	6/6	0.93	0.10	31,32,34,35	0
3	MIG	A	1001	14/14	0.95	0.08	20,23,33,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.