



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 11, 2024 – 06:21 PM EDT

PDB ID : 1L7D
Title : Crystal Structure of R. rubrum Transhydrogenase Domain I without Bound NAD(H)
Authors : Prasad, G.S.; Wahlberg, M.; Sridhar, V.; Yamaguchi, M.; Hatefi, Y.; Stout, C.D.
Deposited on : 2002-03-14
Resolution : 1.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	384	 70% 20% • • 8%
1	B	384	 73% 19% • 7%
1	C	384	 69% 20% • 8%
1	D	384	 70% 21% • 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called nicotinamide nucleotide Transhydrogenase, subunit alpha 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	0	0
			2589	1638	448	487	16			
1	B	358	Total	C	N	O	S	0	0	0
			2620	1656	454	494	16			
1	C	355	Total	C	N	O	S	0	0	0
			2604	1648	451	489	16			
1	D	353	Total	C	N	O	S	0	0	0
			2592	1640	449	487	16			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	184	Total	O	0	0
			184	184		
2	B	243	Total	O	0	0
			243	243		
2	C	291	Total	O	0	0
			291	291		
2	D	221	Total	O	0	0
			221	221		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: nicotinamide nucleotide Transhydrogenase, subunit alpha 1

Chain A: 




- Molecule 1: nicotinamide nucleotide Transhydrogenase, subunit alpha 1

Chain B: 




- Molecule 1: nicotinamide nucleotide Transhydrogenase, subunit alpha 1

Chain C: 





● Molecule 1: nicotinamide nucleotide Transhydrogenase, subunit alpha 1



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.96Å 117.13Å 94.23Å 90.00° 108.26° 90.00°	Depositor
Resolution (Å)	50.00 – 1.81	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-1.81)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.220 , 0.262	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11344	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.49	4/2623 (0.2%)	0.76	8/3560 (0.2%)
1	B	0.32	0/2654	0.62	1/3600 (0.0%)
1	C	0.37	2/2638 (0.1%)	0.69	5/3578 (0.1%)
1	D	0.35	1/2626 (0.0%)	0.66	4/3561 (0.1%)
All	All	0.38	7/10541 (0.1%)	0.69	18/14299 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	3	6
1	B	1	0
All	All	4	6

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	348	ASP	CA-C	-10.05	1.26	1.52
1	A	348	ASP	C-N	8.76	1.54	1.34
1	A	377	PRO	N-CD	6.40	1.56	1.47
1	C	1150	ASP	C-N	5.68	1.47	1.34
1	A	347	VAL	C-N	-5.62	1.21	1.34

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	348	ASP	CA-C-N	-10.66	93.76	117.20
1	A	348	ASP	O-C-N	9.64	138.13	122.70
1	D	1576	HIS	C-N-CD	9.34	148.01	128.40
1	A	377	PRO	CA-N-CD	-8.41	99.72	111.50
1	C	1150	ASP	O-C-N	7.55	134.78	122.70

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	374	ILE	CB
1	A	375	VAL	CA
1	A	376	HIS	CA
1	B	401	MET	CA

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	347	VAL	Mainchain
1	A	350	ASP	Mainchain
1	A	375	VAL	Mainchain,Peptide
1	A	376	HIS	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2589	0	2714	85	1
1	B	2620	0	2754	67	0
1	C	2604	0	2744	80	0
1	D	2592	0	2730	68	0
2	A	184	0	0	3	0
2	B	243	0	0	5	1
2	C	291	0	0	7	0
2	D	221	0	0	1	0
All	All	11344	0	10942	291	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 291 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:ARG:H	1:A:127:ARG:HH21	1.11	0.94
1:B:401:MET:N	1:B:469:ASP:OD2	2.07	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:837:GLN:HE21	1:C:859:SER:HA	1.41	0.84
1:D:1286:GLU:HA	1:D:1289:LEU:HD23	1.60	0.83
1:D:1308:VAL:O	1:D:1312:LEU:HD23	1.79	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:377:PRO:O	2:B:2379:HOH:O[2_545]	1.71	0.49

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/384 (91%)	329 (94%)	20 (6%)	1 (0%)	41	27
1	B	354/384 (92%)	340 (96%)	14 (4%)	0	100	100
1	C	351/384 (91%)	332 (95%)	14 (4%)	5 (1%)	11	3
1	D	349/384 (91%)	337 (97%)	12 (3%)	0	100	100
All	All	1404/1536 (91%)	1338 (95%)	60 (4%)	6 (0%)	34	21

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	376	HIS
1	C	1150	ASP
1	C	866	SER
1	C	892	GLU
1	C	906	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/296 (93%)	266 (97%)	8 (3%)	42	28
1	B	278/296 (94%)	271 (98%)	7 (2%)	47	33
1	C	277/296 (94%)	267 (96%)	10 (4%)	35	19
1	D	276/296 (93%)	271 (98%)	5 (2%)	59	48
All	All	1105/1184 (93%)	1075 (97%)	30 (3%)	44	30

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	685	LYS
1	D	1337	LEU
1	C	999	MET
1	D	1571	ASP
1	C	1150	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	1342	ASN
1	D	1386	GLN
1	D	1538	ASN
1	B	610	GLN
1	B	542	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.