



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2024 – 05:55 PM EST

PDB ID : 3LAC
Title : Crystal structure of Bacillus anthracis pyrrolidone-carboxylate peptidase, pcP
Authors : Anderson, S.M.; Wawrzak, Z.; Onopriyenko, O.; Hasseman, J.; Edwards, A.; Savchenko, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2010-01-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	215	 76% 19% 5%
1	B	215	 84% 11% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	A	217	-	-	X	-
2	PEG	A	218	-	-	X	-
2	PEG	B	221	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyrrolidone-carboxylate peptidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	Se	0	2	0
			1593	1021	263	302	2	5			
1	B	204	Total	C	N	O	S	Se	0	1	0
			1585	1016	260	302	2	5			

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Mg 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	127	Total O 128 128	0	1
4	B	160	Total O 162 162	0	2


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyrrolidone-carboxylate peptidase



- Molecule 1: Pyrrolidone-carboxylate peptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	78.55Å 78.55Å 141.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.30 – 2.00 34.30 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.0 (34.30-2.00) 99.1 (34.30-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.181 , 0.223 0.207 , 0.249	Depositor DCC
R_{free} test set	1532 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3533	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CSO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/1623	0.77	1/2194 (0.0%)
1	B	0.73	0/1612	0.84	0/2180
All	All	0.67	0/3235	0.80	1/4374 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	ARG	NE-CZ-NH2	-5.96	117.32	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1593	0	1610	33	0
1	B	1585	0	1597	24	2
2	A	28	0	40	14	0
2	B	35	0	50	7	0
3	B	2	0	0	0	0
4	A	128	0	0	4	0
4	B	162	0	0	2	0
All	All	3533	0	3297	62	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:GLN:HE22	2:A:217:PEG:H22	0.99	1.13
1:B:154:LEU:HD22	1:B:161:MSE:HE3	1.48	0.95
1:A:41:GLN:NE2	2:A:217:PEG:H22	1.80	0.95
1:A:141:PHE:HB2	2:A:218:PEG:H11	1.53	0.91
2:A:218:PEG:H22	4:A:239:HOH:O	1.74	0.88
1:A:186:LEU:HD23	1:A:190[A]:ARG:HH12	1.38	0.88
1:A:41:GLN:HE22	2:A:217:PEG:C2	1.85	0.85
1:B:154:LEU:CD2	1:B:161:MSE:HE3	2.09	0.82
1:A:28:HIS:HD2	1:A:39:SER:H	1.30	0.80
1:A:70:GLN:HE22	1:A:174:GLN:HE22	1.39	0.71
1:A:99:PRO:HB2	1:A:102:VAL:HG22	1.75	0.68
1:B:143:CSO:HB2	1:B:165:PHE:CZ	2.32	0.65
1:B:70:GLN:HB3	2:B:221:PEG:H12	1.80	0.62
1:A:28:HIS:CD2	1:A:39:SER:H	2.15	0.60
1:B:18:ASN:HD21	2:B:221:PEG:H22	1.67	0.60
1:A:119:LYS:HG2	2:A:219:PEG:H32	1.84	0.59
1:B:65:ILE:N	1:B:65:ILE:HD12	2.17	0.59
1:B:6:LEU:HD12	1:B:66:ILE:HB	1.84	0.59
1:B:54:LYS:O	1:B:58:GLU:HG3	2.03	0.58
1:B:70:GLN:CB	2:B:221:PEG:H12	2.34	0.58
1:A:11:PRO:HD3	2:A:217:PEG:H21	1.85	0.58
1:A:26:SER:OG	1:A:190[A]:ARG:HD3	2.04	0.58
1:A:102:VAL:HG23	4:A:273:HOH:O	2.05	0.56
1:B:143:CSO:SG	2:B:221:PEG:H42	2.46	0.55
1:A:19:PRO:O	1:A:23:VAL:HG23	2.07	0.55
1:A:99:PRO:HB2	1:A:102:VAL:CG2	2.37	0.54
1:B:154:LEU:CD2	1:B:161:MSE:CE	2.83	0.54
1:A:154:LEU:CD2	1:A:161:MSE:HE3	2.38	0.53
1:B:50:ILE:HG22	1:B:54:LYS:HE2	1.91	0.53
1:B:3:THR:OG1	1:B:62:PRO:HA	2.08	0.52
1:A:6:LEU:HD22	1:A:68:ILE:HD11	1.90	0.52
1:B:130:ILE:HD11	1:B:191:LYS:HD3	1.91	0.52
1:B:162:LYS:NZ	1:B:204:GLU:OE1	2.41	0.51
2:A:216:PEG:H31	4:A:343:HOH:O	2.10	0.51
1:A:154:LEU:HD22	1:A:161:MSE:HE3	1.95	0.49
1:B:18:ASN:HD21	2:B:221:PEG:C2	2.26	0.49
2:A:219:PEG:H22	2:A:219:PEG:H42	1.60	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:GLN:O	2:B:221:PEG:H41	2.14	0.48
1:B:58:GLU:CB	4:B:273:HOH:O	2.61	0.48
1:A:70:GLN:NE2	1:A:174:GLN:HE22	2.09	0.47
1:B:58:GLU:HB2	4:B:273:HOH:O	2.14	0.47
1:A:46:PHE:O	1:A:47:HIS:HB2	2.14	0.47
1:A:157:HIS:CG	1:A:161:MSE:HE2	2.51	0.46
1:A:9:PHE:CD1	1:A:42:VAL:HG23	2.51	0.45
2:A:218:PEG:H41	2:A:218:PEG:H21	1.80	0.45
1:A:119:LYS:HB3	2:A:219:PEG:H41	1.98	0.45
1:A:30:LYS:HE2	1:A:194:GLU:OE2	2.17	0.45
1:A:11:PRO:CD	2:A:217:PEG:H21	2.46	0.45
2:A:218:PEG:C2	4:A:239:HOH:O	2.49	0.44
1:A:27:LEU:HD23	1:A:27:LEU:HA	1.89	0.44
1:A:186:LEU:CD2	1:A:190[A]:ARG:HH12	2.19	0.44
1:A:65:ILE:HD12	1:A:65:ILE:N	2.34	0.43
1:A:143:CSO:HB2	1:A:165:PHE:CZ	2.53	0.43
1:A:93:ASP:OD1	1:A:93:ASP:C	2.57	0.43
1:B:90[B]:ARG:NE	1:B:90[B]:ARG:HA	2.34	0.43
1:A:89:ALA:HA	2:A:218:PEG:H12	2.01	0.43
1:B:23:VAL:HG23	1:B:186:LEU:HD12	2.00	0.43
1:B:77:ILE:O	1:B:132:ALA:HA	2.19	0.42
1:A:77:ILE:O	1:A:132:ALA:HA	2.19	0.42
1:B:18:ASN:HD21	2:B:221:PEG:C1	2.33	0.41
1:B:162:LYS:CE	1:B:204:GLU:OE1	2.69	0.40
1:A:64:PHE:HA	1:A:162:LYS:O	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:GLU:CA	1:B:107:GLU:CB[5_556]	1.98	0.22
1:B:107:GLU:O	1:B:107:GLU:CB[5_556]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/215 (94%)	197 (97%)	6 (3%)	0	100	100
1	B	202/215 (94%)	198 (98%)	4 (2%)	0	100	100
All	All	405/430 (94%)	395 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/177 (99%)	172 (98%)	3 (2%)	56	61
1	B	174/177 (98%)	173 (99%)	1 (1%)	84	88
All	All	349/354 (99%)	345 (99%)	4 (1%)	67	76

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	SER
1	A	36	LYS
1	A	204	GLU
1	B	186	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	41	GLN
1	A	70	GLN
1	B	18	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	143	1	3,6,7	0.36	0	1,6,8	0.92	0
1	CSO	B	143	1	3,6,7	1.06	0	1,6,8	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	143	1	-	0/1/5/7	-
1	CSO	B	143	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	143	CSO	1	0
1	B	143	CSO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	A	219	-	6,6,6	0.46	0	5,5,5	0.35	0
2	PEG	B	218	-	6,6,6	0.58	0	5,5,5	0.73	0
2	PEG	A	217	-	6,6,6	0.53	0	5,5,5	0.28	0
2	PEG	B	219	-	6,6,6	0.48	0	5,5,5	0.21	0
2	PEG	A	216	-	6,6,6	0.51	0	5,5,5	0.39	0
2	PEG	A	218	-	6,6,6	0.54	0	5,5,5	0.26	0
2	PEG	B	222	-	6,6,6	0.53	0	5,5,5	0.24	0
2	PEG	B	220	-	6,6,6	0.49	0	5,5,5	0.28	0
2	PEG	B	221	-	6,6,6	0.63	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	219	-	-	3/4/4/4	-
2	PEG	B	218	-	-	4/4/4/4	-
2	PEG	A	217	-	-	3/4/4/4	-
2	PEG	B	219	-	-	2/4/4/4	-
2	PEG	A	216	-	-	1/4/4/4	-
2	PEG	A	218	-	-	2/4/4/4	-
2	PEG	B	222	-	-	2/4/4/4	-
2	PEG	B	220	-	-	3/4/4/4	-
2	PEG	B	221	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	219	PEG	C4-C3-O2-C2
2	A	218	PEG	C4-C3-O2-C2
2	A	218	PEG	O2-C3-C4-O4
2	B	218	PEG	O2-C3-C4-O4
2	B	222	PEG	O1-C1-C2-O2
2	B	221	PEG	O1-C1-C2-O2
2	A	217	PEG	O1-C1-C2-O2
2	B	220	PEG	O2-C3-C4-O4
2	A	219	PEG	O2-C3-C4-O4
2	A	217	PEG	O2-C3-C4-O4
2	A	219	PEG	O1-C1-C2-O2
2	A	217	PEG	C1-C2-O2-C3
2	A	216	PEG	C1-C2-O2-C3
2	B	218	PEG	C1-C2-O2-C3
2	B	219	PEG	C1-C2-O2-C3
2	B	220	PEG	C4-C3-O2-C2
2	B	220	PEG	C1-C2-O2-C3
2	B	221	PEG	C4-C3-O2-C2
2	B	218	PEG	O1-C1-C2-O2
2	B	221	PEG	O2-C3-C4-O4
2	B	222	PEG	O2-C3-C4-O4
2	B	219	PEG	C4-C3-O2-C2
2	B	218	PEG	C4-C3-O2-C2

There are no ring outliers.

5 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	219	PEG	3	0
2	A	217	PEG	5	0
2	A	216	PEG	1	0
2	A	218	PEG	5	0
2	B	221	PEG	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Warning: The R factor obtained from EDS is 0.2311, which does not match the depositor's R factor of 0.181. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/215 (92%)	-0.29	2 (1%) 79 78	6, 16, 30, 52	2 (1%)
1	B	198/215 (92%)	-0.30	4 (2%) 64 63	3, 10, 26, 37	2 (1%)
All	All	396/430 (92%)	-0.29	6 (1%) 71 70	3, 13, 29, 52	4 (1%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	203	VAL	2.7
1	B	204	GLU	2.3
1	B	107	GLU	2.1
1	B	158	ASP	2.1
1	A	159	THR	2.1
1	B	159	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	B	143	7/8	0.96	0.10	3,4,13,14	0
1	CSO	A	143	7/8	0.98	0.09	8,11,17,18	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	B	218	7/7	0.60	0.17	38,39,51,54	0
2	PEG	B	222	7/7	0.65	0.20	61,65,67,68	0
2	PEG	A	217	7/7	0.67	0.21	65,67,72,72	0
2	PEG	A	218	7/7	0.72	0.17	42,50,53,56	0
2	PEG	B	221	7/7	0.73	0.16	27,43,50,57	0
2	PEG	A	216	7/7	0.73	0.14	52,52,56,58	0
2	PEG	B	219	7/7	0.75	0.17	53,54,58,59	0
2	PEG	A	219	7/7	0.79	0.13	41,42,45,50	0
2	PEG	B	220	7/7	0.82	0.12	48,52,54,54	0
3	MG	B	217	1/1	0.97	0.12	24,24,24,24	0
3	MG	B	216	1/1	0.98	0.04	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.