



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2025 – 04:08 PM JST

PDB ID : 9LBG / pdb_00009lbg
Title : The crystal structure of the truncated PAK2 containing K278R mutant
Authors : Hu, H.-F.; Luo, Z.P.; Wu, J.-W.; Wang, Z.-X.
Deposited on : 2025-01-03
Resolution : 2.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2335 (2.90-2.90)
Clashscore	180529	2564 (2.90-2.90)
Ramachandran outliers	177936	2514 (2.90-2.90)
Sidechain outliers	177891	2516 (2.90-2.90)
RSRZ outliers	164620	2337 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 74%, grey 24%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 74% 25% </div> </div>
1	B	466	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 70%, grey 28%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 70% 28% </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5426 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PAK 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	351	Total	C	N	O	S	0	0	0
			2761	1768	454	521	18			
1	B	336	Total	C	N	O	S	0	0	0
			2640	1686	437	500	17			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	59	MET	-	initiating methionine	UNP Q13177
A	60	GLY	-	expression tag	UNP Q13177
A	61	SER	-	expression tag	UNP Q13177
A	62	SER	-	expression tag	UNP Q13177
A	63	HIS	-	expression tag	UNP Q13177
A	64	HIS	-	expression tag	UNP Q13177
A	65	HIS	-	expression tag	UNP Q13177
A	66	HIS	-	expression tag	UNP Q13177
A	67	HIS	-	expression tag	UNP Q13177
A	68	HIS	-	expression tag	UNP Q13177
A	69	SER	-	expression tag	UNP Q13177
A	70	SER	-	expression tag	UNP Q13177
A	71	GLY	-	expression tag	UNP Q13177
A	72	LEU	-	expression tag	UNP Q13177
A	73	VAL	-	expression tag	UNP Q13177
A	74	PRO	-	expression tag	UNP Q13177
A	75	ARG	-	expression tag	UNP Q13177
A	76	GLY	-	expression tag	UNP Q13177
A	77	SER	-	expression tag	UNP Q13177
A	78	HIS	-	expression tag	UNP Q13177
A	79	MET	-	expression tag	UNP Q13177
A	80	ALA	-	expression tag	UNP Q13177
A	81	SER	-	expression tag	UNP Q13177
A	278	ARG	LYS	engineered mutation	UNP Q13177
B	59	MET	-	initiating methionine	UNP Q13177

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	60	GLY	-	expression tag	UNP Q13177
B	61	SER	-	expression tag	UNP Q13177
B	62	SER	-	expression tag	UNP Q13177
B	63	HIS	-	expression tag	UNP Q13177
B	64	HIS	-	expression tag	UNP Q13177
B	65	HIS	-	expression tag	UNP Q13177
B	66	HIS	-	expression tag	UNP Q13177
B	67	HIS	-	expression tag	UNP Q13177
B	68	HIS	-	expression tag	UNP Q13177
B	69	SER	-	expression tag	UNP Q13177
B	70	SER	-	expression tag	UNP Q13177
B	71	GLY	-	expression tag	UNP Q13177
B	72	LEU	-	expression tag	UNP Q13177
B	73	VAL	-	expression tag	UNP Q13177
B	74	PRO	-	expression tag	UNP Q13177
B	75	ARG	-	expression tag	UNP Q13177
B	76	GLY	-	expression tag	UNP Q13177
B	77	SER	-	expression tag	UNP Q13177
B	78	HIS	-	expression tag	UNP Q13177
B	79	MET	-	expression tag	UNP Q13177
B	80	ALA	-	expression tag	UNP Q13177
B	81	SER	-	expression tag	UNP Q13177
B	278	ARG	LYS	engineered mutation	UNP Q13177

- Molecule 2 is D-MALATE (CCD ID: MLT) (formula: C₄H₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			9	4	5		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		
4	B	6	Total	O	0	0
			6	6		

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	166.94Å 166.94Å 208.99Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.14 – 2.89 49.14 – 2.89	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.14-2.89) 98.7 (49.14-2.89)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0415	Depositor
R, R_{free}	0.195 , 0.233 0.196 , 0.228	Depositor DCC
R_{free} test set	1251 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	85.4	Xtriage
Anisotropy	0.546	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 80.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -2/3*h-1/3*k+2/3*l,-1/3*h-2/3*k-2/3*l,2/3*h-2/3*k+1/3*l 0.018 for -h,1/3*h-1/3*k+2/3*l,2/3*h+4/3*k+1/3*l 0.008 for -1/3*h+1/3*k-2/3*l,-k,-4/3*h-2/3*k+1/3*l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5426	wwPDB-VP
Average B, all atoms (Å ²)	104.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLT, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/2810	1.09	0/3800
1	B	0.72	0/2684	1.09	3/3629 (0.1%)
All	All	0.72	0/5494	1.09	3/7429 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	475	ASP	CA-CB-CG	7.29	119.89	112.60
1	B	341	ASP	CA-CB-CG	6.08	118.69	112.60
1	B	133	ASN	CA-CB-CG	5.82	118.42	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2761	0	2820	3	0
1	B	2640	0	2701	3	0
2	B	9	0	4	0	0
3	B	5	0	0	0	0
4	A	5	0	0	0	0
4	B	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5426	0	5525	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:472:ILE:HD11	1:A:498:PRO:HG2	1.79	0.65
1:A:378:MET:SD	1:A:517:LYS:HE3	2.56	0.46
1:B:238:THR:HG23	1:B:239:ILE:HG23	1.99	0.43
1:B:375:LEU:C	1:B:376:LEU:HD12	2.44	0.43
1:B:236:LEU:HD23	1:B:239:ILE:HD11	2.02	0.41
1:A:472:ILE:HD12	1:A:472:ILE:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/466 (74%)	329 (95%)	16 (5%)	0	100	100
1	B	330/466 (71%)	315 (96%)	15 (4%)	0	100	100
All	All	675/932 (72%)	644 (95%)	31 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/404 (76%)	305 (99%)	2 (1%)	81	94
1	B	293/404 (72%)	289 (99%)	4 (1%)	62	86
All	All	600/808 (74%)	594 (99%)	6 (1%)	73	91

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	VAL
1	A	143	THR
1	B	91	VAL
1	B	133	ASN
1	B	326	LEU
1	B	350	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	279	GLN
1	B	133	ASN
1	B	362	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLT	B	601	-	8,8,8	1.13	0	10,10,10	1.42	2 (20%)
3	SO4	B	602	-	4,4,4	0.26	0	6,6,6	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLT	B	601	-	-	4/8/8/8	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	MLT	O1-C1-C2	-2.46	117.74	122.54
2	B	601	MLT	O2-C1-C2	2.32	117.81	112.72

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	MLT	O1-C1-C2-O3
2	B	601	MLT	O2-C1-C2-O3
2	B	601	MLT	O1-C1-C2-C3
2	B	601	MLT	O2-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	351/466 (75%)	-0.31	4 (1%) 77 72	55, 90, 154, 198	0
1	B	336/466 (72%)	-0.17	4 (1%) 76 71	54, 95, 195, 228	0
All	All	687/932 (73%)	-0.24	8 (1%) 76 71	54, 92, 180, 228	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	91	VAL	4.1
1	B	124	LEU	3.8
1	B	395	PRO	2.9
1	A	142	PHE	2.8
1	B	91	VAL	2.3
1	A	460	THR	2.2
1	B	106	LEU	2.1
1	A	93	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MLT	B	601	9/9	0.63	0.16	104,131,145,159	0
3	SO4	B	602	5/5	0.90	0.07	85,113,119,121	5

6.5 Other polymers [i](#)

There are no such residues in this entry.