



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 02:04 AM EDT

PDB ID : 5LNL
Title : Crystal structure of Hsf 1608-1749 putative domain 1
Authors : Thomsen, M.; Wright, J.; Ridley, J.; Goldman, A.
Deposited on : 2016-08-05
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	
1	B	151	
1	C	151	
1	D	151	
1	E	151	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	151	<div><div>%</div><div><div></div><div></div><div></div></div><div>74%11%15%</div></div>
1	G	151	<div><div>%</div><div><div></div><div></div><div></div></div><div>79%8%13%</div></div>
1	H	151	<div><div></div><div><div></div><div></div><div></div></div><div>81%5%14%</div></div>
1	I	151	<div><div></div><div><div></div><div></div><div></div></div><div>73%13%14%</div></div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 8073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hsf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	0	0
			899	551	154	193	1			
1	B	132	Total	C	N	O		0	0	0
			907	556	157	194				
1	C	133	Total	C	N	O		0	0	0
			905	553	157	195				
1	D	129	Total	C	N	O		0	0	1
			883	541	153	189				
1	E	132	Total	C	N	O		0	0	0
			894	548	156	190				
1	F	129	Total	C	N	O		0	0	0
			893	546	152	195				
1	G	132	Total	C	N	O		0	0	0
			911	557	157	197				
1	H	130	Total	C	N	O		0	0	1
			883	542	154	187				
1	I	130	Total	C	N	O		0	0	0
			898	548	154	196				

There are 81 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1607	MET	-	initiating methionine	UNP P71401
A	1750	LEU	-	expression tag	UNP P71401
A	1751	GLU	-	expression tag	UNP P71401
A	1752	HIS	-	expression tag	UNP P71401
A	1753	HIS	-	expression tag	UNP P71401
A	1754	HIS	-	expression tag	UNP P71401
A	1755	HIS	-	expression tag	UNP P71401
A	1756	HIS	-	expression tag	UNP P71401
A	1757	HIS	-	expression tag	UNP P71401
B	1607	MET	-	initiating methionine	UNP P71401
B	1750	LEU	-	expression tag	UNP P71401

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1751	GLU	-	expression tag	UNP P71401
B	1752	HIS	-	expression tag	UNP P71401
B	1753	HIS	-	expression tag	UNP P71401
B	1754	HIS	-	expression tag	UNP P71401
B	1755	HIS	-	expression tag	UNP P71401
B	1756	HIS	-	expression tag	UNP P71401
B	1757	HIS	-	expression tag	UNP P71401
C	1607	MET	-	initiating methionine	UNP P71401
C	1750	LEU	-	expression tag	UNP P71401
C	1751	GLU	-	expression tag	UNP P71401
C	1752	HIS	-	expression tag	UNP P71401
C	1753	HIS	-	expression tag	UNP P71401
C	1754	HIS	-	expression tag	UNP P71401
C	1755	HIS	-	expression tag	UNP P71401
C	1756	HIS	-	expression tag	UNP P71401
C	1757	HIS	-	expression tag	UNP P71401
D	1607	MET	-	initiating methionine	UNP P71401
D	1750	LEU	-	expression tag	UNP P71401
D	1751	GLU	-	expression tag	UNP P71401
D	1752	HIS	-	expression tag	UNP P71401
D	1753	HIS	-	expression tag	UNP P71401
D	1754	HIS	-	expression tag	UNP P71401
D	1755	HIS	-	expression tag	UNP P71401
D	1756	HIS	-	expression tag	UNP P71401
D	1757	HIS	-	expression tag	UNP P71401
E	1607	MET	-	initiating methionine	UNP P71401
E	1750	LEU	-	expression tag	UNP P71401
E	1751	GLU	-	expression tag	UNP P71401
E	1752	HIS	-	expression tag	UNP P71401
E	1753	HIS	-	expression tag	UNP P71401
E	1754	HIS	-	expression tag	UNP P71401
E	1755	HIS	-	expression tag	UNP P71401
E	1756	HIS	-	expression tag	UNP P71401
E	1757	HIS	-	expression tag	UNP P71401
F	1607	MET	-	initiating methionine	UNP P71401
F	1750	LEU	-	expression tag	UNP P71401
F	1751	GLU	-	expression tag	UNP P71401
F	1752	HIS	-	expression tag	UNP P71401
F	1753	HIS	-	expression tag	UNP P71401
F	1754	HIS	-	expression tag	UNP P71401
F	1755	HIS	-	expression tag	UNP P71401
F	1756	HIS	-	expression tag	UNP P71401

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	1757	HIS	-	expression tag	UNP P71401
G	1607	MET	-	initiating methionine	UNP P71401
G	1750	LEU	-	expression tag	UNP P71401
G	1751	GLU	-	expression tag	UNP P71401
G	1752	HIS	-	expression tag	UNP P71401
G	1753	HIS	-	expression tag	UNP P71401
G	1754	HIS	-	expression tag	UNP P71401
G	1755	HIS	-	expression tag	UNP P71401
G	1756	HIS	-	expression tag	UNP P71401
G	1757	HIS	-	expression tag	UNP P71401
H	1607	MET	-	initiating methionine	UNP P71401
H	1750	LEU	-	expression tag	UNP P71401
H	1751	GLU	-	expression tag	UNP P71401
H	1752	HIS	-	expression tag	UNP P71401
H	1753	HIS	-	expression tag	UNP P71401
H	1754	HIS	-	expression tag	UNP P71401
H	1755	HIS	-	expression tag	UNP P71401
H	1756	HIS	-	expression tag	UNP P71401
H	1757	HIS	-	expression tag	UNP P71401
I	1607	MET	-	initiating methionine	UNP P71401
I	1750	LEU	-	expression tag	UNP P71401
I	1751	GLU	-	expression tag	UNP P71401
I	1752	HIS	-	expression tag	UNP P71401
I	1753	HIS	-	expression tag	UNP P71401
I	1754	HIS	-	expression tag	UNP P71401
I	1755	HIS	-	expression tag	UNP P71401
I	1756	HIS	-	expression tag	UNP P71401
I	1757	HIS	-	expression tag	UNP P71401

- Molecule 1: Hsf



HIS	V1607		I1634	K1642	D1643	H1644	N1645	L1648	L1654	G1690	W1691	R1692	G1697	ALA	ASN	GLY	GLN	ASP	D1704	F1705	A1706	T1707	V1708	A1709	T1712	N1713	V1714	D1718	T1723	A1724	T1727	T1735	V1736	K1737	V1738	M1739	D1744	GLY	LEU	LYS	LEU	ASP	LEU	GLU	HIS
-----	-------	--	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-----	-----	-----	-----	-----	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-----	-----	-----	-----	-----	-----	-----	-----


V1607
V1621
K1624
V1632
N1645
R1692
G1697
ALA ASN GLY GLN GLU ASP D1703
G1711
D1718
T1723
A1724
E1725
V1726
T1727
K1728
I1734
T1735
V1736
N1739
D1744
LEU LYS LEU LEU ASP LEU LEU GLI HIS HIS HIS HIS HIS HIS HIS

HIS	V1607		V1619	T1620	V1621		V1624	ASP	V1626		V1632	V1633	I1634		I1641	D1642	D1643	H1644	N1645		L1648		V1692	V1693	K1694		G1697	ALA	ASN	GLY	GLN	ASN	D1703	D1704	F1705	A1706		T1707		G1711		D1718		G1721		A1724		K1741	VAL	ALA	ASP	GLY	LEU	LYS	LEU	ASP	LEU	GLU	HIS	HIS
-----	-------	--	-------	-------	-------	--	-------	-----	-------	--	-------	-------	-------	--	-------	-------	-------	-------	-------	--	-------	--	-------	-------	-------	--	-------	-----	-----	-----	-----	-----	-------	-------	-------	-------	--	-------	--	-------	--	-------	--	-------	--	-------	--	-------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Chain E:  77% 10% 13%





● Molecule 1: Hsf





HIS
HIS
HIS
HIS
HIS


● Molecule 1: Hsf



● Molecule 1: Hsf



● Molecule 1: Hsf



HIS
HIS

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.44Å 50.40Å 256.80Å 90.00° 101.94° 90.00°	Depositor
Resolution (Å)	29.60 – 3.30 29.60 – 3.30	Depositor EDS
% Data completeness (in resolution range)	97.6 (29.60-3.30) 97.6 (29.60-3.30)	Depositor EDS
R_{merge}	0.33	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 3.31Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.296 , 0.334 0.302 , 0.334	Depositor DCC
R_{free} test set	1179 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	56.8	Xtriage
Anisotropy	0.979	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.19 , -32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.085 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	8073	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/904	0.41	0/1229
1	B	0.21	0/913	0.40	0/1243
1	C	0.21	0/911	0.41	0/1242
1	D	0.21	0/888	0.40	0/1206
1	E	0.21	0/899	0.39	0/1224
1	F	0.21	0/898	0.40	0/1224
1	G	0.21	0/917	0.40	0/1249
1	H	0.21	0/888	0.39	0/1209
1	I	0.21	0/903	0.39	0/1231
All	All	0.21	0/8121	0.40	0/11057

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	899	0	838	9	0
1	B	907	0	846	18	0
1	C	905	0	828	9	0
1	D	883	0	819	11	0
1	E	894	0	824	9	0
1	F	893	0	821	11	0
1	G	911	0	845	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	883	0	814	5	0
1	I	898	0	826	12	0
All	All	8073	0	7461	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1645:ASN:HB3	1:B:1692:ARG:HD2	1.74	0.69
1:A:1723:THR:HB	1:A:1739:ASN:HB2	1.76	0.68
1:A:1692:ARG:NH2	1:A:1707:THR:OG1	2.29	0.66
1:D:1692:ARG:NH1	1:D:1707:THR:OG1	2.29	0.65
1:B:1723:THR:HB	1:B:1739:ASN:HB2	1.79	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	125/151 (83%)	122 (98%)	3 (2%)	0	100	100
1	B	128/151 (85%)	125 (98%)	3 (2%)	0	100	100
1	C	129/151 (85%)	125 (97%)	3 (2%)	1 (1%)	19	51
1	D	123/151 (82%)	121 (98%)	2 (2%)	0	100	100
1	E	126/151 (83%)	124 (98%)	2 (2%)	0	100	100
1	F	123/151 (82%)	118 (96%)	5 (4%)	0	100	100
1	G	128/151 (85%)	126 (98%)	2 (2%)	0	100	100
1	H	124/151 (82%)	122 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	124/151 (82%)	122 (98%)	1 (1%)	1 (1%)	19	51
All	All	1130/1359 (83%)	1105 (98%)	23 (2%)	2 (0%)	47	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	1703	ASP
1	C	1624	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/121 (74%)	89 (100%)	0	100	100
1	B	89/121 (74%)	89 (100%)	0	100	100
1	C	87/121 (72%)	87 (100%)	0	100	100
1	D	86/121 (71%)	86 (100%)	0	100	100
1	E	85/121 (70%)	85 (100%)	0	100	100
1	F	89/121 (74%)	89 (100%)	0	100	100
1	G	90/121 (74%)	90 (100%)	0	100	100
1	H	84/121 (69%)	84 (100%)	0	100	100
1	I	90/121 (74%)	90 (100%)	0	100	100
All	All	789/1089 (72%)	789 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	131/151 (86%)	-0.53	0	100	100	29, 55, 95, 107	0
1	B	132/151 (87%)	-0.61	0	100	100	27, 51, 84, 101	0
1	C	133/151 (88%)	-0.60	0	100	100	30, 58, 88, 112	0
1	D	129/151 (85%)	-0.62	0	100	100	32, 58, 79, 105	0
1	E	132/151 (87%)	-0.55	0	100	100	35, 60, 94, 116	0
1	F	129/151 (85%)	-0.61	1 (0%)	86	86	30, 54, 100, 123	0
1	G	132/151 (87%)	-0.56	1 (0%)	86	86	25, 58, 87, 105	0
1	H	130/151 (86%)	-0.64	0	100	100	34, 59, 90, 99	0
1	I	130/151 (86%)	-0.59	0	100	100	34, 57, 83, 91	0
All	All	1178/1359 (86%)	-0.59	2 (0%)	95	96	25, 57, 90, 123	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	1742	VAL	5.0
1	F	1663	THR	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.