



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 01:59 AM EDT

PDB ID : 3LOF  
Title : C-terminal domain of human heat shock 70kDa protein 1B.  
Authors : Osipiuk, J.; Gu, M.; Mihelic, M.; Orton, K.; Morimoto, R.I.; Joachimiak, A.;  
Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2010-02-03  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	113	<div> <div>%</div> <div> <div></div> <div>62%</div> <div>10%</div> <div>.</div> <div>26%</div> </div> </div>
1	B	113	<div> <div>60%</div> <div>7%</div> <div>.</div> <div>29%</div> </div>
1	C	113	<div> <div>%</div> <div> <div></div> <div>57%</div> <div>12%</div> <div>.</div> <div>29%</div> </div> </div>
1	D	113	<div> <div>%</div> <div> <div></div> <div>59%</div> <div>6%</div> <div>.</div> <div>30%</div> </div> </div>
1	E	113	<div> <div>58%</div> <div>11%</div> <div>.</div> <div>27%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	113	<div><div></div><div>4%</div><div>59%</div><div>9%</div><div>29%</div></div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock 70 kDa protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	84	Total	C	N	O	S	Se	0	1	0
			659	410	112	134	2	1			
1	B	80	Total	C	N	O	S	Se	0	0	0
			622	386	106	127	2	1			
1	C	80	Total	C	N	O	S	Se	0	1	0
			628	391	107	127	2	1			
1	D	79	Total	C	N	O	S	Se	0	1	0
			623	388	106	126	2	1			
1	E	82	Total	C	N	O	S	Se	0	1	0
			641	398	109	131	2	1			
1	F	80	Total	C	N	O	S	Se	0	0	0
			627	390	107	127	2	1			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	529	SER	-	expression tag	UNP P08107
A	530	ASN	-	expression tag	UNP P08107
A	531	ALA	-	expression tag	UNP P08107
A	532	ALA	-	expression tag	UNP P08107
A	533	ALA	-	expression tag	UNP P08107
B	529	SER	-	expression tag	UNP P08107
B	530	ASN	-	expression tag	UNP P08107
B	531	ALA	-	expression tag	UNP P08107
B	532	ALA	-	expression tag	UNP P08107
B	533	ALA	-	expression tag	UNP P08107
C	529	SER	-	expression tag	UNP P08107
C	530	ASN	-	expression tag	UNP P08107
C	531	ALA	-	expression tag	UNP P08107
C	532	ALA	-	expression tag	UNP P08107
C	533	ALA	-	expression tag	UNP P08107
D	529	SER	-	expression tag	UNP P08107
D	530	ASN	-	expression tag	UNP P08107

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	531	ALA	-	expression tag	UNP P08107
D	532	ALA	-	expression tag	UNP P08107
D	533	ALA	-	expression tag	UNP P08107
E	529	SER	-	expression tag	UNP P08107
E	530	ASN	-	expression tag	UNP P08107
E	531	ALA	-	expression tag	UNP P08107
E	532	ALA	-	expression tag	UNP P08107
E	533	ALA	-	expression tag	UNP P08107
F	529	SER	-	expression tag	UNP P08107
F	530	ASN	-	expression tag	UNP P08107
F	531	ALA	-	expression tag	UNP P08107
F	532	ALA	-	expression tag	UNP P08107
F	533	ALA	-	expression tag	UNP P08107

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	28	Total O 28 28	0	0
2	B	14	Total O 14 14	0	0
2	C	26	Total O 26 26	0	0
2	D	31	Total O 31 31	0	0
2	E	24	Total O 24 24	0	0
2	F	11	Total O 11 11	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heat shock 70 kDa protein 1




- Molecule 1: Heat shock 70 kDa protein 1




- Molecule 1: Heat shock 70 kDa protein 1




- Molecule 1: Heat shock 70 kDa protein 1



- Molecule 1: Heat shock 70 kDa protein 1



● Molecule 1: Heat shock 70 kDa protein 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.70Å 71.88Å 143.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.00 – 2.40 32.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.0 (32.00-2.40) 99.0 (32.00-2.40)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.51 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.192 , 0.235 0.188 , 0.233	Depositor DCC
$R_{free}$ test set	1475 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.2	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3934	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.97	0/668	0.80	0/892
1	B	0.97	2/627 (0.3%)	0.81	0/838
1	C	1.05	0/636	0.82	0/849
1	D	0.95	0/631	0.90	3/842 (0.4%)
1	E	0.93	0/649	0.81	0/866
1	F	0.93	0/632	0.84	1/844 (0.1%)
All	All	0.97	2/3843 (0.1%)	0.83	4/5131 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	612	GLN	CB-CG	-5.05	1.39	1.52
1	B	576	GLU	CG-CD	5.01	1.59	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	582	ASP	CB-CG-OD2	-5.76	113.12	118.30
1	D	586	LEU	CA-CB-CG	-5.72	102.15	115.30
1	F	596	ARG	NE-CZ-NH2	5.32	122.96	120.30
1	D	582	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	659	0	657	10	0
1	B	622	0	610	3	0
1	C	628	0	623	9	0
1	D	623	0	618	6	0
1	E	641	0	632	10	0
1	F	627	0	620	5	0
2	A	28	0	0	2	0
2	B	14	0	0	0	0
2	C	26	0	0	3	0
2	D	31	0	0	2	0
2	E	24	0	0	1	0
2	F	11	0	0	1	0
All	All	3934	0	3760	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:585:THR:CG2	2:C:33:HOH:O	2.22	0.87
1:C:553:VAL:HG11	1:C:571:LEU:HD13	1.70	0.71
1:A:553:VAL:HG21	1:A:571:LEU:HD13	1.72	0.68
1:B:553:VAL:HG11	1:B:571:LEU:HD13	1.75	0.68
1:D:585:THR:CG2	2:D:89:HOH:O	2.43	0.67
1:C:585:THR:HG21	2:C:33:HOH:O	1.93	0.64
2:A:14:HOH:O	1:C:585:THR:HG23	2.02	0.59
1:A:576:GLU:HG3	1:A:577:VAL:N	2.19	0.57
1:A:553:VAL:CG2	1:A:571:LEU:HD13	2.36	0.56
1:A:553:VAL:CG2	1:A:571:LEU:CD1	2.85	0.55
1:B:542:LEU:HD11	1:B:577:VAL:HG11	1.90	0.54
1:E:595:LYS:NZ	2:E:110:HOH:O	2.41	0.54
1:F:555:ASP:OD2	1:F:555:ASP:C	2.46	0.53
1:A:566:ASP:HB3	1:A:606:ILE:HD12	1.92	0.51
1:E:563:SER:OG	1:E:566:ASP:OD2	2.27	0.51
1:F:610:LEU:HD23	1:F:610:LEU:N	2.29	0.48
1:E:539:LYS:HG3	1:E:592:PHE:CZ	2.47	0.48
1:E:573:LYS:HG3	1:E:599:LEU:HD12	1.94	0.48
1:F:604:ASN:ND2	2:F:128:HOH:O	2.39	0.47
1:D:585:THR:HG23	2:D:90:HOH:O	2.15	0.46
1:C:582:ASP:O	2:C:33:HOH:O	2.21	0.46
1:B:585:THR:HG22	1:B:586:LEU:HD22	1.97	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:608:SER:OG	1:E:610:LEU:HD22	2.16	0.45
1:D:593:GLU:O	1:D:597:LYS:HD3	2.16	0.45
1:A:612:GLN:NE2	2:A:75:HOH:O	2.50	0.45
1:D:553:VAL:HG21	1:D:571:LEU:HD13	1.99	0.45
1:A:585:THR:HG22	1:A:586:LEU:HD22	1.98	0.44
1:D:584:ASN:HB3	1:D:587:ALA:HB2	2.00	0.43
1:E:608:SER:CB	1:E:610:LEU:HD22	2.48	0.43
1:F:550:LYS:NZ	1:F:575:GLN:OE1	2.40	0.43
1:C:545:TYR:CD2	1:C:599:LEU:HD23	2.53	0.43
1:A:582:ASP:HA	1:C:586:LEU:HD23	2.01	0.43
1:F:553:VAL:HG11	1:F:571:LEU:HD13	2.00	0.42
1:A:530:ASN:OD1	1:A:530:ASN:C	2.58	0.42
1:E:545:TYR:OH	1:E:604:ASN:ND2	2.53	0.42
1:E:550:LYS:HG3	1:E:571:LEU:HD12	2.01	0.42
1:E:553:VAL:HG11	1:E:571:LEU:HD13	2.01	0.42
1:D:604:ASN:HD22	1:D:604:ASN:HA	1.71	0.41
1:E:573:LYS:HG3	1:E:599:LEU:CD1	2.50	0.41
1:A:549:MSE:HE3	1:A:570:VAL:HG11	2.03	0.40
1:C:545:TYR:HD2	1:C:599:LEU:HD23	1.85	0.40
1:C:550:LYS:O	1:C:553:VAL:HG12	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	83/113 (74%)	82 (99%)	1 (1%)	0	100	100
1	B	76/113 (67%)	75 (99%)	1 (1%)	0	100	100
1	C	77/113 (68%)	74 (96%)	3 (4%)	0	100	100
1	D	76/113 (67%)	76 (100%)	0	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	79/113 (70%)	76 (96%)	3 (4%)	0	100	100
1	F	76/113 (67%)	71 (93%)	5 (7%)	0	100	100
All	All	467/678 (69%)	454 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	71/85 (84%)	67 (94%)	4 (6%)	17	30
1	B	66/85 (78%)	58 (88%)	8 (12%)	4	5
1	C	67/85 (79%)	56 (84%)	11 (16%)	2	2
1	D	67/85 (79%)	60 (90%)	7 (10%)	5	8
1	E	68/85 (80%)	60 (88%)	8 (12%)	4	6
1	F	67/85 (79%)	59 (88%)	8 (12%)	4	6
All	All	406/510 (80%)	360 (89%)	46 (11%)	4	7

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	530	ASN
1	A	571	LEU
1	A	576	GLU
1	A	594	HIS
1	B	550	LYS
1	B	551	SER
1	B	562	ILE
1	B	568	LYS
1	B	571	LEU
1	B	576	GLU
1	B	586	LEU
1	B	612	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	530	ASN
1	C	544	SER
1	C	553	VAL
1	C	554	GLU
1	C	561	LYS
1	C	562	ILE
1	C	563	SER
1	C	568	LYS
1	C	569	LYS
1	C	571	LEU
1	C	585	THR
1	D	550	LYS
1	D	563	SER
1	D	571	LEU
1	D	585	THR
1	D	586	LEU
1	D	597	LYS
1	D	604	ASN
1	E	550	LYS
1	E	553	VAL
1	E	563	SER
1	E	569	LYS
1	E	571	LEU
1	E	576	GLU
1	E	586	LEU
1	E	610	LEU
1	F	553	VAL
1	F	554	GLU
1	F	555	ASP
1	F	563	SER
1	F	571	LEU
1	F	589	LYS
1	F	601	GLN
1	F	608	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	540	ASN
1	A	601	GLN
1	A	612	GLN
1	B	584	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	612	GLN
1	C	540	ASN
1	C	604	ASN
1	C	612	GLN
1	D	540	ASN
1	D	601	GLN
1	D	604	ASN
1	E	604	ASN
1	F	604	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	83/113 (73%)	-0.24	1 (1%) 76 73	12, 28, 46, 50	1 (1%)
1	B	79/113 (69%)	-0.26	0 100 100	15, 31, 54, 62	0
1	C	79/113 (69%)	-0.34	1 (1%) 74 71	10, 25, 45, 53	1 (1%)
1	D	78/113 (69%)	-0.64	1 (1%) 74 71	14, 22, 50, 58	1 (1%)
1	E	81/113 (71%)	-0.30	0 100 100	13, 27, 46, 57	1 (1%)
1	F	79/113 (69%)	0.47	5 (6%) 27 25	12, 29, 47, 50	0
All	All	479/678 (70%)	-0.22	8 (1%) 69 65	10, 27, 49, 62	4 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	560	GLY	3.4
1	F	613	GLY	3.2
1	F	565	ALA	2.6
1	A	560	GLY	2.6
1	F	611	TYR	2.5
1	F	614	ALA	2.5
1	F	532	ALA	2.3
1	C	562	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.