



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2024 – 04:56 AM EDT

PDB ID : 1LZ4
Title : ENTHALPIC DESTABILIZATION OF A MUTANT HUMAN LYSOZYME
LACKING A DISULFIDE BRIDGE BETWEEN CYSTEINE-77 AND
CYSTEINE-95
Authors : Inaka, K.; Matsushima, M.; Kuroki, R.; Taniyama, Y.; Kikuchi, M.
Deposited on : 1993-02-03
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	130	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	130	Total	C	N	O	S	0	0	0
			1016	627	194	186	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	77	ALA	CYS	conflict	UNP P61626

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	57	Total	O	0	0
			57	57		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HUMAN LYSOZYME

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.82Å 60.86Å 33.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (5.00-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.188 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1073	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	0/1036	1.99	32/1401 (2.3%)

There are no bond length outliers.

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	50	ARG	NE-CZ-NH1	15.68	128.14	120.30
1	A	98	ARG	NE-CZ-NH2	-14.95	112.83	120.30
1	A	21	ARG	NE-CZ-NH2	-13.58	113.51	120.30
1	A	98	ARG	NE-CZ-NH1	11.04	125.82	120.30
1	A	50	ARG	CD-NE-CZ	10.56	138.39	123.60
1	A	115	ARG	NE-CZ-NH2	-10.41	115.09	120.30
1	A	53	ASP	CB-CG-OD2	-9.67	109.59	118.30
1	A	113	ARG	NE-CZ-NH2	-9.51	115.55	120.30
1	A	107	ARG	NE-CZ-NH1	9.01	124.80	120.30
1	A	10	ARG	NE-CZ-NH1	-8.91	115.84	120.30
1	A	18	ASP	CB-CG-OD1	8.68	126.11	118.30
1	A	62	ARG	NE-CZ-NH2	7.38	123.99	120.30
1	A	7	GLU	CG-CD-OE2	-7.33	103.65	118.30
1	A	62	ARG	NE-CZ-NH1	-7.32	116.64	120.30
1	A	50	ARG	NE-CZ-NH2	-7.04	116.78	120.30
1	A	107	ARG	NE-CZ-NH2	-7.01	116.80	120.30
1	A	119	ARG	NE-CZ-NH1	-6.94	116.83	120.30
1	A	4	GLU	OE1-CD-OE2	6.87	131.55	123.30
1	A	119	ARG	NE-CZ-NH2	6.86	123.73	120.30
1	A	87	ASP	CB-CG-OD1	6.76	124.39	118.30
1	A	21	ARG	NE-CZ-NH1	6.65	123.63	120.30
1	A	102	ASP	CB-CG-OD1	6.53	124.18	118.30
1	A	115	ARG	NE-CZ-NH1	6.40	123.50	120.30
1	A	7	GLU	CG-CD-OE1	6.25	130.79	118.30
1	A	120	ASP	CB-CG-OD2	-6.21	112.71	118.30
1	A	38	TYR	CB-CG-CD1	5.96	124.58	121.00
1	A	101	ARG	NE-CZ-NH1	5.91	123.26	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	TYR	CB-CG-CD2	5.40	124.24	121.00
1	A	18	ASP	CB-CG-OD2	-5.34	113.49	118.30
1	A	64	TRP	N-CA-CB	-5.12	101.39	110.60
1	A	102	ASP	CB-CG-OD2	-5.11	113.70	118.30
1	A	73	ALA	N-CA-CB	-5.10	102.96	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1016	0	972	10	0
2	A	57	0	0	1	0
All	All	1073	0	972	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:ARG:NH1	1:A:71:PRO:O	2.25	0.69
1:A:75:ASN:ND2	1:A:78:HIS:HA	2.12	0.65
1:A:64:TRP:O	1:A:77:ALA:HB3	1.96	0.65
1:A:64:TRP:CE3	1:A:76:ALA:HB3	2.48	0.49
1:A:1:LYS:HE2	1:A:3:PHE:CZ	2.49	0.48
1:A:75:ASN:CG	1:A:78:HIS:HA	2.36	0.46
1:A:112:TRP:CZ3	1:A:117:GLN:HB2	2.51	0.45
1:A:113:ARG:HA	1:A:117:GLN:HE21	1.83	0.44
1:A:74:VAL:N	2:A:186:HOH:O	2.50	0.44
1:A:104:GLN:O	1:A:107:ARG:HB2	2.18	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/130 (98%)	125 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/104 (98%)	99 (97%)	3 (3%)	37	26

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	TYR
1	A	123	GLN
1	A	126	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.