



wwPDB NMR Structure Validation Summary Report ⓘ

Oct 30, 2024 – 07:35 AM EDT

PDB ID : 2LZS
BMRB ID : 18771
Title : TatA oligomer
Authors : Rodriguez, F.M.; Berks, B.C.; Schnell, J.R.
Deposited on : 2012-10-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 4%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	55	64% 20% 5% 11%
1	B	55	82% 7% 11%
1	C	55	69% 20% 11%
1	D	55	65% 22% • 11%
1	E	55	73% 15% • 11%
1	F	55	73% 15% • 11%
1	G	55	67% 16% 5% 11%
1	H	55	71% 18% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	55	 A horizontal bar chart showing the quality of chain 1. The bar is divided into four segments: green (62%), yellow (22%), orange (5%), and grey (11%).

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 6804 atoms, of which 3555 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Sec-independent protein translocase protein TatA.

Mol	Chain	Residues	Atoms						Trace
1	A	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	B	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	C	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	D	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	E	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	F	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	G	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	H	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	
1	I	49	Total	C	H	N	O	S	0
			756	238	395	57	64	2	

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	HIS	-	expression tag	UNP P69428
A	51	HIS	-	expression tag	UNP P69428
A	52	HIS	-	expression tag	UNP P69428
A	53	HIS	-	expression tag	UNP P69428
A	54	HIS	-	expression tag	UNP P69428
A	55	HIS	-	expression tag	UNP P69428
B	50	HIS	-	expression tag	UNP P69428
B	51	HIS	-	expression tag	UNP P69428
B	52	HIS	-	expression tag	UNP P69428
B	53	HIS	-	expression tag	UNP P69428
B	54	HIS	-	expression tag	UNP P69428
B	55	HIS	-	expression tag	UNP P69428
C	50	HIS	-	expression tag	UNP P69428
C	51	HIS	-	expression tag	UNP P69428
C	52	HIS	-	expression tag	UNP P69428

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	53	HIS	-	expression tag	UNP P69428
C	54	HIS	-	expression tag	UNP P69428
C	55	HIS	-	expression tag	UNP P69428
D	50	HIS	-	expression tag	UNP P69428
D	51	HIS	-	expression tag	UNP P69428
D	52	HIS	-	expression tag	UNP P69428
D	53	HIS	-	expression tag	UNP P69428
D	54	HIS	-	expression tag	UNP P69428
D	55	HIS	-	expression tag	UNP P69428
E	50	HIS	-	expression tag	UNP P69428
E	51	HIS	-	expression tag	UNP P69428
E	52	HIS	-	expression tag	UNP P69428
E	53	HIS	-	expression tag	UNP P69428
E	54	HIS	-	expression tag	UNP P69428
E	55	HIS	-	expression tag	UNP P69428
F	50	HIS	-	expression tag	UNP P69428
F	51	HIS	-	expression tag	UNP P69428
F	52	HIS	-	expression tag	UNP P69428
F	53	HIS	-	expression tag	UNP P69428
F	54	HIS	-	expression tag	UNP P69428
F	55	HIS	-	expression tag	UNP P69428
G	50	HIS	-	expression tag	UNP P69428
G	51	HIS	-	expression tag	UNP P69428
G	52	HIS	-	expression tag	UNP P69428
G	53	HIS	-	expression tag	UNP P69428
G	54	HIS	-	expression tag	UNP P69428
G	55	HIS	-	expression tag	UNP P69428
H	50	HIS	-	expression tag	UNP P69428
H	51	HIS	-	expression tag	UNP P69428
H	52	HIS	-	expression tag	UNP P69428
H	53	HIS	-	expression tag	UNP P69428
H	54	HIS	-	expression tag	UNP P69428
H	55	HIS	-	expression tag	UNP P69428
I	50	HIS	-	expression tag	UNP P69428
I	51	HIS	-	expression tag	UNP P69428
I	52	HIS	-	expression tag	UNP P69428
I	53	HIS	-	expression tag	UNP P69428
I	54	HIS	-	expression tag	UNP P69428
I	55	HIS	-	expression tag	UNP P69428

4 Residue-property plots [i](#)


These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Sec-independent protein translocase protein TatA

Chain A: 



- Molecule 1: Sec-independent protein translocase protein TatA

Chain B: 



- Molecule 1: Sec-independent protein translocase protein TatA

Chain C: 




- Molecule 1: Sec-independent protein translocase protein TatA

Chain D: 



- Molecule 1: Sec-independent protein translocase protein TatA

Chain E: 



- Molecule 1: Sec-independent protein translocase protein TatA

Chain F: 



- Molecule 1: Sec-independent protein translocase protein TatA



- Molecule 1: Sec-independent protein translocase protein TatA



- Molecule 1: Sec-independent protein translocase protein TatA



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 10 calculated structures, 1 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	212
Number of shifts mapped to atoms	212
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	4%

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FME

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	361	395	396	11
1	B	361	395	396	2
1	C	361	395	396	4
1	D	361	395	396	7
1	E	361	395	396	6
1	F	361	395	396	6
1	G	361	395	396	8
1	H	361	395	396	5
1	I	361	395	396	16
All	All	3249	3555	3564	63

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

5 of 63 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:I:4:ILE:HD13	1:I:4:ILE:H	0.68	1.46
1:I:8:GLN:NE2	1:I:8:GLN:N	0.67	2.43
1:D:8:GLN:N	1:D:8:GLN:OE1	0.65	2.30
1:B:1:FME:SD	1:B:1:FME:N	0.59	2.75
1:F:1:FME:SD	1:F:1:FME:N	0.59	2.76

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	47/55 (85%)	43 (91%)	4 (9%)	0 (0%)	100	100
1	B	47/55 (85%)	41 (87%)	6 (13%)	0 (0%)	100	100
1	C	47/55 (85%)	43 (91%)	3 (6%)	1 (2%)	8	48
1	D	47/55 (85%)	40 (85%)	4 (9%)	3 (6%)	2	18
1	E	47/55 (85%)	44 (94%)	3 (6%)	0 (0%)	100	100
1	F	47/55 (85%)	42 (89%)	5 (11%)	0 (0%)	100	100
1	G	47/55 (85%)	44 (94%)	3 (6%)	0 (0%)	100	100
1	H	47/55 (85%)	43 (91%)	3 (6%)	1 (2%)	8	48
1	I	47/55 (85%)	43 (91%)	4 (9%)	0 (0%)	100	100
All	All	423/495 (85%)	383 (91%)	35 (8%)	5 (1%)	14	62

All 5 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	C	4	ILE
1	D	4	ILE
1	D	44	SER
1	D	47	GLU
1	H	4	ILE

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	38/44 (86%)	36 (95%)	2 (5%)	21	74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	38/44 (86%)	37 (97%)	1 (3%)	42	89
1	C	38/44 (86%)	37 (97%)	1 (3%)	42	89
1	D	38/44 (86%)	38 (100%)	0 (0%)	100	100
1	E	38/44 (86%)	37 (97%)	1 (3%)	42	89
1	F	38/44 (86%)	38 (100%)	0 (0%)	100	100
1	G	38/44 (86%)	36 (95%)	2 (5%)	21	74
1	H	38/44 (86%)	38 (100%)	0 (0%)	100	100
1	I	38/44 (86%)	34 (89%)	4 (11%)	8	53
All	All	342/396 (86%)	331 (97%)	11 (3%)	36	86

5 of 11 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	5	SER
1	A	43	MET
1	B	22	THR
1	C	22	THR
1	E	22	THR

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	FME	I	1	1	8,9,10	0.86	1 (12%)

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	FME	D	1	1	8,9,10	0.82	0 (0%)
1	FME	H	1	1	8,9,10	0.86	1 (12%)
1	FME	C	1	1	8,9,10	0.86	1 (12%)
1	FME	B	1	1	8,9,10	0.85	0 (0%)
1	FME	G	1	1	8,9,10	1.06	1 (12%)
1	FME	A	1	1	8,9,10	0.95	1 (12%)
1	FME	F	1	1	8,9,10	0.99	1 (12%)
1	FME	E	1	1	8,9,10	0.90	1 (12%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	FME	I	1	1	8,9,11	1.15	0 (0%)
1	FME	D	1	1	8,9,11	1.14	0 (0%)
1	FME	H	1	1	8,9,11	1.15	0 (0%)
1	FME	C	1	1	8,9,11	1.14	0 (0%)
1	FME	B	1	1	8,9,11	1.14	0 (0%)
1	FME	G	1	1	8,9,11	1.04	0 (0%)
1	FME	A	1	1	8,9,11	1.12	0 (0%)
1	FME	F	1	1	8,9,11	1.02	0 (0%)
1	FME	E	1	1	8,9,11	1.13	0 (0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FME	I	1	1	-	0,7,9,11	-
1	FME	D	1	1	-	0,7,9,11	-
1	FME	H	1	1	-	1,7,9,11	-
1	FME	C	1	1	-	0,7,9,11	-
1	FME	B	1	1	-	0,7,9,11	-
1	FME	G	1	1	-	0,7,9,11	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FME	A	1	1	-	0,7,9,11	-
1	FME	F	1	1	-	0,7,9,11	-
1	FME	E	1	1	-	0,7,9,11	-

5 of 7 bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	1	FME	CA-N	2.68	1.50	1.46
1	F	1	FME	CA-N	2.50	1.50	1.46
1	A	1	FME	CA-N	2.17	1.49	1.46
1	E	1	FME	CA-N	2.09	1.49	1.46
1	I	1	FME	CA-N	2.05	1.49	1.46

There are no bond-angle outliers.

There are no chirality outliers.

All torsion outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	H	1	FME	O1-CN-N-CA

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 4% for the well-defined parts and 4% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	212
Number of shifts mapped to atoms	212
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	None (insufficient data)
$^{13}\text{C}_\beta$	3	—	None (insufficient data)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	44	-0.06 ± 0.21	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 4%, i.e. 208 atoms were assigned a chemical shift out of a possible 5796. 0 out of 81 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	88/2205 (4%)	44/918 (5%)	0/864 (0%)	44/423 (10%)
Sidechain	120/3303 (4%)	90/2196 (4%)	30/1044 (3%)	0/63 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	0/288 (0%)	0/144 (0%)	0/135 (0%)	0/9 (0%)
Overall	208/5796 (4%)	134/3258 (4%)	30/2043 (1%)	44/495 (9%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:

