



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 12, 2024 – 03:15 PM EST

PDB ID : 3M5J  
Title : Crystal structure of a H7 influenza virus hemagglutinin complexed with LSTb  
Authors : Yang, H.; Chen, L.M.; Carney, P.J.; Donis, R.O.; Stevens, J.  
Deposited on : 2010-03-12  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	<div> <div>%</div> <div>91% 6% .</div> </div>
1	C	317	<div> <div>90% 8% .</div> </div>
1	E	317	<div> <div>%</div> <div>92% 6% .</div> </div>
2	B	182	<div> <div>%</div> <div>92% 5% .</div> </div>
2	D	182	<div> <div>%</div> <div>90% . . 5%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	182	 90% 6%
3	G	4	 50% 50%
4	H	2	 50% 50%
4	I	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	G	2	X	-	-	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2392	1488	433	458	13			
1	C	309	Total	C	N	O	S	0	0	0
			2383	1482	431	457	13			
1	E	309	Total	C	N	O	S	0	0	0
			2383	1482	431	457	13			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ALA	-	expression tag	UNP B7NY59
A	8	ASP	-	expression tag	UNP B7NY59
A	9	PRO	-	expression tag	UNP B7NY59
A	216	ASN	SER	SEE REMARK 999	UNP B7NY59
A	257	THR	ALA	SEE REMARK 999	UNP B7NY59
A	309	VAL	ILE	SEE REMARK 999	UNP B7NY59
C	7	ALA	-	expression tag	UNP B7NY59
C	8	ASP	-	expression tag	UNP B7NY59
C	9	PRO	-	expression tag	UNP B7NY59
C	216	ASN	SER	SEE REMARK 999	UNP B7NY59
C	257	THR	ALA	SEE REMARK 999	UNP B7NY59
C	309	VAL	ILE	SEE REMARK 999	UNP B7NY59
E	7	ALA	-	expression tag	UNP B7NY59
E	8	ASP	-	expression tag	UNP B7NY59
E	9	PRO	-	expression tag	UNP B7NY59
E	216	ASN	SER	SEE REMARK 999	UNP B7NY59
E	257	THR	ALA	SEE REMARK 999	UNP B7NY59
E	309	VAL	ILE	SEE REMARK 999	UNP B7NY59


- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	176	Total	C	N	O	S	0	0	0
			1419	876	248	288	7			
2	D	172	Total	C	N	O	S	0	0	0
			1393	861	244	281	7			
2	F	171	Total	C	N	O	S	0	0	0
			1384	856	242	279	7			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	176	GLY	VAL	SEE REMARK 999	UNP B7NYS1
B	177	ARG	LYS	SEE REMARK 999	UNP B7NYS1
B	179	VAL	-	expression tag	UNP B7NYS1
B	180	PRO	-	expression tag	UNP B7NYS1
B	181	ARG	-	expression tag	UNP B7NYS1
B	182	GLY	-	expression tag	UNP B7NYS1
D	176	GLY	VAL	SEE REMARK 999	UNP B7NYS1
D	177	ARG	LYS	SEE REMARK 999	UNP B7NYS1
D	179	VAL	-	expression tag	UNP B7NYS1
D	180	PRO	-	expression tag	UNP B7NYS1
D	181	ARG	-	expression tag	UNP B7NYS1
D	182	GLY	-	expression tag	UNP B7NYS1
F	176	GLY	VAL	SEE REMARK 999	UNP B7NYS1
F	177	ARG	LYS	SEE REMARK 999	UNP B7NYS1
F	179	VAL	-	expression tag	UNP B7NYS1
F	180	PRO	-	expression tag	UNP B7NYS1
F	181	ARG	-	expression tag	UNP B7NYS1
F	182	GLY	-	expression tag	UNP B7NYS1

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-3)-[N-acetyl-alpha-neuraminic acid-(2-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	4	Total	C	N	O	0	0	0
			57	31	2	24			

- Molecule 4 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	2	Total	C	N	O	0	0	0
			33	19	2	12			
4	I	2	Total	C	N	O	0	0	0
			33	19	2	12			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	D	1	Total	C	O	0	0
			6	3	3		
6	F	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	100	Total	O	0	0
			100	100		
7	B	74	Total	O	0	0
			74	74		
7	C	90	Total	O	0	0
			90	90		
7	D	49	Total	O	0	0
			49	49		
7	E	134	Total	O	0	0
			134	134		
7	F	67	Total	O	0	0
			67	67		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin




- Molecule 1: Hemagglutinin




- Molecule 1: Hemagglutinin




- Molecule 2: Hemagglutinin





- Molecule 2: Hemagglutinin







• Molecule 2: Hemagglutinin



• Molecule 3: beta-D-galactopyranose-(1-3)-[N-acetyl-alpha-neuraminic acid-(2-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose



• Molecule 4: N-acetyl-alpha-neuraminic acid-(2-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: N-acetyl-alpha-neuraminic acid-(2-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.08Å 116.52Å 251.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.60 50.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	91.4 (50.00-2.60) 91.4 (50.00-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.97 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.204 , 0.247 0.201 , 0.243	Depositor DCC
$R_{free}$ test set	2842 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.6	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 19.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12105	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GOL, GAL, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2440	0.48	0/3297
1	C	0.34	0/2431	0.48	0/3286
1	E	0.36	0/2431	0.52	2/3286 (0.1%)
2	B	0.33	0/1443	0.44	0/1945
2	D	0.33	0/1417	0.46	0/1910
2	F	0.33	0/1408	0.46	0/1898
All	All	0.34	0/11570	0.48	2/15622 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	93	GLY	O-C-N	-6.34	112.56	122.70
1	E	93	GLY	C-N-CA	5.04	134.31	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2392	0	2353	9	0
1	C	2383	0	2340	11	0
1	E	2383	0	2339	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1419	0	1324	5	0
2	D	1393	0	1301	7	0
2	F	1384	0	1293	3	0
3	G	57	0	46	0	0
4	H	33	0	24	0	0
4	I	33	0	24	0	0
5	A	14	0	13	0	0
5	B	14	0	13	0	0
5	C	14	0	13	0	0
5	D	14	0	13	0	0
5	E	14	0	13	0	0
5	F	14	0	13	0	0
6	B	18	0	24	0	0
6	D	6	0	8	1	0
6	F	6	0	8	0	0
7	A	100	0	0	0	0
7	B	74	0	0	0	0
7	C	90	0	0	0	0
7	D	49	0	0	0	0
7	E	134	0	0	0	0
7	F	67	0	0	0	0
All	All	12105	0	11162	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:124:LYS:HD3	2:D:134:GLY:HA2	1.81	0.61
1:C:134:GLY:HA3	1:C:153:TRP:HB3	1.85	0.59
1:E:50:LYS:HG2	1:E:275:ASP:HB2	1.88	0.55
1:C:211:GLN:HG2	1:C:213:PHE:CZ	2.43	0.53
2:F:19:ASN:HB2	2:F:36:ALA:HB2	1.92	0.51
1:A:113:ARG:HG2	1:A:267:GLY:HA3	1.93	0.51
1:C:237:LEU:HD13	1:C:243:VAL:HG23	1.93	0.51
1:C:13:ILE:HG22	2:D:140:ILE:HD11	1.92	0.50
1:C:116:GLY:HA2	1:C:265:SER:HB3	1.94	0.50
1:C:237:LEU:HD22	1:C:243:VAL:HG23	1.94	0.50
1:A:132:THR:HG23	1:A:152:LYS:HD2	1.93	0.49
1:E:132:THR:HG22	1:E:152:LYS:HD2	1.94	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:LEU:HD11	1:A:112:LEU:HD11	1.96	0.47
1:C:180:TRP:CE2	1:C:204:VAL:HG21	2.50	0.47
1:A:186:GLU:HB2	1:A:220:ARG:HG3	1.96	0.47
2:B:77:ILE:HD11	2:D:80:VAL:HG21	1.96	0.47
1:C:70:LEU:HD11	1:C:112:LEU:HD11	1.97	0.46
1:A:15:LEU:HD22	2:B:119:TYR:HA	1.98	0.45
1:E:174:LYS:HB2	1:E:175:PRO:HD2	1.97	0.45
2:D:54:ARG:HH21	6:D:184:GOL:H11	1.83	0.44
1:A:180:TRP:CE2	1:A:204:VAL:HG21	2.53	0.44
1:E:70:LEU:HD11	1:E:112:LEU:HD11	2.00	0.44
2:D:19:ASN:OD1	2:D:19:ASN:N	2.45	0.44
2:D:131:GLU:OE2	2:D:170:ARG:HD2	2.18	0.43
2:B:106:HIS:O	2:B:110:LEU:HB2	2.18	0.43
2:F:131:GLU:OE2	2:F:170:ARG:HD2	2.17	0.43
2:B:44:ALA:O	2:B:48:ILE:HG12	2.18	0.43
1:E:180:TRP:CE2	1:E:204:VAL:HG21	2.55	0.42
1:A:203:THR:HB	1:A:246:THR:OG1	2.20	0.42
2:D:140:ILE:HG22	2:D:142:HIS:H	1.84	0.42
1:E:203:THR:OG1	1:E:246:THR:HB	2.19	0.41
1:C:158(B):ASN:HA	1:C:192:THR:O	2.21	0.41
1:C:71:ILE:HG13	1:C:73:PRO:HD3	2.03	0.41
1:A:118:ILE:HD12	1:A:120:LYS:HZ2	1.86	0.41
1:C:107:SER:HB3	2:F:75:GLN:HB2	2.03	0.40
1:A:201:LEU:HD11	1:A:212:SER:HB3	2.03	0.40
1:E:203:THR:HG22	1:E:212:SER:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	308/317 (97%)	296 (96%)	11 (4%)	1 (0%)	37 59

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	307/317 (97%)	298 (97%)	8 (3%)	1 (0%)	37	59
1	E	307/317 (97%)	300 (98%)	7 (2%)	0	100	100
2	B	174/182 (96%)	166 (95%)	8 (5%)	0	100	100
2	D	170/182 (93%)	163 (96%)	7 (4%)	0	100	100
2	F	169/182 (93%)	164 (97%)	5 (3%)	0	100	100
All	All	1435/1497 (96%)	1387 (97%)	46 (3%)	2 (0%)	48	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	142	SER
1	A	144	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/271 (98%)	261 (98%)	4 (2%)	60	81
1	C	264/271 (97%)	258 (98%)	6 (2%)	45	71
1	E	264/271 (97%)	260 (98%)	4 (2%)	60	81
2	B	150/155 (97%)	148 (99%)	2 (1%)	65	84
2	D	147/155 (95%)	145 (99%)	2 (1%)	62	82
2	F	146/155 (94%)	144 (99%)	2 (1%)	62	82
All	All	1236/1278 (97%)	1216 (98%)	20 (2%)	58	79

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	132	THR
1	A	155	LEU
1	A	321	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	58	LYS
2	B	66	ILE
1	C	18	HIS
1	C	75	GLN
1	C	152	LYS
1	C	190	GLU
1	C	240	ASN
1	C	321	ARG
2	D	12	ASN
2	D	19	ASN
1	E	75	GLN
1	E	141	ARG
1	E	155	LEU
1	E	173	ASN
2	F	65	LEU
2	F	67	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	HIS
1	A	216	ASN
2	B	26	HIS
2	F	26	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GAL	G	1	3	12,12,12	0.50	0	17,17,17	0.49	0
3	NAG	G	2	3	14,14,15	0.51	0	17,19,21	2.18	2 (11%)
3	GAL	G	3	3	11,11,12	0.56	0	15,15,17	0.68	0
3	SIA	G	4	3	20,20,21	4.07	2 (10%)	21,28,31	1.71	5 (23%)
4	NAG	H	1	4	13,13,15	0.72	0	15,17,21	0.85	0
4	SIA	H	2	4	20,20,21	4.11	2 (10%)	21,28,31	1.53	4 (19%)
4	NAG	I	1	4	13,13,15	0.78	1 (7%)	15,17,21	1.12	1 (6%)
4	SIA	I	2	4	20,20,21	4.08	2 (10%)	21,28,31	1.54	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GAL	G	1	3	-	2/2/22/22	0/1/1/1
3	NAG	G	2	3	1/1/5/7	0/6/23/26	0/1/1/1
3	GAL	G	3	3	-	0/2/19/22	0/1/1/1
3	SIA	G	4	3	-	3/18/34/38	0/1/1/1
4	NAG	H	1	4	-	0/6/19/26	0/1/1/1
4	SIA	H	2	4	-	4/18/34/38	0/1/1/1
4	NAG	I	1	4	-	0/6/19/26	0/1/1/1
4	SIA	I	2	4	-	2/18/34/38	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	2	SIA	C11-C10	-13.57	1.22	1.50
3	G	4	SIA	C11-C10	-13.50	1.22	1.50
4	I	2	SIA	C11-C10	-13.45	1.22	1.50
4	H	2	SIA	O10-C10	12.13	1.50	1.23
4	I	2	SIA	O10-C10	12.06	1.50	1.23
3	G	4	SIA	O10-C10	11.96	1.50	1.23
4	I	1	NAG	C1-C2	2.00	1.53	1.51

All (16) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	2	NAG	O3-C3-C2	7.71	125.41	109.40
3	G	4	SIA	C11-C10-N5	4.45	123.50	116.12
4	H	2	SIA	C11-C10-N5	4.26	123.19	116.12
4	I	2	SIA	C11-C10-N5	4.22	123.11	116.12
3	G	4	SIA	O10-C10-N5	-4.10	114.73	121.98
4	I	2	SIA	O10-C10-N5	-3.90	115.09	121.98
4	H	2	SIA	O10-C10-N5	-3.70	115.44	121.98
3	G	2	NAG	O3-C3-C4	3.51	118.64	110.38
4	I	1	NAG	C3-C2-C1	3.47	112.69	109.50
3	G	4	SIA	O6-C2-C1	2.89	113.17	107.72
3	G	4	SIA	O1B-C1-C2	2.70	119.75	112.71
4	H	2	SIA	O6-C2-C1	2.48	112.41	107.72
4	H	2	SIA	O1B-C1-C2	2.37	118.89	112.71
4	I	2	SIA	O1B-C1-C2	2.31	118.72	112.71
4	I	2	SIA	O6-C2-C1	2.26	111.98	107.72
3	G	4	SIA	O1A-C1-C2	-2.13	118.25	122.85

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	G	2	NAG	C3


All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	G	1	GAL	O5-C5-C6-O6
3	G	1	GAL	C4-C5-C6-O6
4	H	2	SIA	O8-C8-C9-O9
3	G	4	SIA	C11-C10-N5-C5
3	G	4	SIA	O10-C10-N5-C5
4	H	2	SIA	C11-C10-N5-C5
4	H	2	SIA	O10-C10-N5-C5
4	I	2	SIA	C11-C10-N5-C5
4	I	2	SIA	O10-C10-N5-C5
4	H	2	SIA	C7-C8-C9-O9
3	G	4	SIA	C6-C7-C8-O8

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	NAG	C	1	1	14,14,15	0.48	0	17,19,21	0.72	0
5	NAG	A	1	1	14,14,15	0.56	0	17,19,21	1.34	1 (5%)
6	GOL	B	184	-	5,5,5	0.35	0	5,5,5	0.32	0
5	NAG	B	183	2	14,14,15	0.50	0	17,19,21	1.05	1 (5%)
6	GOL	D	184	-	5,5,5	0.34	0	5,5,5	0.40	0
6	GOL	B	185	-	5,5,5	0.34	0	5,5,5	0.41	0
5	NAG	D	183	2	14,14,15	0.55	0	17,19,21	0.86	0
5	NAG	F	183	2	14,14,15	0.47	0	17,19,21	1.02	1 (5%)
6	GOL	F	184	-	5,5,5	0.34	0	5,5,5	0.40	0
5	NAG	E	1	1	14,14,15	0.56	0	17,19,21	1.13	1 (5%)
6	GOL	B	186	-	5,5,5	0.38	0	5,5,5	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1	1	-	2/6/23/26	0/1/1/1
5	NAG	A	1	1	-	2/6/23/26	0/1/1/1
6	GOL	B	184	-	-	0/4/4/4	-
5	NAG	B	183	2	-	0/6/23/26	0/1/1/1
6	GOL	D	184	-	-	4/4/4/4	-
6	GOL	B	185	-	-	2/4/4/4	-
5	NAG	D	183	2	-	1/6/23/26	0/1/1/1
5	NAG	F	183	2	-	0/6/23/26	0/1/1/1
6	GOL	F	184	-	-	2/4/4/4	-
5	NAG	E	1	1	-	2/6/23/26	0/1/1/1
6	GOL	B	186	-	-	0/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1	NAG	C2-N2-C7	3.64	127.78	122.90
5	F	183	NAG	C1-O5-C5	3.34	116.67	112.19
5	E	1	NAG	C2-N2-C7	3.24	127.24	122.90
5	B	183	NAG	C1-O5-C5	2.91	116.09	112.19

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	D	184	GOL	C1-C2-C3-O3
6	F	184	GOL	C1-C2-C3-O3
6	F	184	GOL	O2-C2-C3-O3
6	B	185	GOL	O2-C2-C3-O3
6	B	185	GOL	C1-C2-C3-O3
6	D	184	GOL	O1-C1-C2-C3
6	D	184	GOL	O1-C1-C2-O2
6	D	184	GOL	O2-C2-C3-O3
5	C	1	NAG	C4-C5-C6-O6
5	A	1	NAG	C1-C2-N2-C7
5	E	1	NAG	C3-C2-N2-C7
5	C	1	NAG	O5-C5-C6-O6
5	E	1	NAG	C1-C2-N2-C7
5	A	1	NAG	C3-C2-N2-C7
5	D	183	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	184	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	310/317 (97%)	-0.22	4 (1%) 74 70	6, 15, 25, 35	0
1	C	309/317 (97%)	-0.23	1 (0%) 90 88	4, 15, 24, 29	0
1	E	309/317 (97%)	-0.45	2 (0%) 85 83	5, 11, 19, 24	0
2	B	176/182 (96%)	-0.13	1 (0%) 85 83	8, 17, 26, 29	0
2	D	172/182 (94%)	-0.22	1 (0%) 85 83	4, 12, 20, 26	0
2	F	171/182 (93%)	-0.20	2 (1%) 76 72	7, 15, 22, 27	0
All	All	1447/1497 (96%)	-0.26	11 (0%) 82 79	4, 14, 24, 35	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	144	GLY	6.6
1	A	145	SER	3.7
2	F	133	ASP	2.9
2	B	176	GLY	2.8
2	F	19	ASN	2.7
2	D	154	ASN	2.6
1	A	142	SER	2.5
1	A	277	CYS	2.0
1	E	173	ASN	2.0
1	C	196	GLY	2.0
1	E	216	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	I	1	13/15	0.40	0.28	79,89,93,93	0
3	GAL	G	3	11/12	0.56	0.25	89,91,91,92	0
3	GAL	G	1	12/12	0.58	0.23	77,81,81,82	0
4	NAG	H	1	13/15	0.61	0.24	76,84,86,86	0
4	SIA	I	2	20/21	0.77	0.21	59,62,69,69	0
4	SIA	H	2	20/21	0.78	0.20	59,63,67,68	0
3	NAG	G	2	14/15	0.81	0.17	61,73,76,86	0
3	SIA	G	4	20/21	0.89	0.14	40,44,49,49	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain G:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	GOL	B	184	6/6	0.59	0.34	50,52,53,53	0
5	NAG	E	1	14/15	0.70	0.15	42,50,53,53	0
5	NAG	C	1	14/15	0.73	0.15	41,47,49,50	0
5	NAG	D	183	14/15	0.75	0.16	35,43,44,44	0
5	NAG	A	1	14/15	0.76	0.16	43,51,54,55	0
6	GOL	B	186	6/6	0.79	0.26	47,48,50,51	0
5	NAG	B	183	14/15	0.80	0.14	35,40,43,45	0
5	NAG	F	183	14/15	0.81	0.13	36,41,41,43	0
6	GOL	D	184	6/6	0.82	0.21	39,41,43,45	0
6	GOL	F	184	6/6	0.84	0.19	39,44,45,47	0
6	GOL	B	185	6/6	0.85	0.22	51,51,52,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.