



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 11:01 PM EDT

PDB ID : 3M6K  
Title : Crystal Structure of N-terminal 44 kDa fragment of topoisomerase V in the presence of guanidium hydrochloride  
Authors : Rajan, R.; Taneja, B.; Mondragon, A.  
Deposited on : 2010-03-15  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	
1	B	380	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Topoisomerase V.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	370	Total	C	N	O	S	0	0	0
			3040	1907	542	582	9			
1	B	357	Total	C	N	O	S	0	1	0
			2931	1842	517	563	9			

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total 20	O 20	0	0
3	B	10	Total 10	O 10	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Topoisomerase V



#### • Molecule 1: Topoisomerase V



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.09Å 70.09Å 349.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.14 – 2.60 29.14 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.14-2.60) 100.0 (29.14-2.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.82 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.241 , 0.289 0.243 , 0.292	Depositor DCC
$R_{free}$ test set	1431 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.0	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 37.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6016	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/3095	0.65	0/4185
1	B	0.54	1/2988 (0.0%)	0.67	0/4041
All	All	0.54	1/6083 (0.0%)	0.66	0/8226

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	361	GLY	C-O	7.42	1.35	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3040	0	3007	28	0
1	B	2931	0	2898	40	0
2	B	15	0	0	0	0
3	A	20	0	0	0	0
3	B	10	0	0	1	0
All	All	6016	0	5905	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:264:LEU:HD23	1:B:264:LEU:O	1.86	0.75
1:B:122:GLU:O	1:B:122:GLU:HG2	1.88	0.73
1:B:83:ARG:NE	1:B:325:MET:HE2	2.07	0.70
1:A:300:LYS:HD3	1:A:323:ILE:HD12	1.72	0.70
1:B:122:GLU:O	1:B:122:GLU:CG	2.39	0.70
1:B:109:ARG:HG3	1:B:325:MET:HE1	1.82	0.62
1:A:264:LEU:HD23	1:A:264:LEU:O	2.00	0.61
1:B:54:GLU:CG	1:B:159:VAL:HG11	2.31	0.60
1:B:83:ARG:HE	1:B:325:MET:HE2	1.69	0.57
1:B:177:TYR:O	1:B:179:VAL:N	2.39	0.56
1:B:54:GLU:HG3	1:B:159:VAL:HG11	1.88	0.55
1:B:115:ALA:HA	1:B:120:ILE:HD13	1.89	0.55
1:B:329:THR:HG23	1:B:330:PRO:HD2	1.88	0.55
1:B:177:TYR:O	1:B:179:VAL:HG13	2.07	0.54
1:B:243:ALA:O	1:B:247:VAL:HG23	2.08	0.54
1:B:155:MET:HE2	1:B:156:PRO:HD2	1.89	0.53
1:B:207:LEU:HB2	1:B:209:LEU:HD22	1.90	0.53
1:A:208:GLY:O	1:A:351:ARG:HD3	2.10	0.52
1:A:95:SER:OG	1:A:96:THR:HG23	2.11	0.50
1:A:262:GLU:O	1:A:265:SER:OG	2.28	0.50
1:B:155:MET:CE	1:B:156:PRO:HD2	2.42	0.50
1:B:103:TYR:HH	1:B:217:GLU:CD	2.16	0.48
1:A:373:HIS:O	1:A:374:PHE:C	2.52	0.48
1:A:295:VAL:O	1:A:299:LEU:HD22	2.14	0.47
1:A:120:ILE:HD13	1:A:120:ILE:N	2.28	0.47
1:A:269:LEU:O	1:A:269:LEU:HD12	2.14	0.47
1:B:222:SER:OG	1:B:223:SER:N	2.48	0.47
1:A:36:THR:OG1	1:A:56:HIS:ND1	2.34	0.46
1:B:281:HIS:HA	3:B:385:HOH:O	2.15	0.46
1:A:79:ASN:N	1:A:88:ASP:OD1	2.49	0.46
1:B:55:LEU:HD21	1:B:261:LEU:HB2	1.96	0.46
1:A:179:VAL:HG13	1:A:231:ARG:HD2	1.96	0.46
1:A:264:LEU:HD23	1:A:264:LEU:C	2.35	0.46
1:A:290:LEU:HD11	1:A:320:ASP:HB3	1.97	0.46
1:B:10:VAL:O	1:B:10:VAL:HG22	2.16	0.46
1:B:264:LEU:HD23	1:B:264:LEU:C	2.37	0.45
1:B:102:ARG:O	1:B:105:ILE:HG22	2.16	0.45
1:B:6:ASP:OD2	1:B:7:ALA:N	2.50	0.45

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:103:TYR:OH	1:B:217:GLU:CD	2.55	0.45
1:B:140:PRO:O	1:B:143:TYR:HB3	2.17	0.45
1:A:49:ASP:OD2	1:A:51:GLU:N	2.50	0.45
1:A:111:TRP:CD2	1:A:127:ALA:HA	2.53	0.44
1:A:152:VAL:HG11	1:A:247:VAL:HG23	1.99	0.44
1:B:102:ARG:HD2	1:B:147:VAL:O	2.18	0.44
1:B:204:ALA:HB1	1:B:209:LEU:O	2.17	0.44
1:B:109:ARG:HG3	1:B:325:MET:CE	2.47	0.44
1:A:49:ASP:OD2	1:A:49:ASP:C	2.57	0.43
1:B:199:PRO:HB2	1:B:201:ASP:OD1	2.17	0.43
1:A:327:ALA:O	1:A:350:ASN:CG	2.57	0.43
1:B:35:ALA:O	1:B:39:LEU:HD13	2.17	0.43
1:B:329:THR:CG2	1:B:330:PRO:HD2	2.48	0.42
1:B:31:ASP:OD2	1:B:123:GLU:OE2	2.37	0.42
1:B:58:ASN:HB3	1:B:257:LEU:HD11	2.01	0.42
1:B:118:LEU:HB2	1:B:120:ILE:CD1	2.50	0.42
1:A:271:HIS:O	1:A:271:HIS:CG	2.72	0.41
1:A:271:HIS:HA	1:A:272:PRO:HD2	1.92	0.41
1:A:252:ARG:O	1:A:256:GLU:HG2	2.21	0.41
1:B:356:LEU:HD13	1:B:367:ALA:HA	2.03	0.41
1:A:125:VAL:O	1:A:127:ALA:N	2.54	0.41
1:A:341:LEU:HD12	1:A:345:GLU:OE1	2.21	0.41
1:B:39:LEU:O	1:B:40:MET:O	2.39	0.41
1:A:20:ARG:HA	1:A:118:LEU:HD13	2.02	0.41
1:A:323:ILE:HA	1:A:326:ILE:HD12	2.01	0.41
1:A:38:TYR:N	1:A:289:TYR:OH	2.53	0.41
1:A:217:GLU:HB3	1:A:222:SER:HB2	2.03	0.40
1:B:125:VAL:HG12	1:B:125:VAL:O	2.21	0.40
1:B:207:LEU:CB	1:B:209:LEU:HD22	2.50	0.40
1:B:308:ARG:HD3	1:B:345:GLU:OE2	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/380 (96%)	345 (94%)	20 (6%)	1 (0%)	41	64
1	B	352/380 (93%)	335 (95%)	16 (4%)	1 (0%)	41	64
All	All	718/760 (94%)	680 (95%)	36 (5%)	2 (0%)	41	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	178	ASP
1	A	95	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/336 (98%)	323 (98%)	6 (2%)	59	80
1	B	318/336 (95%)	300 (94%)	18 (6%)	20	41
All	All	647/672 (96%)	623 (96%)	24 (4%)	35	60

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ARG
1	A	148	ARG
1	A	269	LEU
1	A	299	LEU
1	A	351	ARG
1	A	373	HIS
1	B	21	GLU
1	B	38	TYR
1	B	88	ASP
1	B	122	GLU
1	B	124	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	169	GLU
1	B	183[A]	GLU
1	B	183[B]	GLU
1	B	199	PRO
1	B	205	ARG
1	B	209	LEU
1	B	210	SER
1	B	227	SER
1	B	241	ARG
1	B	248	LYS
1	B	259	GLU
1	B	286	MET
1	B	290	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	B	381	-	4,4,4	1.03	0	6,6,6	0.81	0
2	PO4	B	383	-	4,4,4	0.89	0	6,6,6	0.65	0
2	PO4	B	382	-	4,4,4	1.10	0	6,6,6	0.82	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	370/380 (97%)	0.10	10 (2%) 54 48	36, 52, 72, 87	5 (1%)
1	B	357/380 (93%)	0.10	11 (3%) 49 42	33, 56, 79, 92	3 (0%)
All	All	727/760 (95%)	0.10	21 (2%) 51 45	33, 54, 76, 92	8 (1%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	ARG	4.0
1	A	93	ARG	3.6
1	A	119	ASP	3.6
1	B	183[A]	GLU	3.5
1	A	244	LYS	3.4
1	B	4	VAL	3.3
1	B	332	ARG	2.9
1	B	92	GLN	2.9
1	A	237	GLU	2.8
1	B	93	ARG	2.8
1	B	10	VAL	2.8
1	A	92	GLN	2.6
1	B	263	GLU	2.5
1	A	147	VAL	2.5
1	B	264	LEU	2.5
1	A	8	GLU	2.3
1	B	156	PRO	2.1
1	B	50	GLU	2.1
1	A	280	GLU	2.0
1	A	4	VAL	2.0
1	B	244	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	B	383	5/5	0.96	0.14	70,71,71,72	0
2	PO4	B	382	5/5	0.97	0.15	58,58,59,61	0
2	PO4	B	381	5/5	0.97	0.13	54,56,56,57	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.