



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2025 – 11:32 PM EDT

PDB ID : 6MEH / pdb_00006meh
Title : Crystal structure of broadly neutralizing antibody HEPC74 in complex with Hepatitis C virus envelope glycoprotein E2 ectodomain
Authors : Flyak, A.I.; Bjorkman, P.J.
Deposited on : 2018-09-06
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	262	<div> <div>6%</div> <div> <div></div> <div>55%</div> <div>8%</div> <div>37%</div> </div> </div>
2	H	239	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>8%</div> <div>6%</div> </div> </div>
3	L	214	<div> <div>%</div> <div> <div></div> <div>93%</div> <div>7%</div> </div> </div>
4	A	2	<div> <div></div> <div>100%</div> </div>
4	B	2	<div> <div></div> <div>50%</div> <div>50%</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5089 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E2 glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	165	Total	C	N	O	S	0	0	0
			1281	816	223	228	14			

- Molecule 2 is a protein called antibody HEPC74 Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	225	Total	C	N	O	S	0	0	0
			1663	1045	279	331	8			

- Molecule 3 is a protein called antibody HEPC74 Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	0	0
			1644	1034	274	331	5			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	A	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	B	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	65	Total	O	0	0
			65	65		
6	H	186	Total	O	0	0
			186	186		
6	L	152	Total	O	0	0
			152	152		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E2 glycoprotein




- Molecule 2: antibody HEPC74 Heavy Chain



- Molecule 3: antibody HEPC74 Light Chain



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  50% 50%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.31Å 69.77Å 168.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.76 – 1.99 43.76 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.5 (43.76-1.99) 98.6 (43.76-1.99)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.186 , 0.230 0.190 , 0.233	Depositor DCC
R_{free} test set	2723 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	36.0	Xtriage
Anisotropy	0.485	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.027 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5089	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.37	0/1326	0.56	0/1815
2	H	0.42	0/1701	0.59	0/2321
3	L	0.39	0/1682	0.58	0/2286
All	All	0.40	0/4709	0.58	0/6422

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	190	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1281	0	1187	14	0
2	H	1663	0	1640	10	0
3	L	1644	0	1599	12	0
4	A	28	0	25	0	0
4	B	28	0	25	0	0
5	C	42	0	39	1	0
6	C	65	0	0	0	1
6	H	186	0	0	3	1
6	L	152	0	0	6	0
All	All	5089	0	4515	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:117:LYS:NZ	6:H:303:HOH:O	2.08	0.85
2:H:214:LYS:O	6:H:302:HOH:O	2.02	0.76
1:C:516:VAL:HG22	1:C:542:THR:HG21	1.77	0.67
1:C:531:GLU:OE2	2:H:100(B):GLY:N	2.24	0.64
3:L:123:GLU:HG3	6:L:374:HOH:O	1.98	0.64
2:H:61:GLN:HG3	2:H:64:GLN:HB2	1.83	0.60
3:L:108:ARG:HH21	3:L:109:THR:HG23	1.68	0.59
2:H:62:LYS:N	6:H:301:HOH:O	1.97	0.56
3:L:24:ARG:NH1	3:L:70:GLU:OE2	2.40	0.55
3:L:37:GLN:HB2	3:L:47:LEU:HD11	1.90	0.53
1:C:506:VAL:HG21	1:C:537:LEU:HD13	1.92	0.52
1:C:422:ILE:HD13	1:C:424:ARG:HB2	1.92	0.51
3:L:33:LEU:HG	3:L:34:ALA:N	2.28	0.48
3:L:2:ILE:N	6:L:301:HOH:O	2.16	0.47
3:L:152:ASN:CB	6:L:305:HOH:O	2.64	0.46
1:C:548:ASN:ND2	1:C:568:PRO:HG3	2.30	0.46
1:C:443:TYR:HB3	1:C:445:HIS:CE1	2.50	0.46
2:H:57:PRO:HB2	2:H:59:TYR:CE1	2.51	0.45
1:C:502:VAL:HG21	1:C:537:LEU:CD1	2.46	0.45
3:L:126:LYS:HB3	3:L:126:LYS:HE3	1.67	0.44
3:L:33:LEU:HD22	3:L:71:PHE:CG	2.53	0.44
2:H:100:CYS:HA	2:H:100(E):CYS:HA	2.00	0.43
1:C:603:LEU:HD22	1:C:631:MET:CE	2.49	0.43
1:C:509:PHE:O	1:C:549:TRP:CZ3	2.72	0.43
3:L:152:ASN:CG	6:L:305:HOH:O	2.57	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:519:THR:HG22	1:C:525:PRO:HA	2.01	0.42
1:C:422:ILE:C	1:C:422:ILE:HD12	2.39	0.42
2:H:93:ALA:HB2	2:H:103:TRP:CD2	2.55	0.42
1:C:423:ASN:HB2	1:C:527:TYR:O	2.20	0.41
2:H:17:SER:HB3	2:H:82(A):SER:HA	2.02	0.41
2:H:30:ILE:HG22	2:H:31:ASN:ND2	2.35	0.41
3:L:2:ILE:HB	6:L:301:HOH:O	2.21	0.41
1:C:622:VAL:HG12	5:C:707:NAG:H82	2.03	0.41
1:C:509:PHE:O	1:C:549:TRP:HZ3	2.04	0.40
3:L:152:ASN:HB2	6:L:305:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:C:854:HOH:O	6:H:335:HOH:O[3_654]	2.04	0.16

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	159/262 (61%)	153 (96%)	5 (3%)	1 (1%)	22	17
2	H	221/239 (92%)	217 (98%)	4 (2%)	0	100	100
3	L	211/214 (99%)	205 (97%)	6 (3%)	0	100	100
All	All	591/715 (83%)	575 (97%)	15 (2%)	1 (0%)	44	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	510	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	140/219 (64%)	139 (99%)	1 (1%)	81	86
2	H	190/204 (93%)	187 (98%)	3 (2%)	58	64
3	L	188/189 (100%)	187 (100%)	1 (0%)	86	90
All	All	518/612 (85%)	513 (99%)	5 (1%)	73	78

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	549	TRP
2	H	35	SER
2	H	59	TYR
2	H	96	LEU
3	L	105	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	64	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1	1,4	14,14,15	0.35	0	17,19,21	0.70	0
4	NAG	A	2	4	14,14,15	0.31	0	17,19,21	0.62	0
4	NAG	B	1	1,4	14,14,15	0.52	0	17,19,21	1.01	2 (11%)
4	NAG	B	2	4	14,14,15	0.35	0	17,19,21	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	2	4	-	5/6/23/26	0/1/1/1
4	NAG	B	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	2	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1	NAG	O5-C1-C2	-2.43	107.53	111.29
4	B	1	NAG	C1-O5-C5	2.17	115.09	112.19

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	2	NAG	C1-C2-N2-C7
4	A	2	NAG	C8-C7-N2-C2
4	A	2	NAG	O7-C7-N2-C2
4	A	2	NAG	O5-C5-C6-O6
4	A	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	C	701	1	14,14,15	0.29	0	17,19,21	0.66	0
5	NAG	C	704	1	14,14,15	0.37	0	17,19,21	0.92	1 (5%)
5	NAG	C	707	1	14,14,15	0.34	0	17,19,21	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	701	1	-	0/6/23/26	0/1/1/1
5	NAG	C	704	1	-	0/6/23/26	0/1/1/1
5	NAG	C	707	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	704	NAG	C2-N2-C7	-2.11	120.07	122.90

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	707	NAG	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	707	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	165/262 (62%)	0.59	15 (9%) 16 15	36, 57, 101, 134	0
2	H	225/239 (94%)	-0.07	8 (3%) 46 44	24, 39, 68, 86	0
3	L	213/214 (99%)	-0.04	2 (0%) 81 80	25, 47, 69, 93	0
All	All	603/715 (84%)	0.12	25 (4%) 42 40	24, 47, 84, 134	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	59	TYR	4.3
2	H	126	PRO	4.1
1	C	443	TYR	3.6
1	C	491	PRO	3.6
1	C	569	CYS	3.5
2	H	215	SER	3.2
2	H	132	SER	3.1
3	L	94	TYR	2.9
1	C	549	TRP	2.9
1	C	521	ARG	2.9
2	H	127	SER	2.9
2	H	190	GLY	2.6
1	C	534	THR	2.6
1	C	497	ILE	2.4
2	H	61	GLN	2.4
1	C	418	GLY	2.4
1	C	546	GLN	2.4
2	H	133	GLY	2.3
1	C	598	GLY	2.2
3	L	109	THR	2.2
1	C	631	MET	2.2
1	C	635	GLY	2.1
1	C	522	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	454	GLU	2.0
1	C	540	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NAG	C	701	14/15	0.74	0.12	61,78,91,99	0
5	NAG	C	707	14/15	0.74	0.13	86,92,105,106	0
5	NAG	C	704	14/15	0.93	0.07	32,46,57,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.