



wwPDB EM Validation Summary Report ⓘ

Dec 16, 2024 – 07:32 PM EST

PDB ID : 7ML4
EMDB ID : EMD-23908
Title : RNA polymerase II initially transcribing complex (ITC)
Authors : Yang, C.; Fujiwara, R.; Kim, H.J.; Gorbea Colon, J.J.; Steimle, S.; Garcia, B.A.; Murakami, K.
Deposited on : 2021-04-27
Resolution : 3.10 Å(reported)
Based on initial model : 5OQJ

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|--------------------------|--------------------------|
| Clashscore | 210492 | 15764 |
| Ramachandran outliers | 207382 | 16835 |
| Sidechain outliers | 206894 | 16415 |
| RNA backbone | 6643 | 2191 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---------------------------------------|
| 1 | Q | 735 | <div><div>7%11%•80%</div></div> |
| 2 | R | 398 | <div><div>5%18%18%•61%</div></div> |
| 3 | D | 221 | <div><div>30%27%41%•29%</div></div> |
| 4 | G | 171 | <div><div>18%32%60%8%</div></div> |
| 5 | M | 345 | <div><div>11%26%36%5%•32%</div></div> |
| 6 | A | 1733 | <div><div>•36%38%7%19%</div></div> |
| 7 | B | 1224 | <div><div>•45%38%8%9%</div></div> |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 8 | C | 318 | |
| 9 | E | 215 | |
| 10 | F | 155 | |
| 11 | H | 146 | |
| 12 | I | 122 | |
| 13 | J | 70 | |
| 14 | K | 120 | |
| 15 | L | 70 | |
| 16 | 3 | 321 | |
| 17 | 0 | 778 | |
| 18 | 4 | 338 | |
| 19 | 6 | 461 | |
| 20 | 1 | 543 | |
| 21 | 7 | 843 | |
| 22 | 5 | 72 | |
| 23 | 2 | 513 | |
| 24 | X | 328 | |
| 25 | U | 286 | |
| 26 | V | 122 | |
| 27 | N | 38 | |
| 28 | T | 148 | |
| 29 | O | 240 | |
| 30 | W | 482 | |
| 31 | P | 5 | |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 34 | SF4 | 0 | 801 | - | - | X | - |

2 Entry composition

There are 34 unique types of molecules in this entry. The entry contains 62733 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription initiation factor IIF subunit alpha.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 1 | Q | 148 | Total | C | N | O | S | 0 | 0 |
| | | | 1141 | 731 | 195 | 212 | 3 | | |

- Molecule 2 is a protein called Transcription initiation factor IIF subunit beta.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 2 | R | 154 | Total | C | N | O | S | 0 | 0 |
| | | | 1039 | 652 | 190 | 193 | 4 | | |

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 3 | D | 157 | Total | C | N | O | S | 0 | 0 |
| | | | 1253 | 779 | 220 | 252 | 2 | | |

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB7.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 4 | G | 171 | Total | C | N | O | S | 0 | 0 |
| | | | 1340 | 861 | 222 | 249 | 8 | | |

- Molecule 5 is a protein called Transcription initiation factor IIB.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 5 | M | 234 | Total | C | N | O | S | 0 | 0 |
| | | | 1805 | 1152 | 304 | 333 | 16 | | |

- Molecule 6 is a protein called DNA-directed RNA polymerase subunit.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| 6 | A | 1405 | Total | C | N | O | S | 0 | 0 |
| | | | 11039 | 6962 | 1935 | 2081 | 61 | | |

- Molecule 7 is a protein called DNA-directed RNA polymerase subunit beta.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| 7 | B | 1114 | Total | C | N | O | S | 0 | 0 |
| | | | 8861 | 5610 | 1549 | 1647 | 55 | | |

- Molecule 8 is a protein called DNA-directed RNA polymerase II subunit RPB3.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 8 | C | 266 | Total | C | N | O | S | 0 | 0 |
| | | | 2095 | 1317 | 348 | 417 | 13 | | |

- Molecule 9 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 9 | E | 214 | Total | C | N | O | S | 0 | 0 |
| | | | 1752 | 1111 | 309 | 321 | 11 | | |

- Molecule 10 is a protein called DNA-directed RNA polymerases I,II,and III subunit RPABC2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 10 | F | 85 | Total | C | N | O | S | 0 | 0 |
| | | | 688 | 439 | 116 | 130 | 3 | | |

- Molecule 11 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 11 | H | 133 | Total | C | N | O | S | 0 | 0 |
| | | | 1068 | 673 | 180 | 211 | 4 | | |

- Molecule 12 is a protein called DNA-directed RNA polymerase II subunit RPB9.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
| 12 | I | 119 | Total | C | N | O | S | 0 | 0 |
| | | | 971 | 596 | 179 | 186 | 10 | | |

- Molecule 13 is a protein called DNA-directed RNA polymerases II subunit RPABC5.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 13 | J | 65 | Total | C | N | O | S | 0 | 0 |
| | | | 532 | 339 | 93 | 94 | 6 | | |

- Molecule 14 is a protein called DNA-directed RNA polymerase II subunit RPB11.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 14 | K | 114 | Total | C | N | O | S | 0 | 0 |
| | | | 919 | 590 | 156 | 171 | 2 | | |

- Molecule 15 is a protein called DNA-directed RNA polymerases II subunit RPABC4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 15 | L | 46 | Total | C | N | O | S | 0 | 0 |
| | | | 363 | 224 | 72 | 63 | 4 | | |

- Molecule 16 is a protein called BJ4_G0050160.mRNA.1.CDS.1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 16 | 3 | 72 | Total | C | N | O | S | 0 | 0 |
| | | | 361 | 215 | 72 | 74 | | | |

- Molecule 17 is a protein called General transcription and DNA repair factor IIIH helicase subunit XPD.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| 17 | 0 | 754 | Total | C | N | O | S | 0 | 0 |
| | | | 6108 | 3891 | 1032 | 1147 | 38 | | |

- Molecule 18 is a protein called General transcription and DNA repair factor IIIH subunit TFB4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 18 | 4 | 284 | Total | C | N | O | S | 0 | 0 |
| | | | 2041 | 1310 | 343 | 376 | 12 | | |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|----------------|
| 4 | 113 | UNK | ASP | conflict | UNP A0A7I9C5C2 |
| 4 | 114 | UNK | MET | conflict | UNP A0A7I9C5C2 |

- Molecule 19 is a protein called General transcription and DNA repair factor IIIH.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 19 | 6 | 351 | Total | C | N | O | S | 0 | 0 |
| | | | 2527 | 1590 | 454 | 456 | 27 | | |

There are 13 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|----------------|
| 6 | 412 | UNK | ILE | conflict | UNP A0A7I9FQL5 |
| 6 | 413 | UNK | LEU | conflict | UNP A0A7I9FQL5 |
| 6 | 414 | UNK | LYS | conflict | UNP A0A7I9FQL5 |
| 6 | 415 | UNK | ASN | conflict | UNP A0A7I9FQL5 |
| 6 | 416 | UNK | HIS | conflict | UNP A0A7I9FQL5 |
| 6 | 417 | UNK | LYS | conflict | UNP A0A7I9FQL5 |
| 6 | 418 | UNK | ASN | conflict | UNP A0A7I9FQL5 |
| 6 | 419 | UNK | ASP | conflict | UNP A0A7I9FQL5 |
| 6 | 420 | UNK | LYS | conflict | UNP A0A7I9FQL5 |
| 6 | 421 | UNK | LEU | conflict | UNP A0A7I9FQL5 |
| 6 | 422 | UNK | LEU | conflict | UNP A0A7I9FQL5 |
| 6 | 423 | UNK | THR | conflict | UNP A0A7I9FQL5 |
| 6 | 424 | UNK | SER | conflict | UNP A0A7I9FQL5 |

- Molecule 20 is a protein called Tfb1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 20 | 1 | 367 | Total | C | N | O | S | 0 | 0 |
| | | | 2411 | 1536 | 438 | 430 | 7 | | |

- Molecule 21 is a protein called General transcription and DNA repair factor IIH helicase subunit XPB.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 21 | 7 | 634 | Total | C | N | O | S | 0 | 0 |
| | | | 4447 | 2722 | 827 | 874 | 24 | | |

- Molecule 22 is a protein called General transcription and DNA repair factor IIH subunit TFB5.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 22 | 5 | 66 | Total | C | N | O | S | 0 | 0 |
| | | | 498 | 314 | 89 | 93 | 2 | | |

- Molecule 23 is a protein called RNA polymerase II transcription factor B subunit 2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 23 | 2 | 460 | Total | C | N | O | S | 0 | 0 |
| | | | 3011 | 1856 | 562 | 584 | 9 | | |

- Molecule 24 is a protein called Transcription initiation factor IIE subunit beta.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 24 | X | 149 | Total | C | N | O | S | 0 | 0 |
| | | | 921 | 569 | 168 | 180 | 4 | | |

- Molecule 25 is a protein called Transcription initiation factor IIA large subunit.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 25 | U | 46 | Total | C | N | O | S | 0 | 0 |
| | | | 383 | 242 | 67 | 71 | 3 | | |

- Molecule 26 is a protein called Transcription initiation factor IIA subunit 2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 26 | V | 49 | Total | C | N | O | S | 0 | 0 |
| | | | 381 | 241 | 63 | 74 | 3 | | |

- Molecule 27 is a DNA chain called non-template strand DNA.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
| 27 | N | 38 | Total | C | N | O | P | 0 | 0 |
| | | | 791 | 376 | 161 | 216 | 38 | | |

- Molecule 28 is a DNA chain called template strand DNA.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
| 28 | T | 48 | Total | C | N | O | P | 0 | 0 |
| | | | 968 | 468 | 150 | 302 | 48 | | |

- Molecule 29 is a protein called TATA-box-binding protein.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 29 | O | 180 | Total | C | N | O | S | 0 | 0 |
| | | | 1416 | 921 | 242 | 247 | 6 | | |

- Molecule 30 is a protein called Transcription initiation factor IIE subunit alpha.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 30 | W | 191 | Total | C | N | O | S | 0 | 0 |
| | | | 1469 | 932 | 254 | 277 | 6 | | |

- Molecule 31 is a RNA chain called RNA.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|-------|
| 31 | P | 5 | Total | C | N | O | P | 0 | 0 |
| | | | 110 | 50 | 25 | 31 | 4 | | |


- Molecule 32 is ZINC ION (three-letter code: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|-------|----|---------|
| 32 | M | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |
| 32 | A | 2 | Total | Zn | 0 |
| | | | 2 | 2 | |
| 32 | B | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |
| 32 | C | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |
| 32 | I | 2 | Total | Zn | 0 |
| | | | 2 | 2 | |
| 32 | J | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |
| 32 | L | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |
| 32 | 4 | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |
| 32 | 6 | 4 | Total | Zn | 0 |
| | | | 4 | 4 | |
| 32 | W | 1 | Total | Zn | 0 |
| | | | 1 | 1 | |

- Molecule 33 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|-------|----|---------|
| 33 | A | 1 | Total | Mg | 0 |
| | | | 1 | 1 | |

- Molecule 34 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




| Mol | Chain | Residues | Atoms | | | AltConf |
|-----|-------|----------|-------|----|---|---------|
| | | | Total | Fe | S | |
| 34 | 0 | 1 | 8 | 4 | 4 | 0 |


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription initiation factor IIF subunit alpha




- Molecule 2: Transcription initiation factor IIF subunit beta




- Molecule 3: DNA-directed RNA polymerase II subunit RPB4





- Molecule 4: DNA-directed RNA polymerase II subunit RPB7






• Molecule 5: Transcription initiation factor IIB




• Molecule 6: DNA-directed RNA polymerase subunit
















- Molecule 10: DNA-directed RNA polymerases I,II,and III subunit RPABC2

Chain F:




- Molecule 11: DNA-directed RNA polymerases I, II, and III subunit RPABC3

Chain H:



- Molecule 12: DNA-directed RNA polymerase II subunit RPB9

Chain I:




- Molecule 13: DNA-directed RNA polymerases II subunit RPABC5


Chain J:




- Molecule 14: DNA-directed RNA polymerase II subunit RPB11










- Molecule 21: General transcription and DNA repair factor IIH helicase subunit XPB








- Molecule 22: General transcription and DNA repair factor IIH subunit TFB5




- Molecule 23: RNA polymerase II transcription factor B subunit 2




- Molecule 24: Transcription initiation factor IIE subunit beta







• Molecule 25: Transcription initiation factor IIA large subunit



• Molecule 26: Transcription initiation factor IIA subunit 2



• Molecule 27: non-template strand DNA



• Molecule 28: template strand DNA

ASP VAL MET ASP ASP ASP GLU THR ALA GLU ASN ALA LEU GLU ASP ASP ASP PHE GLU ASP VAL THR ASP THR ALA GLY THR ALA LYS THR GLU SER ASN THR SER ASN ASP VAL LYS GLN GLU SER THR ASN ASP LYS THR GLU ASP ALA VAL ASN ALA THR ALA THR ALA SER GLY

PRO SER ALA ASN ALA LYS PRO ASN ASP GLY ASP ASP ASP ASP ASP ASP ASP MET ASP THR ILE GLU PHE GLU VAL

● Molecule 31: RNA



G6 A7 G8 G9 A10

4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, Not provided | |
| Number of particles used | 254448 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 45 | Depositor |
| Minimum defocus (nm) | 500 | Depositor |
| Maximum defocus (nm) | 2500 | Depositor |
| Magnification | Not provided | |
| Image detector | GATAN K3 BIOQUANTUM (6k x 4k) | Depositor |
| Maximum map value | 0.057 | Depositor |
| Minimum map value | -0.001 | Depositor |
| Average map value | 0.001 | Depositor |
| Map value standard deviation | 0.002 | Depositor |
| Recommended contour level | 0.015 | Depositor |
| Map size (Å) | 444.13998, 503.49997, 508.8 | wwPDB |
| Map dimensions | 419, 475, 480 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 1.06, 1.06, 1.06 | Depositor |

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | Q | 0.49 | 0/1165 | 0.55 | 0/1576 |
| 2 | R | 0.35 | 0/1047 | 0.50 | 0/1422 |
| 3 | D | 0.32 | 0/1262 | 0.52 | 0/1693 |
| 4 | G | 0.53 | 0/1368 | 0.57 | 0/1844 |
| 5 | M | 0.40 | 0/1828 | 0.54 | 0/2459 |
| 6 | A | 1.01 | 5/11237 (0.0%) | 0.74 | 4/15195 (0.0%) |
| 7 | B | 1.08 | 0/9033 | 0.73 | 1/12181 (0.0%) |
| 8 | C | 0.98 | 0/2133 | 0.71 | 0/2891 |
| 9 | E | 0.94 | 0/1788 | 0.65 | 1/2406 (0.0%) |
| 10 | F | 1.17 | 0/700 | 0.76 | 1/945 (0.1%) |
| 11 | H | 0.95 | 0/1086 | 0.74 | 0/1470 |
| 12 | I | 0.76 | 0/989 | 0.63 | 0/1331 |
| 13 | J | 1.19 | 0/541 | 0.79 | 0/727 |
| 14 | K | 0.94 | 0/937 | 0.66 | 0/1265 |
| 15 | L | 0.82 | 0/365 | 0.77 | 0/485 |
| 16 | 3 | 0.23 | 0/360 | 0.35 | 0/501 |
| 17 | 0 | 0.25 | 0/6226 | 0.43 | 0/8407 |
| 18 | 4 | 0.26 | 0/2062 | 0.46 | 0/2805 |
| 19 | 6 | 0.25 | 0/2506 | 0.44 | 0/3402 |
| 20 | 1 | 0.24 | 0/1896 | 0.39 | 0/2543 |
| 21 | 7 | 0.25 | 0/4521 | 0.43 | 0/6036 |
| 22 | 5 | 0.23 | 0/502 | 0.42 | 0/677 |
| 23 | 2 | 0.24 | 0/3057 | 0.42 | 0/4071 |
| 24 | X | 0.25 | 0/929 | 0.43 | 0/1272 |
| 25 | U | 0.25 | 0/389 | 0.48 | 0/523 |
| 26 | V | 0.26 | 0/384 | 0.44 | 0/518 |
| 27 | N | 0.53 | 0/893 | 1.35 | 16/1377 (1.2%) |
| 28 | T | 0.51 | 0/1076 | 1.45 | 24/1654 (1.5%) |
| 29 | O | 0.29 | 0/1443 | 0.47 | 0/1942 |
| 30 | W | 0.27 | 0/1490 | 0.44 | 0/2014 |
| 31 | P | 0.69 | 0/124 | 1.59 | 3/193 (1.6%) |
| All | All | 0.73 | 5/63337 (0.0%) | 0.65 | 50/85825 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | Q | 0 | 1 |
| 4 | G | 0 | 1 |
| 5 | M | 0 | 3 |
| 6 | A | 0 | 9 |
| 7 | B | 0 | 5 |
| 10 | F | 0 | 1 |
| 11 | H | 0 | 3 |
| All | All | 0 | 23 |

All (5) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 6 | A | 786 | HIS | CA-C | -5.82 | 1.37 | 1.52 |
| 6 | A | 656 | TRP | CB-CG | -5.68 | 1.40 | 1.50 |
| 6 | A | 55 | ASP | C-N | -5.65 | 1.23 | 1.34 |
| 6 | A | 512 | VAL | C-N | -5.34 | 1.21 | 1.34 |
| 6 | A | 556 | TRP | CB-CG | -5.17 | 1.41 | 1.50 |

The worst 5 of 50 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|--------|-------------|----------|
| 27 | N | 23 | DA | O4'-C4'-C3' | -12.58 | 98.45 | 106.00 |
| 28 | T | 20 | DC | O4'-C4'-C3' | -8.81 | 100.72 | 106.00 |
| 27 | N | 23 | DA | C4'-C3'-C2' | -8.56 | 95.39 | 103.10 |
| 27 | N | 16 | DC | O4'-C1'-N1 | 8.37 | 113.86 | 108.00 |
| 28 | T | 19 | DT | O4'-C4'-C3' | -8.21 | 101.07 | 106.00 |

There are no chirality outliers.

5 of 23 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 4 | G | 56 | ILE | Peptide |
| 5 | M | 267 | LYS | Peptide |
| 5 | M | 268 | GLU | Peptide |
| 5 | M | 270 | ALA | Peptide |
| 1 | Q | 125 | LYS | Peptide |

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | Q | 1141 | 0 | 1027 | 130 | 0 |
| 2 | R | 1039 | 0 | 857 | 124 | 0 |
| 3 | D | 1253 | 0 | 1275 | 102 | 0 |
| 4 | G | 1340 | 0 | 1357 | 118 | 0 |
| 5 | M | 1805 | 0 | 1895 | 115 | 0 |
| 6 | A | 11039 | 0 | 11122 | 727 | 0 |
| 7 | B | 8861 | 0 | 8884 | 494 | 0 |
| 8 | C | 2095 | 0 | 2051 | 112 | 0 |
| 9 | E | 1752 | 0 | 1776 | 112 | 0 |
| 10 | F | 688 | 0 | 707 | 73 | 0 |
| 11 | H | 1068 | 0 | 1040 | 123 | 0 |
| 12 | I | 971 | 0 | 927 | 96 | 0 |
| 13 | J | 532 | 0 | 542 | 27 | 0 |
| 14 | K | 919 | 0 | 929 | 47 | 0 |
| 15 | L | 363 | 0 | 389 | 50 | 0 |
| 16 | 3 | 361 | 0 | 150 | 4 | 0 |
| 17 | 0 | 6108 | 0 | 6167 | 270 | 0 |
| 18 | 4 | 2041 | 0 | 1954 | 108 | 0 |
| 19 | 6 | 2527 | 0 | 2321 | 96 | 0 |
| 20 | 1 | 2411 | 0 | 1879 | 57 | 0 |
| 21 | 7 | 4447 | 0 | 3905 | 197 | 0 |
| 22 | 5 | 498 | 0 | 506 | 11 | 0 |
| 23 | 2 | 3011 | 0 | 2600 | 82 | 0 |
| 24 | X | 921 | 0 | 650 | 33 | 0 |
| 25 | U | 383 | 0 | 384 | 26 | 0 |
| 26 | V | 381 | 0 | 388 | 22 | 0 |
| 27 | N | 791 | 0 | 426 | 40 | 0 |
| 28 | T | 968 | 0 | 552 | 55 | 0 |
| 29 | O | 1416 | 0 | 1493 | 151 | 0 |
| 30 | W | 1469 | 0 | 1433 | 88 | 0 |
| 31 | P | 110 | 0 | 56 | 13 | 0 |
| 32 | 4 | 1 | 0 | 0 | 0 | 0 |
| 32 | 6 | 4 | 0 | 0 | 0 | 0 |
| 32 | A | 2 | 0 | 0 | 0 | 0 |
| 32 | B | 1 | 0 | 0 | 0 | 0 |
| 32 | C | 1 | 0 | 0 | 0 | 0 |
| 32 | I | 2 | 0 | 0 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 32 | J | 1 | 0 | 0 | 0 | 0 |
| 32 | L | 1 | 0 | 0 | 0 | 0 |
| 32 | M | 1 | 0 | 0 | 0 | 0 |
| 32 | W | 1 | 0 | 0 | 0 | 0 |
| 33 | A | 1 | 0 | 0 | 0 | 0 |
| 34 | 0 | 8 | 0 | 0 | 2 | 0 |
| All | All | 62733 | 0 | 59642 | 3374 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 3374 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|-----------------|--------------------------|-------------------|
| 6:A:68:GLN:NE2 | 6:A:80:HIS:ND1 | 1.92 | 1.16 |
| 11:H:55:LEU:H | 11:H:146:ARG:HA | 1.12 | 1.10 |
| 6:A:1444:MET:HA | 10:F:133:VAL:HA | 1.19 | 1.09 |
| 6:A:68:GLN:NE2 | 6:A:80:HIS:CE1 | 2.23 | 1.07 |
| 6:A:68:GLN:HE22 | 6:A:80:HIS:CG | 1.74 | 1.05 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|----------|-------------|-----|
| 1 | Q | 140/735 (19%) | 119 (85%) | 21 (15%) | 0 | 100 | 100 |
| 2 | R | 142/398 (36%) | 126 (89%) | 16 (11%) | 0 | 100 | 100 |
| 3 | D | 153/221 (69%) | 140 (92%) | 13 (8%) | 0 | 100 | 100 |
| 4 | G | 169/171 (99%) | 156 (92%) | 13 (8%) | 0 | 100 | 100 |
| 5 | M | 228/345 (66%) | 196 (86%) | 29 (13%) | 3 (1%) | 10 | 36 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|------------------|------------|-----------|----------|-------------|-----|
| 6 | A | 1395/1733 (80%) | 1227 (88%) | 161 (12%) | 7 (0%) | 25 | 58 |
| 7 | B | 1096/1224 (90%) | 942 (86%) | 149 (14%) | 5 (0%) | 25 | 58 |
| 8 | C | 264/318 (83%) | 241 (91%) | 22 (8%) | 1 (0%) | 30 | 63 |
| 9 | E | 212/215 (99%) | 199 (94%) | 13 (6%) | 0 | 100 | 100 |
| 10 | F | 83/155 (54%) | 73 (88%) | 10 (12%) | 0 | 100 | 100 |
| 11 | H | 129/146 (88%) | 97 (75%) | 31 (24%) | 1 (1%) | 16 | 48 |
| 12 | I | 117/122 (96%) | 95 (81%) | 22 (19%) | 0 | 100 | 100 |
| 13 | J | 63/70 (90%) | 52 (82%) | 11 (18%) | 0 | 100 | 100 |
| 14 | K | 112/120 (93%) | 101 (90%) | 11 (10%) | 0 | 100 | 100 |
| 15 | L | 44/70 (63%) | 30 (68%) | 14 (32%) | 0 | 100 | 100 |
| 16 | 3 | 70/321 (22%) | 65 (93%) | 5 (7%) | 0 | 100 | 100 |
| 17 | 0 | 752/778 (97%) | 701 (93%) | 51 (7%) | 0 | 100 | 100 |
| 18 | 4 | 279/338 (82%) | 246 (88%) | 33 (12%) | 0 | 100 | 100 |
| 19 | 6 | 336/461 (73%) | 295 (88%) | 39 (12%) | 2 (1%) | 22 | 53 |
| 20 | 1 | 256/543 (47%) | 237 (93%) | 17 (7%) | 2 (1%) | 16 | 48 |
| 21 | 7 | 630/843 (75%) | 578 (92%) | 52 (8%) | 0 | 100 | 100 |
| 22 | 5 | 64/72 (89%) | 59 (92%) | 5 (8%) | 0 | 100 | 100 |
| 23 | 2 | 456/513 (89%) | 412 (90%) | 44 (10%) | 0 | 100 | 100 |
| 24 | X | 145/328 (44%) | 126 (87%) | 19 (13%) | 0 | 100 | 100 |
| 25 | U | 44/286 (15%) | 39 (89%) | 5 (11%) | 0 | 100 | 100 |
| 26 | V | 45/122 (37%) | 45 (100%) | 0 | 0 | 100 | 100 |
| 29 | O | 178/240 (74%) | 165 (93%) | 13 (7%) | 0 | 100 | 100 |
| 30 | W | 189/482 (39%) | 179 (95%) | 10 (5%) | 0 | 100 | 100 |
| All | All | 7791/11370 (68%) | 6941 (89%) | 829 (11%) | 21 (0%) | 38 | 68 |

5 of 21 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 5 | M | 269 | ILE |
| 5 | M | 270 | ALA |
| 6 | A | 68 | GLN |
| 6 | A | 72 | GLU |
| 6 | A | 609 | ASP |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|-------------|-----|
| 1 | Q | 108/641 (17%) | 84 (78%) | 24 (22%) | 1 | 3 |
| 2 | R | 78/362 (22%) | 62 (80%) | 16 (20%) | 1 | 4 |
| 3 | D | 139/200 (70%) | 126 (91%) | 13 (9%) | 7 | 27 |
| 4 | G | 152/152 (100%) | 137 (90%) | 15 (10%) | 6 | 24 |
| 5 | M | 202/299 (68%) | 171 (85%) | 31 (15%) | 2 | 10 |
| 6 | A | 1224/1520 (80%) | 1032 (84%) | 192 (16%) | 2 | 9 |
| 7 | B | 967/1061 (91%) | 833 (86%) | 134 (14%) | 3 | 12 |
| 8 | C | 234/274 (85%) | 198 (85%) | 36 (15%) | 2 | 9 |
| 9 | E | 196/197 (100%) | 169 (86%) | 27 (14%) | 3 | 13 |
| 10 | F | 75/137 (55%) | 63 (84%) | 12 (16%) | 2 | 9 |
| 11 | H | 117/128 (91%) | 95 (81%) | 22 (19%) | 1 | 5 |
| 12 | I | 113/116 (97%) | 83 (74%) | 30 (26%) | 0 | 1 |
| 13 | J | 60/65 (92%) | 51 (85%) | 9 (15%) | 2 | 10 |
| 14 | K | 99/102 (97%) | 89 (90%) | 10 (10%) | 6 | 23 |
| 15 | L | 40/57 (70%) | 28 (70%) | 12 (30%) | 0 | 0 |
| 16 | 3 | 1/303 (0%) | 1 (100%) | 0 | 100 | 100 |
| 17 | 0 | 686/707 (97%) | 657 (96%) | 29 (4%) | 25 | 56 |
| 18 | 4 | 198/298 (66%) | 185 (93%) | 13 (7%) | 14 | 41 |
| 19 | 6 | 247/406 (61%) | 237 (96%) | 10 (4%) | 27 | 58 |
| 20 | 1 | 169/396 (43%) | 160 (95%) | 9 (5%) | 19 | 48 |
| 21 | 7 | 414/737 (56%) | 394 (95%) | 20 (5%) | 21 | 51 |
| 22 | 5 | 53/66 (80%) | 53 (100%) | 0 | 100 | 100 |
| 23 | 2 | 258/468 (55%) | 250 (97%) | 8 (3%) | 35 | 63 |
| 24 | X | 54/295 (18%) | 54 (100%) | 0 | 100 | 100 |
| 25 | U | 42/260 (16%) | 41 (98%) | 1 (2%) | 44 | 70 |
| 26 | V | 46/108 (43%) | 44 (96%) | 2 (4%) | 25 | 55 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|-------------|----|
| 29 | O | 152/205 (74%) | 144 (95%) | 8 (5%) | 19 | 48 |
| 30 | W | 155/429 (36%) | 147 (95%) | 8 (5%) | 19 | 48 |
| All | All | 6279/9989 (63%) | 5588 (89%) | 691 (11%) | 7 | 21 |

5 of 691 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 9 | E | 5 | ASN |
| 15 | L | 34 | CYS |
| 9 | E | 92 | THR |
| 9 | E | 4 | GLU |
| 11 | H | 143 | LEU |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 11 | H | 128 | ASN |
| 17 | 0 | 726 | GLN |
| 12 | I | 12 | ASN |
| 17 | 0 | 224 | ASN |
| 19 | 6 | 351 | ASN |

5.3.3 RNA ⓘ

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|-----------|-------------------|-----------------|
| 31 | P | 4/5 (80%) | 1 (25%) | 0 |

All (1) RNA backbone outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 31 | P | 8 | G |

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 16 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | $\# Z > 2$ | Counts | RMSZ | $\# Z > 2$ |
| 34 | SF4 | 0 | 801 | 17 | 0,12,12 | - | - | - | | |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|----------|---------|
| 34 | SF4 | 0 | 801 | 17 | - | - | 0/6/5/5 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 34 | 0 | 801 | SF4 | 2 | 0 |

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

| Mol | Chain | Number of breaks |
|-----|-------|------------------|
| 20 | 1 | 2 |

All chain breaks are listed below:

| Model | Chain | Residue-1 | Atom-1 | Residue-2 | Atom-2 | Distance (Å) |
|-------|-------|-----------|--------|-----------|--------|--------------|
| 1 | 1 | 355:UNK | C | 368:UNK | N | 14.81 |
| 1 | 1 | 519:UNK | C | 537:GLU | N | 11.86 |


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-23908. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 209



Y Index: 237




Z Index: 240


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 109



Y Index: 253




Z Index: 242


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)



The volume at the recommended contour level is 863 nm³; this corresponds to an approximate mass of 779 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-23908 and PDB model 7ML4. Per-residue inclusion information can be found in [section 3](#) on [page 12](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).

































































9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion | Q-score |
|-------|--|--|
| All |  0.7930 |  0.2810 |
| 0 |  0.6660 |  0.0610 |
| 1 |  0.5790 |  0.0820 |
| 2 |  0.6950 |  0.0880 |
| 3 |  0.4380 |  0.1210 |
| 4 |  0.7980 |  0.0710 |
| 5 |  0.4680 |  0.0950 |
| 6 |  0.7830 |  0.0790 |
| 7 |  0.4380 |  0.0470 |
| A |  0.9260 |  0.4730 |
| B |  0.9410 |  0.4960 |
| C |  0.9570 |  0.4930 |
| D |  0.4630 |  0.2750 |
| E |  0.9250 |  0.4610 |
| F |  0.9540 |  0.5000 |
| G |  0.6800 |  0.3600 |
| H |  0.9410 |  0.4440 |
| I |  0.8650 |  0.4200 |
| J |  0.9790 |  0.5080 |
| K |  0.8770 |  0.4750 |
| L |  0.9600 |  0.4430 |
| M |  0.7540 |  0.1890 |
| N |  0.9170 |  0.1710 |
| O |  0.8960 |  0.0690 |
| P |  0.6540 |  0.2610 |
| Q |  0.8950 |  0.3570 |
| R |  0.8490 |  0.2590 |
| T |  0.8460 |  0.1830 |
| U |  0.6490 |  0.0900 |
| V |  0.7880 |  0.1090 |
| W |  0.6690 |  0.1000 |
| X |  0.7940 |  0.0640 |

