



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 14, 2025 – 07:26 PM EDT

PDB ID : 3MOD / pdb\_00003mod  
Title : Crystal structure of the neutralizing HIV antibody 2F5 Fab fragment (recombinantly produced IgG) with 11 aa gp41 MPER-derived peptide  
Authors : Nicely, N.I.; Dennison, S.M.; Kelsoe, G.; Liao, H.-X.; Alam, S.M.; Haynes, B.F.  
Deposited on : 2010-04-22  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtrriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	12	<div> <div>8%</div> <div>100%</div> </div>
2	L	214	<div> <div>90%</div> <div>10%</div> </div>
3	H	237	<div> <div>4%</div> <div>85%</div> <div>13%</div> <div>.</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called gp41 MPER-derived peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	P	12	Total	C	N	O	0	0	1
			98	67	15	16			

- Molecule 2 is a protein called ANTI-HIV-1 ANTIBODY 2F5 LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1639	1023	281	330	5			

- Molecule 3 is a protein called ANTI-HIV-1 ANTIBODY 2F5 HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	237	Total	C	N	O	S	0	0	0
			1774	1124	301	342	7			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	4	Total	O	0	0
			4	4		
4	L	91	Total	O	0	0
			91	91		
4	H	120	Total	O	0	0
			120	120		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: gp41 MPER-derived peptide



- Molecule 2: ANTI-HIV-1 ANTIBODY 2F5 LIGHT CHAIN



- Molecule 3: ANTI-HIV-1 ANTIBODY 2F5 HEAVY CHAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.83Å 64.43Å 178.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.40 – 2.20 43.40 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.1 (43.40-2.20) 98.1 (43.40-2.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.80 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_271)	Depositor
R, $R_{free}$	0.184 , 0.214 0.176 , 0.204	Depositor DCC
$R_{free}$ test set	1987 reflections (5.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.6	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 39.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3726	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	P	1.69	0/100	0.69	0/136
2	L	0.52	0/1676	0.57	0/2279
3	H	1.57	7/1817 (0.4%)	0.69	0/2483
All	All	1.20	7/3593 (0.2%)	0.64	0/4898

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	22	CYS	CB-SG	-6.06	1.72	1.82
3	H	52	TYR	CD1-CE1	-5.98	1.30	1.39
3	H	92	CYS	CB-SG	-5.71	1.72	1.81
3	H	34	VAL	CB-CG2	-5.61	1.41	1.52
3	H	150	VAL	CB-CG1	-5.49	1.41	1.52
3	H	196	CYS	CB-SG	-5.25	1.73	1.81
3	H	59	TYR	CD2-CE2	-5.21	1.31	1.39

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	98	0	96	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	L	1639	0	1587	16	0
3	H	1774	0	1794	25	0
4	H	120	0	0	3	0
4	L	91	0	0	2	0
4	P	4	0	0	0	0
All	All	3726	0	3477	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:127:SER:O	3:H:131:THR:HG23	1.67	0.94
3:H:148:GLU:OE1	4:H:223:HOH:O	1.92	0.85
2:L:161:GLU:HG3	4:L:215:HOH:O	1.83	0.78
2:L:116:PHE:HD2	3:H:130:SER:HA	1.52	0.73
2:L:37:ARG:HD2	4:L:236:HOH:O	1.95	0.66
2:L:22:THR:HG22	2:L:72:THR:HG22	1.83	0.60
3:H:31:ASP:OD2	4:H:330:HOH:O	2.17	0.59
3:H:127:SER:O	3:H:131:THR:CG2	2.48	0.58
3:H:111:ILE:N	3:H:111:ILE:HD12	2.19	0.57
3:H:117:LYS:HD3	3:H:144:ASP:O	2.05	0.57
2:L:163:VAL:HG22	2:L:175:LEU:HD12	1.87	0.57
3:H:114:THR:CG2	3:H:115:SER:N	2.68	0.56
3:H:100(A):LEU:O	3:H:100(D):VAL:HG22	2.06	0.56
3:H:178:LEU:C	3:H:178:LEU:HD12	2.28	0.54
3:H:127:SER:OG	3:H:129:LYS:N	2.40	0.54
2:L:124:GLN:HG2	2:L:129:THR:O	2.10	0.52
3:H:114:THR:HG22	3:H:115:SER:N	2.24	0.51
2:L:116:PHE:HD2	3:H:130:SER:CA	2.21	0.51
2:L:169:LYS:N	2:L:169:LYS:HD2	2.26	0.49
2:L:12:SER:HB3	2:L:140:TYR:OH	2.14	0.48
3:H:114:THR:HG21	3:H:146:PHE:CG	2.50	0.47
3:H:178:LEU:HA	4:H:313:HOH:O	2.14	0.47
3:H:123:PRO:HD3	3:H:209:LYS:HE2	1.97	0.46
3:H:143:LYS:HA	3:H:177:SER:HB2	1.98	0.46
2:L:39:LYS:HE2	2:L:81:GLU:O	2.16	0.45
2:L:116:PHE:CD2	3:H:130:SER:HA	2.41	0.45
2:L:186:TYR:CZ	2:L:211:ARG:HD2	2.52	0.45
3:H:132:SER:N	3:H:135:THR:O	2.44	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:100(H):ARG:O	3:H:100(I):GLY:C	2.55	0.44
3:H:6:GLU:HG3	3:H:92:CYS:SG	2.58	0.43
3:H:100(H):ARG:O	3:H:100(K):VAL:HG22	2.19	0.42
2:L:12:SER:CB	2:L:140:TYR:OH	2.68	0.42
2:L:12:SER:HB3	2:L:105:ASP:OD2	2.19	0.41
3:H:148:GLU:HA	3:H:149:PRO:HA	1.78	0.41
3:H:22:CYS:HB2	3:H:36:TRP:CZ2	2.56	0.41
2:L:32:ALA:HB3	2:L:92:HIS:HB2	2.02	0.41
3:H:100(D):VAL:HA	3:H:100(E):PRO:HD3	1.80	0.41
2:L:11:LEU:C	2:L:11:LEU:HD12	2.40	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	10/12 (83%)	10 (100%)	0	0	100	100
2	L	212/214 (99%)	208 (98%)	4 (2%)	0	100	100
3	H	235/237 (99%)	227 (97%)	6 (3%)	2 (1%)	14	14
All	All	457/463 (99%)	445 (97%)	10 (2%)	2 (0%)	30	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	H	100(I)	GLY
3	H	149	PRO



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	10/10 (100%)	10 (100%)	0	100	100
2	L	186/186 (100%)	183 (98%)	3 (2%)	58	73
3	H	207/207 (100%)	204 (99%)	3 (1%)	62	77
All	All	403/403 (100%)	397 (98%)	6 (2%)	60	75

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	2	LEU
2	L	12	SER
2	L	18	ARG
3	H	150	VAL
3	H	172	SER
3	H	177	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	P	11/12 (91%)	-0.26	1 (9%) 16 14	28, 35, 52, 68	0
2	L	214/214 (100%)	-0.36	1 (0%) 87 85	23, 39, 60, 77	0
3	H	237/237 (100%)	-0.24	10 (4%) 41 37	20, 36, 81, 130	0
All	All	462/463 (99%)	-0.30	12 (2%) 57 54	20, 37, 65, 130	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	129	LYS	5.1
3	H	131	THR	5.1
3	H	130	SER	4.4
3	H	128	SER	4.1
3	H	133	GLY	3.9
2	L	1	ALA	3.6
3	H	216	CYS	3.1
3	H	132	SER	3.1
3	H	215	SER	2.9
1	P	660	LEU	2.5
3	H	100(D)	VAL	2.3
3	H	100(B)	PHE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.