



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 01:47 AM EDT

PDB ID : 5MTW
Title : Mycobacterium tuberculosis Rv1957 SecB-like chaperone in complex with a ChAD peptide from Rv1956 HigA1 antitoxin
Authors : Guillet, V.; Mourey, L.
Deposited on : 2017-01-10
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	<div> <div>4%</div> <div>70%</div> <div>5%</div> <div>25%</div> </div>
1	B	184	<div> <div>6%</div> <div>66%</div> <div>9%</div> <div>25%</div> </div>
1	C	184	<div> <div>3%</div> <div>65%</div> <div>9%</div> <div>25%</div> </div>
1	D	184	<div> <div>4%</div> <div>61%</div> <div>12%</div> <div>26%</div> </div>
2	E	13	<div> <div>15%</div> <div>62%</div> <div>31%</div> <div>8%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	13	<div><div></div><div>8%</div><div>85%</div><div>8%</div><div>8%</div></div>
2	G	13	<div><div></div><div>8%</div><div>62%</div><div>31%</div><div>8%</div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SecB-like chaperone Rv1957.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	138	Total	C	N	O	S	0	0	0
			1060	676	180	203	1			
1	D	136	Total	C	N	O	S	0	2	0
			1069	683	184	201	1			
1	A	138	Total	C	N	O	S	0	0	0
			1072	682	184	205	1			
1	B	138	Total	C	N	O	S	0	0	0
			1077	686	186	204	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP P95257
C	-1	SER	-	expression tag	UNP P95257
C	0	HIS	-	expression tag	UNP P95257
D	-2	GLY	-	expression tag	UNP P95257
D	-1	SER	-	expression tag	UNP P95257
D	0	HIS	-	expression tag	UNP P95257
A	-2	GLY	-	expression tag	UNP P95257
A	-1	SER	-	expression tag	UNP P95257
A	0	HIS	-	expression tag	UNP P95257
B	-2	GLY	-	expression tag	UNP P95257
B	-1	SER	-	expression tag	UNP P95257
B	0	HIS	-	expression tag	UNP P95257

- Molecule 2 is a protein called Antitoxin HigA1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	12	Total	C	N	O	0	0	0
			108	69	21	18			
2	F	12	Total	C	N	O	0	0	0
			108	69	21	18			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	12	Total	C	N	O	0	0	0
			102	66	18	18			

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	13	Total	O	0	0
			13	13		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	16	Total 16	O 16	0	0
5	A	11	Total 11	O 11	0	0
5	B	4	Total 4	O 4	0	0
5	E	1	Total 1	O 1	0	0
5	G	4	Total 4	O 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SecB-like chaperone Rv1957




- Molecule 1: SecB-like chaperone Rv1957




- Molecule 1: SecB-like chaperone Rv1957




- Molecule 1: SecB-like chaperone Rv1957








● Molecule 2: Antitoxin HigA1



● Molecule 2: Antitoxin HigA1



● Molecule 2: Antitoxin HigA1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.86Å 89.96Å 91.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.09 – 1.84 62.94 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.9 (64.09-1.84) 99.9 (62.94-1.84)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.212 , 0.265 0.217 , 0.267	Depositor DCC
R_{free} test set	3397 reflections (5.38%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.024 for -h,l,k 0.027 for -l,-k,-h 0.025 for k,h,-l 0.009 for k,l,h 0.009 for l,h,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4656	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	0/1092	0.97	2/1489 (0.1%)
1	B	0.98	0/1097	1.00	3/1496 (0.2%)
1	C	1.02	1/1081 (0.1%)	1.11	9/1477 (0.6%)
1	D	0.98	2/1097 (0.2%)	1.02	3/1496 (0.2%)
2	E	0.98	0/112	0.95	1/152 (0.7%)
2	F	0.78	0/112	0.95	1/152 (0.7%)
2	G	1.20	0/106	0.91	0/145
All	All	0.98	3/4697 (0.1%)	1.02	19/6407 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	160	GLU	CD-OE1	7.21	1.33	1.25
1	D	72	SER	CB-OG	-7.16	1.32	1.42
1	D	143	GLU	CD-OE2	5.64	1.31	1.25

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	32	ARG	NE-CZ-NH1	8.88	124.74	120.30
1	C	60	ASP	CB-CG-OD1	7.95	125.46	118.30
1	D	32	ARG	NE-CZ-NH1	7.40	124.00	120.30
1	C	27	ASP	CB-CG-OD1	7.19	124.77	118.30
1	C	60	ASP	CB-CG-OD2	-7.03	111.97	118.30
1	D	151	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	A	151	ARG	NE-CZ-NH2	-6.34	117.13	120.30
1	B	40	ALA	C-N-CD	6.21	141.45	128.40
1	C	47	LEU	CB-CG-CD2	-5.88	101.00	111.00
2	F	110	ARG	NE-CZ-NH1	5.81	123.20	120.30
2	E	110	ARG	NE-CZ-NH1	5.50	123.05	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	118	ASP	CB-CG-OD2	-5.44	113.41	118.30
1	C	27	ASP	CB-CG-OD2	-5.34	113.49	118.30
1	B	18	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	C	102	ASP	CB-CG-OD1	5.19	122.97	118.30
1	C	18	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	D	18	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	B	32	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	C	47	LEU	CB-CA-C	5.01	119.72	110.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1072	0	1029	8	0
1	B	1077	0	1040	8	0
1	C	1060	0	1015	13	0
1	D	1069	0	1040	13	0
2	E	108	0	101	4	0
2	F	108	0	101	0	0
2	G	102	0	90	6	1
3	B	4	0	6	0	0
3	C	4	0	6	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	11	0	0	1	1
5	B	4	0	0	0	0
5	C	13	0	0	0	0
5	D	16	0	0	1	0
5	E	1	0	0	0	0
5	G	4	0	0	0	0
All	All	4656	0	4428	41	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:47:LEU:HD21	2:E:114:TYR:CD2	2.06	0.90
1:A:19:LEU:HD11	1:A:120:PRO:HB3	1.58	0.84
1:A:26:ARG:NE	1:A:110:ASP:OD2	2.14	0.81
1:C:47:LEU:HD21	2:E:114:TYR:CE2	2.19	0.77
1:D:44:ALA:HB2	2:G:111:LEU:HD22	1.67	0.77
1:D:147:ASP:OD1	5:D:301:HOH:O	2.07	0.71
1:B:114:GLN:HE21	1:B:114:GLN:HA	1.56	0.70
1:D:161:ILE:H	1:B:24:GLN:HE22	1.40	0.67
1:A:26:ARG:HG3	1:A:110:ASP:OD2	1.95	0.66
1:C:18:ARG:NH2	1:C:118:ASP:O	2.27	0.65
1:C:47:LEU:CD2	2:E:114:TYR:CD2	2.78	0.64
1:B:24:GLN:HE21	1:B:25:ILE:H	1.45	0.63
1:B:50:ASP:OD2	1:B:76:ARG:NH2	2.33	0.61
1:C:28:ILE:HD13	1:C:140:TYR:CE1	2.35	0.60
1:A:26:ARG:CG	1:A:110:ASP:OD2	2.50	0.60
1:A:116:GLY:O	5:A:301:HOH:O	2.17	0.59
1:B:58:ASP:OD2	1:B:70:ARG:NH1	2.37	0.56
1:D:44:ALA:CB	2:G:111:LEU:HD22	2.34	0.56
1:C:47:LEU:C	1:C:47:LEU:HD23	2.26	0.56
1:C:71:ILE:HD11	1:C:134:ARG:HA	1.89	0.53
1:C:28:ILE:HD13	1:C:140:TYR:CZ	2.45	0.51
1:C:47:LEU:HD21	2:E:114:TYR:CG	2.44	0.50
1:D:22:ARG:NE	1:D:117:GLU:OE2	2.45	0.50
1:D:43:PRO:HB3	2:G:111:LEU:N	2.27	0.50
1:D:53:PHE:N	1:D:53:PHE:CD1	2.81	0.47
1:B:24:GLN:HE21	1:B:25:ILE:N	2.13	0.47
1:C:113:LEU:C	1:C:117:GLU:HG2	2.35	0.46
1:D:55:PRO:HA	1:D:71:ILE:HD13	1.97	0.46
1:D:118:ASP:N	1:D:118:ASP:OD1	2.47	0.46
1:A:147:ASP:O	1:A:151:ARG:HG3	2.16	0.45
1:B:28:ILE:HD13	1:B:140:TYR:CE1	2.51	0.45
1:D:76:ARG:NH1	1:D:97:ASP:OD2	2.51	0.43
1:D:44:ALA:HB3	2:G:111:LEU:HD13	2.01	0.42
1:C:117:GLU:HG3	1:C:118:ASP:N	2.33	0.42
1:D:74:HIS:ND1	1:D:102:ASP:OD1	2.38	0.42
1:D:41:PRO:HG3	1:D:154:LEU:HD23	2.02	0.42
1:A:26:ARG:CD	1:A:110:ASP:OD2	2.68	0.41
1:B:42:LYS:H	1:B:79:ASN:HD21	1.68	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:138:TYR:CZ	1:C:162:LEU:HG	2.56	0.41
1:A:47:LEU:HD12	2:G:112:SER:OG	2.21	0.41
1:C:64:ILE:HD11	2:G:105:VAL:HB	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:104:GLU:OE2	5:A:301:HOH:O[2_575]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/184 (72%)	129 (98%)	3 (2%)	0	100	100
1	B	132/184 (72%)	127 (96%)	4 (3%)	1 (1%)	19	7
1	C	132/184 (72%)	129 (98%)	3 (2%)	0	100	100
1	D	132/184 (72%)	130 (98%)	2 (2%)	0	100	100
2	E	10/13 (77%)	10 (100%)	0	0	100	100
2	F	10/13 (77%)	10 (100%)	0	0	100	100
2	G	10/13 (77%)	10 (100%)	0	0	100	100
All	All	558/775 (72%)	545 (98%)	12 (2%)	1 (0%)	47	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	114	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/149 (71%)	105 (99%)	1 (1%)	78	71
1	B	107/149 (72%)	103 (96%)	4 (4%)	34	16
1	C	104/149 (70%)	100 (96%)	4 (4%)	33	15
1	D	107/149 (72%)	102 (95%)	5 (5%)	26	10
2	E	12/12 (100%)	10 (83%)	2 (17%)	2	0
2	F	12/12 (100%)	12 (100%)	0	100	100
2	G	11/12 (92%)	11 (100%)	0	100	100
All	All	459/632 (73%)	443 (96%)	16 (4%)	36	18

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	47	LEU
1	C	63	THR
1	C	117	GLU
1	C	163	SER
1	D	45	GLN
1	D	53	PHE
1	D	113	LEU
1	D	118	ASP
1	D	122	GLU
1	A	11	ASP
1	B	10	LEU
1	B	63	THR
1	B	114	GLN
1	B	162	LEU
2	E	111	LEU
2	E	115	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	B	24	GLN
1	B	79	ASN
1	B	114	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DMS	B	201	-	3,3,3	0.59	0	3,3,3	0.94	0
3	DMS	C	201	-	3,3,3	0.70	0	3,3,3	0.61	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/184 (75%)	0.20	8 (5%) 23 21	19, 31, 62, 72	7 (5%)
1	B	138/184 (75%)	0.37	11 (7%) 12 11	18, 35, 59, 69	12 (8%)
1	C	138/184 (75%)	0.32	5 (3%) 42 39	15, 31, 62, 75	7 (5%)
1	D	136/184 (73%)	0.29	7 (5%) 28 25	17, 32, 56, 69	14 (10%)
2	E	12/13 (92%)	0.68	2 (16%) 1 1	33, 38, 52, 70	2 (16%)
2	F	12/13 (92%)	0.26	1 (8%) 11 10	35, 41, 53, 65	1 (8%)
2	G	12/13 (92%)	0.32	1 (8%) 11 10	21, 31, 46, 50	3 (25%)
All	All	586/775 (75%)	0.30	35 (5%) 21 19	15, 33, 60, 75	46 (7%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	13	GLN	7.5
1	D	80	GLN	7.2
1	A	9	ASP	6.0
1	C	166	MET	5.2
1	D	79	ASN	5.1
1	D	116	GLY	4.5
1	B	7	ALA	4.0
2	E	115	ARG	3.8
1	A	81	ALA	3.6
1	B	6	ASP	3.5
1	B	115	GLU	3.5
1	A	10	LEU	3.4
1	D	14	ARG	3.3
1	C	118	ASP	3.3
1	A	163	SER	3.3
1	B	112	HIS	3.3
1	B	130	ALA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	44	ALA	3.2
1	B	10	LEU	3.0
1	B	62	ALA	3.0
1	D	10	LEU	2.9
1	C	46	GLY	2.7
1	A	46	GLY	2.6
1	D	40	ALA	2.5
1	A	53	PHE	2.5
1	A	80	GLN	2.5
1	A	11	ASP	2.5
1	B	42	LYS	2.5
2	G	115	ARG	2.4
2	E	104	GLU	2.3
1	B	63	THR	2.3
2	F	115	ARG	2.3
1	B	9	ASP	2.2
1	D	47	LEU	2.1
1	B	114	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CA	A	201	1/1	0.92	0.23	41,41,41,41	1
4	CA	D	201	1/1	0.95	0.14	37,37,37,37	0
4	CA	B	202	1/1	0.95	0.10	58,58,58,58	0
3	DMS	B	201	4/4	0.96	0.13	57,60,65,66	0
3	DMS	C	201	4/4	0.96	0.15	58,64,65,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.