



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2024 – 05:51 PM EDT

PDB ID : 6N29
Title : Crystal structure of monomeric von Willebrand Factor D'D3 assembly
Authors : Dong, X.; Arndt, J.W.; Springer, T.A.
Deposited on : 2018-11-12
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	 5% 90% 9% .
1	B	483	 5% 88% 11% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called von Willebrand factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	478	Total	C	N	O	S	0	1	0
			3645	2246	633	708	58			
1	B	475	Total	C	N	O	S	0	1	0
			3628	2236	630	704	58			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	852	ARG	GLN	conflict	UNP P04275
A	1099	ALA	CYS	conflict	UNP P04275
A	1142	ALA	CYS	conflict	UNP P04275
A	1245	PRO	-	expression tag	UNP P04275
A	1246	ARG	-	expression tag	UNP P04275
B	852	ARG	GLN	conflict	UNP P04275
B	1099	ALA	CYS	conflict	UNP P04275
B	1142	ALA	CYS	conflict	UNP P04275
B	1245	PRO	-	expression tag	UNP P04275
B	1246	ARG	-	expression tag	UNP P04275

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		
3	B	1	Total	Ca	0	0
			1	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	47	Total	O	0	0
			47	47		
5	B	53	Total	O	0	0
			53	53		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: von Willebrand factor



- Molecule 1: von Willebrand factor



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	174.69Å 174.69Å 104.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.45 – 2.50 48.45 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.45-2.50) 98.6 (48.45-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.202 , 0.241 0.208 , 0.245	Depositor DCC
R_{free} test set	53525 reflections (3.78%)	wwPDB-VP
Wilson B-factor (Å ²)	78.3	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 88.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7460	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/3724	0.50	0/5061
1	B	0.31	0/3706	0.50	0/5036
All	All	0.31	0/7430	0.50	0/10097

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3645	0	3444	25	0
1	B	3628	0	3430	30	0
2	A	42	0	39	0	0
2	B	28	0	26	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	7	0	10	0	0
4	B	7	0	10	0	0
5	A	47	0	0	1	0
5	B	53	0	0	4	0
All	All	7460	0	6959	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1122:THR:HG23	1:B:1125:LEU:H	1.48	0.77
1:A:791:THR:HG23	1:A:793:GLN:H	1.56	0.70
1:A:1122:THR:HG23	1:A:1125:LEU:H	1.57	0.69
1:B:791:THR:HG23	1:B:793:GLN:H	1.58	0.67
1:A:965:LEU:HD21	1:A:1166:PRO:HD3	1.80	0.63
1:B:1225:CYS:HB3	1:B:1232:LEU:HD11	1.83	0.60
1:A:945:ARG:NH1	1:B:946:PRO:O	2.27	0.60
1:B:921:CYS:SG	1:B:922:LYS:N	2.78	0.57
1:B:787:GLU:HB2	5:B:1415:HOH:O	2.04	0.57
1:A:1204:ARG:NH1	5:A:1403:HOH:O	2.37	0.56
1:B:939:GLY:O	1:B:976:ARG:NH1	2.38	0.56
1:B:952:HIS:CE1	1:B:968:LYS:HE3	2.41	0.56
1:A:1052:LYS:NZ	1:A:1091:CYS:O	2.38	0.54
1:B:1206:PHE:CD1	1:B:1212:VAL:HG21	2.43	0.53
1:A:1123:ALA:HA	1:A:1127:PRO:HB3	1.90	0.53
1:A:925:VAL:HB	1:A:936:LEU:HB2	1.92	0.52
1:A:777:PRO:HG2	1:A:780:ASN:HB2	1.92	0.52
1:B:985:LYS:NZ	5:B:1403:HOH:O	2.44	0.51
1:B:953:PHE:HA	1:B:965:LEU:O	2.10	0.51
1:B:796:ASP:OD1	1:B:796:ASP:N	2.37	0.51
1:B:925:VAL:HB	1:B:936:LEU:HB2	1.94	0.50
1:B:965:LEU:HD21	1:B:1166:PRO:HD3	1.91	0.50
1:B:969:ALA:HB1	1:B:985:LYS:HB2	1.93	0.50
1:A:1046:CYS:HB3	1:A:1053[B]:GLN:HG2	1.94	0.49
1:A:796:ASP:N	1:A:796:ASP:OD1	2.45	0.48
1:B:791:THR:HG22	1:B:794:ASN:OD1	2.14	0.47
1:A:965:LEU:CD2	1:A:1166:PRO:HD3	2.44	0.47
1:B:1052:LYS:NZ	1:B:1091:CYS:O	2.32	0.47
1:A:870:ILE:HD11	1:A:876:LEU:HD22	1.97	0.46
1:B:786:LEU:HD11	1:B:807:GLY:HA3	1.98	0.46
1:A:953:PHE:HA	1:A:965:LEU:O	2.16	0.46
1:B:953:PHE:O	1:B:1210:LYS:NZ	2.49	0.46
1:A:791:THR:HG22	1:A:794:ASN:OD1	2.17	0.45
1:B:881:LEU:HB2	1:B:1003:GLN:HG3	1.99	0.45
1:A:921:CYS:SG	1:A:922:LYS:N	2.89	0.44
1:B:894:VAL:HG11	1:B:982:VAL:HG11	1.98	0.44
1:B:1156:THR:HA	1:B:1182:ILE:HG23	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1150:ALA:HB3	1:A:1174:HIS:CE1	2.53	0.44
1:A:1156:THR:HA	1:A:1182:ILE:HG23	1.99	0.44
1:B:1018:PRO:HD2	5:B:1431:HOH:O	2.18	0.43
1:A:1112:ALA:HB1	1:A:1149:CYS:SG	2.59	0.43
1:A:881:LEU:HB2	1:A:1003:GLN:HG3	2.01	0.43
1:B:855:LYS:HG3	5:B:1410:HOH:O	2.18	0.42
1:B:1100:PHE:HE2	1:B:1125:LEU:HD11	1.84	0.42
1:A:782:ARG:HD3	1:A:782:ARG:HA	1.84	0.42
1:B:927:ILE:HG21	1:B:966:LEU:HD13	2.01	0.42
1:A:826:ARG:HD3	1:A:826:ARG:HA	1.81	0.42
1:B:881:LEU:HD21	1:B:1007:LEU:HG	2.01	0.41
1:B:794:ASN:HA	1:B:797:LEU:HD12	2.02	0.41
1:A:1133:ARG:NH2	1:A:1173:CYS:HB2	2.35	0.41
1:A:894:VAL:HG11	1:A:982:VAL:HG11	2.03	0.41
1:A:1208:SER:OG	1:A:1228:ASP:O	2.35	0.41
1:B:785:GLY:N	1:B:804:CYS:HB2	2.36	0.40
1:B:1199:CYS:HB3	1:B:1232:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	477/483 (99%)	455 (95%)	21 (4%)	1 (0%)	44	64
1	B	474/483 (98%)	453 (96%)	20 (4%)	1 (0%)	44	64
All	All	951/966 (98%)	908 (96%)	41 (4%)	2 (0%)	44	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1044	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1044	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	420/427 (98%)	417 (99%)	3 (1%)	81	93
1	B	418/427 (98%)	414 (99%)	4 (1%)	73	88
All	All	838/854 (98%)	831 (99%)	7 (1%)	79	91

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1031	CYS
1	A	1149	CYS
1	A	1165	CYS
1	B	1031	CYS
1	B	1149	CYS
1	B	1165	CYS
1	B	1181	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1302	1	14,14,15	0.41	0	17,19,21	0.51	0
2	NAG	B	1302	1	14,14,15	0.31	0	17,19,21	0.40	0
2	NAG	B	1301	1	14,14,15	0.29	0	17,19,21	0.47	0
4	PEG	A	1306	-	6,6,6	0.51	0	5,5,5	0.29	0
2	NAG	A	1303	1	14,14,15	0.31	0	17,19,21	0.49	0
4	PEG	B	1304	-	6,6,6	0.50	0	5,5,5	0.27	0
2	NAG	A	1301	1	14,14,15	0.32	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1302	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1302	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1301	1	-	2/6/23/26	0/1/1/1
4	PEG	A	1306	-	-	1/4/4/4	-
2	NAG	A	1303	1	-	1/6/23/26	0/1/1/1
4	PEG	B	1304	-	-	1/4/4/4	-
2	NAG	A	1301	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1301	NAG	O5-C5-C6-O6
2	B	1301	NAG	C4-C5-C6-O6
2	A	1303	NAG	O5-C5-C6-O6
4	B	1304	PEG	O1-C1-C2-O2
4	A	1306	PEG	C1-C2-O2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	478/483 (98%)	0.37	24 (5%) 35 32	47, 93, 172, 293	1 (0%)
1	B	475/483 (98%)	0.36	23 (4%) 36 34	49, 94, 163, 249	1 (0%)
All	All	953/966 (98%)	0.36	47 (4%) 36 33	47, 94, 170, 293	2 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	902	PRO	7.5
1	A	781	LEU	5.1
1	B	765	LEU	5.0
1	B	918	SER	4.8
1	B	781	LEU	4.6
1	A	775	VAL	4.4
1	A	919	VAL	4.4
1	B	919	VAL	4.1
1	A	783	ALA	4.1
1	A	918	SER	3.6
1	A	902	PRO	3.5
1	A	853	ASP	3.5
1	A	917	PRO	3.5
1	A	914	CYS	3.2
1	B	1212	VAL	2.9
1	A	765	LEU	2.8
1	A	774	LEU	2.8
1	B	917	PRO	2.8
1	B	1232	LEU	2.6
1	A	920	LYS	2.6
1	A	1031	CYS	2.6
1	A	777	PRO	2.5
1	A	795	TYR	2.5
1	B	848	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	775	VAL	2.5
1	A	1016	GLU	2.5
1	B	1206	PHE	2.5
1	A	796	ASP	2.4
1	A	773	LYS	2.3
1	A	799	CYS	2.3
1	A	764	SER	2.3
1	B	801	SER	2.3
1	B	914	CYS	2.3
1	A	809	LEU	2.3
1	A	1145	ARG	2.3
1	B	800	MET	2.3
1	B	916	HIS	2.2
1	B	783	ALA	2.2
1	B	791	THR	2.1
1	B	802	MET	2.1
1	A	854	ARG	2.1
1	B	1014	VAL	2.1
1	A	1030	GLN	2.1
1	B	1031	CYS	2.1
1	B	782	ARG	2.0
1	B	788	CYS	2.0
1	B	787	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	A	1303	14/15	0.44	0.15	137,163,181,184	0
4	PEG	B	1304	7/7	0.51	0.17	118,137,143,146	0
2	NAG	A	1301	14/15	0.66	0.12	137,157,164,167	0
2	NAG	B	1301	14/15	0.73	0.12	132,145,167,171	0
2	NAG	B	1302	14/15	0.75	0.13	106,128,140,148	0
4	PEG	A	1306	7/7	0.84	0.18	115,119,130,134	0
2	NAG	A	1302	14/15	0.84	0.11	105,116,127,129	0
3	CA	A	1305	1/1	0.94	0.05	144,144,144,144	0
3	CA	A	1304	1/1	0.99	0.04	79,79,79,79	0
3	CA	B	1303	1/1	0.99	0.10	101,101,101,101	0

6.5 Other polymers [i](#)

There are no such residues in this entry.