



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 05:50 PM EDT

PDB ID : 5NBW
Title : Crystal structure of the Fab fragment 22F12 in complex with 3-hydroxybenzo[a]pyrene
Authors : Skerra, A.; Eichinger, A.
Deposited on : 2017-03-02
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 89%, yellow 89%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 89% 11% </div> </div>
1	L	214	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 93%, yellow 93%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 93% 7% </div> </div>
2	B	222	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 5%, green 5%, green 85%, yellow 85%, yellow 11%, grey 11%, grey 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 5% 85% 11% .. </div> </div>
2	H	222	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 4%, green 4%, green 87%, yellow 87%, yellow 9%, grey 9%, grey 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 4% 87% 9% .. </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6920 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fab 22F12 (L,H).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	214	Total	C	N	O	S	0	0	0
			1682	1053	277	344	8			
1	A	214	Total	C	N	O	S	0	0	0
			1682	1053	277	344	8			

- Molecule 2 is a protein called Fab 22F12 (A,B).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	216	Total	C	N	O	S	0	0	0
			1636	1037	268	324	7			
2	B	216	Total	C	N	O	S	0	0	0
			1636	1037	268	324	7			

- Molecule 3 is benzo[a]pyren-3-ol (three-letter code: 8SK) (formula: C₂₀H₁₂O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	C	O	0	0
			21	20	1		
3	A	1	Total	C	O	0	0
			21	20	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	61	Total	O	0	0
			61	61		
4	H	87	Total	O	0	0
			87	87		
4	A	53	Total	O	0	0
			53	53		
4	B	41	Total	O	0	0
			41	41		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab 22F12 (L,H)




- Molecule 1: Fab 22F12 (L,H)



- Molecule 2: Fab 22F12 (A,B)



- Molecule 2: Fab 22F12 (A,B)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.15Å 81.00Å 186.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.10 – 2.40 61.10 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (61.10-2.40) 99.7 (61.10-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.40Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.180 , 0.228 0.185 , 0.230	Depositor DCC
R_{free} test set	1704 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	0.447	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6920	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 8SK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.96	2/1721 (0.1%)	1.00	7/2333 (0.3%)
1	L	0.93	0/1721	0.96	4/2333 (0.2%)
2	B	0.94	2/1682 (0.1%)	0.94	2/2302 (0.1%)
2	H	0.96	0/1682	0.97	4/2302 (0.2%)
All	All	0.95	4/6806 (0.1%)	0.97	17/9270 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	204	SER	CB-OG	6.38	1.50	1.42
2	B	149	GLU	CD-OE1	6.28	1.32	1.25
1	A	191	SER	CB-OG	-5.79	1.34	1.42
1	A	194	CYS	CB-SG	-5.01	1.73	1.81

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	208	ASP	CB-CG-OD1	7.37	124.93	118.30
2	H	57	ARG	NE-CZ-NH1	6.90	123.75	120.30
1	A	18	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	A	188	ARG	NE-CZ-NH2	-6.64	116.98	120.30
1	L	11	MET	CB-CA-C	-6.42	97.56	110.40
1	A	70	ASP	CB-CG-OD1	6.17	123.86	118.30
2	H	214	ARG	NE-CZ-NH2	-5.98	117.31	120.30
2	H	89	ASP	CB-CG-OD2	5.71	123.43	118.30
2	H	178	LEU	CA-CB-CG	5.66	128.32	115.30
1	A	188	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	155	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	L	83	MET	CG-SD-CE	5.28	108.64	100.20
1	A	70	ASP	CB-CG-OD2	-5.25	113.57	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	184	ASP	CB-CG-OD1	5.15	122.93	118.30
2	B	57	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	A	108	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	L	188	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1682	0	1601	8	0
1	L	1682	0	1601	5	0
2	B	1636	0	1586	6	0
2	H	1636	0	1586	12	0
3	A	21	0	0	0	0
3	L	21	0	0	0	0
4	A	53	0	0	1	0
4	B	41	0	0	0	0
4	H	87	0	0	1	0
4	L	61	0	0	0	0
All	All	6920	0	6374	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:173:SER:OG	2:H:174:ASP:N	2.24	0.70
2:H:12:VAL:HG11	2:H:86:LEU:HD13	1.80	0.64
1:L:214:CYS:HB3	2:H:216:CYS:HA	1.80	0.62
2:B:173:SER:OG	2:B:174:ASP:N	2.35	0.56
1:A:106:LEU:HD23	1:A:171:SER:OG	2.12	0.49
1:L:136:LEU:HD12	1:L:136:LEU:N	2.27	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:GLU:OE2	4:A:401:HOH:O	2.20	0.47
1:A:136:LEU:HD12	1:A:136:LEU:N	2.29	0.47
1:A:83:MET:CE	1:A:106:LEU:HD13	2.45	0.47
1:L:192:TYR:O	1:L:208:SER:HB2	2.16	0.46
2:H:87:THR:HG21	2:B:25:PRO:HB2	1.99	0.45
2:H:40:ARG:HD2	4:H:309:HOH:O	2.16	0.45
2:H:178:LEU:C	2:H:178:LEU:HD23	2.37	0.44
2:B:215:ASP:O	2:B:216:CYS:HB3	2.17	0.44
1:A:91:TYR:HA	1:A:96:TYR:CD1	2.53	0.44
1:L:214:CYS:CB	2:H:216:CYS:HA	2.46	0.44
1:A:192:TYR:O	1:A:208:SER:HB2	2.17	0.44
1:L:91:TYR:HA	1:L:96:TYR:CD1	2.54	0.43
2:H:189:TRP:CG	2:H:190:PRO:HA	2.53	0.43
2:B:178:LEU:C	2:B:178:LEU:HD23	2.39	0.43
1:A:83:MET:HE1	1:A:106:LEU:HD13	2.01	0.42
2:H:133:THR:HG23	2:H:133:THR:O	2.20	0.42
2:B:42:GLY:O	2:B:44:GLY:N	2.49	0.41
2:B:27:TYR:O	2:B:28:THR:OG1	2.29	0.41
2:H:133:THR:O	2:H:133:THR:CG2	2.68	0.41
2:H:189:TRP:CD1	2:H:190:PRO:HA	2.55	0.41
2:H:12:VAL:CG1	2:H:86:LEU:HD13	2.49	0.41
1:A:149:LYS:HB2	1:A:193:THR:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
1	L	212/214 (99%)	208 (98%)	4 (2%)	0	100	100
2	B	214/222 (96%)	204 (95%)	4 (2%)	6 (3%)	5	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	214/222 (96%)	205 (96%)	7 (3%)	2 (1%)	17	25
All	All	852/872 (98%)	824 (97%)	20 (2%)	8 (1%)	17	25

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	132	GLN
2	H	130	ALA
2	B	43	HIS
2	B	128	GLY
2	B	100	TYR
2	B	130	ALA
2	H	100	TYR
2	B	25	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/193 (100%)	187 (97%)	6 (3%)	40	60
1	L	193/193 (100%)	189 (98%)	4 (2%)	53	72
2	B	185/191 (97%)	175 (95%)	10 (5%)	22	36
2	H	185/191 (97%)	176 (95%)	9 (5%)	25	40
All	All	756/768 (98%)	727 (96%)	29 (4%)	33	51

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	56	ASP
1	L	169	LYS
1	L	185	GLU
1	L	190	ASN
2	H	3	LYS
2	H	18	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	133	THR
2	H	161	SER
2	H	178	LEU
2	H	196	CYS
2	H	206	LYS
2	H	215	ASP
2	H	216	CYS
1	A	24	LYS
1	A	56	ASP
1	A	116	SER
1	A	169	LYS
1	A	185	GLU
1	A	202	THR
2	B	1	GLU
2	B	3	LYS
2	B	18	VAL
2	B	65	LYS
2	B	132	GLN
2	B	133	THR
2	B	161	SER
2	B	172	GLN
2	B	178	LEU
2	B	196	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	69	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	8SK	A	301	-	25,25,25	1.06	2 (8%)	38,38,38	1.24	2 (5%)
3	8SK	L	301	-	25,25,25	0.93	1 (4%)	38,38,38	1.11	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	8SK	A	301	-	-	-	0/5/5/5
3	8SK	L	301	-	-	-	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	8SK	CAN-CAM	2.17	1.41	1.35
3	L	301	8SK	CAO-CAK	2.10	1.40	1.35
3	A	301	8SK	CAP-CAQ	2.01	1.40	1.36

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	8SK	OAA-CAL-CAF	4.44	121.86	116.31
3	L	301	8SK	OAA-CAL-CAF	3.70	120.93	116.31
3	L	301	8SK	CAL-CAF-CAC	3.23	119.99	118.56
3	A	301	8SK	CAL-CAF-CAC	3.11	119.94	118.56
3	L	301	8SK	CAS-CAI-CAJ	-2.05	118.65	122.00


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/214 (100%)	-0.35	2 (0%) 84 82	21, 33, 55, 113	0
1	L	214/214 (100%)	-0.36	3 (1%) 75 73	24, 33, 54, 121	0
2	B	216/222 (97%)	-0.18	10 (4%) 32 31	21, 37, 76, 118	0
2	H	216/222 (97%)	-0.30	9 (4%) 36 35	22, 29, 72, 113	0
All	All	860/872 (98%)	-0.30	24 (2%) 53 51	21, 33, 66, 121	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	27	TYR	8.8
1	A	214	CYS	8.0
2	H	131	ALA	7.9
1	L	214	CYS	6.8
2	B	216	CYS	5.9
2	H	130	ALA	5.8
2	H	132	GLN	5.7
1	L	213	GLU	3.6
2	B	133	THR	3.6
2	B	1	GLU	3.2
1	L	212	ASN	3.1
2	B	132	GLN	3.1
2	B	129	SER	3.1
2	H	128	GLY	3.0
2	H	27	TYR	2.9
2	H	129	SER	2.9
2	H	133	THR	2.8
2	B	134	ASN	2.8
2	B	26	GLY	2.8
2	H	1	GLU	2.5
2	B	187	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	131	ALA	2.3
2	H	216	CYS	2.3
1	A	213	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	8SK	L	301	21/21	0.95	0.12	25,26,28,28	0
3	8SK	A	301	21/21	0.96	0.13	29,30,31,33	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 8SK L 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 8SK A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.