



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 15, 2024 – 10:44 AM EST

PDB ID : 1NFD
Title : AN ALPHA-BETA T CELL RECEPTOR (TCR) HETERODIMER IN COMPLEX WITH AN ANTI-TCR FAB FRAGMENT DERIVED FROM A MITOGENIC ANTIBODY
Authors : Wang, J.-H.; Lim, K.; Smolyar, A.; Teng, M.-K.; Sacchittini, J.; Reinherz, E.L.
Deposited on : 1997-08-04
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	203	26% 51% 23%
1	C	203	28% 48% 22% .
2	B	239	28% 50% 20% .
2	D	239	27% 52% 20% .
3	E	212	25% 53% 22% .
3	G	212	19% 51% 28% .
4	F	222	16% 58% 24% .
4	H	222	22% 56% 20% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	B	248	X	-	-	-
5	NAG	C	214	X	-	-	-
5	NAG	C	215	X	-	-	-
5	NAG	C	216	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N15 ALPHA-BETA T-CELL RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	0	0
			1589	1002	259	320	8			
1	C	203	Total	C	N	O	S	0	0	0
			1589	1002	259	320	8			

- Molecule 2 is a protein called N15 ALPHA-BETA T-CELL RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	239	Total	C	N	O	S	0	0	0
			1926	1218	338	362	8			
2	D	239	Total	C	N	O	S	0	0	0
			1926	1218	338	362	8			

- Molecule 3 is a protein called H57 FAB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	212	Total	C	N	O	S	0	0	0
			1618	1012	269	330	7			
3	G	212	Total	C	N	O	S	0	0	0
			1618	1012	269	330	7			

- Molecule 4 is a protein called H57 FAB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	222	Total	C	N	O	S	0	0	0
			1711	1081	289	333	8			
4	H	222	Total	C	N	O	S	0	0	0
			1711	1081	289	333	8			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: N15 ALPHA-BETA T-CELL RECEPTOR




• Molecule 1: N15 ALPHA-BETA T-CELL RECEPTOR




• Molecule 2: N15 ALPHA-BETA T-CELL RECEPTOR






• Molecule 2: N15 ALPHA-BETA T-CELL RECEPTOR

Chain D: 27% 52% 20%



• Molecule 3: H57 FAB


Chain E: 25% 53% 22%



• Molecule 3: H57 FAB


Chain G: 19% 51% 28%






• Molecule 4: H57 FAB

Chain F: 16% 58% 24% .



• Molecule 4: H57 FAB

Chain H: 22% 56% 20% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.74Å 122.30Å 115.84Å 90.00° 107.95° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80	Depositor
% Data completeness (in resolution range)	73.5 (15.00-2.80)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.243 , 0.309	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13884	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.37	0/1625	0.66	0/2208
1	C	0.37	0/1625	0.64	2/2208 (0.1%)
2	B	0.74	1/1981 (0.1%)	0.60	0/2683
2	D	0.77	1/1981 (0.1%)	0.62	0/2683
3	E	0.32	0/1653	0.57	0/2249
3	G	0.29	0/1653	0.55	0/2249
4	F	0.30	0/1759	0.56	0/2404
4	H	0.31	0/1759	0.60	0/2404
All	All	0.49	2/14036 (0.0%)	0.60	2/19088 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	191	SER	CB-OG	30.51	1.81	1.42
2	B	191	SER	CB-OG	28.96	1.79	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	203	ASN	N-CA-CB	5.92	121.26	110.60
1	C	203	ASN	CB-CG-ND2	5.45	129.78	116.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1589	0	1526	241	1
1	C	1589	0	1525	166	0
2	B	1926	0	1835	249	2
2	D	1926	0	1836	257	1
3	E	1618	0	1576	269	0
3	G	1618	0	1576	258	0
4	F	1711	0	1647	294	0
4	H	1711	0	1647	283	2
5	A	42	0	39	4	0
5	B	56	0	52	7	0
5	C	42	0	37	8	0
5	D	56	0	52	3	0
All	All	13884	0	13348	1893	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 70.

The worst 5 of 1893 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:117:VAL:HG22	3:E:209:LEU:CD2	1.55	1.35
1:A:34:TRP:HD1	1:A:73:PHE:CE1	1.44	1.34
1:C:203:ASN:ND2	5:C:216:NAG:O7	1.65	1.30
3:E:165:SER:HB2	3:E:173:TYR:CD1	1.66	1.30
1:A:34:TRP:CD1	1:A:73:PHE:HE1	1.49	1.29

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:CYS:O	2:D:57:LYS:NZ[2_546]	2.01	0.19
2:B:53:VAL:O	4:H:70:SER:OG[2_656]	2.09	0.11
2:B:55:ARG:NH2	4:H:19:LYS:NZ[2_656]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/203 (99%)	156 (78%)	33 (16%)	12 (6%)	1	3
1	C	201/203 (99%)	162 (81%)	23 (11%)	16 (8%)	1	1
2	B	237/239 (99%)	198 (84%)	25 (10%)	14 (6%)	1	4
2	D	237/239 (99%)	195 (82%)	30 (13%)	12 (5%)	1	5
3	E	210/212 (99%)	159 (76%)	35 (17%)	16 (8%)	1	2
3	G	210/212 (99%)	158 (75%)	33 (16%)	19 (9%)	0	1
4	F	220/222 (99%)	160 (73%)	39 (18%)	21 (10%)	0	1
4	H	220/222 (99%)	170 (77%)	32 (14%)	18 (8%)	1	1
All	All	1736/1752 (99%)	1358 (78%)	250 (14%)	128 (7%)	1	2

5 of 128 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	58	HIS
1	A	134	ARG
1	A	204	ALA
2	B	39	LEU
1	C	53	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/181 (100%)	127 (70%)	54 (30%)	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	181/181 (100%)	118 (65%)	63 (35%)	0	0
2	B	212/212 (100%)	151 (71%)	61 (29%)	0	1
2	D	212/212 (100%)	147 (69%)	65 (31%)	0	0
3	E	185/185 (100%)	137 (74%)	48 (26%)	0	1
3	G	185/185 (100%)	122 (66%)	63 (34%)	0	0
4	F	192/192 (100%)	135 (70%)	57 (30%)	0	1
4	H	192/192 (100%)	144 (75%)	48 (25%)	0	1
All	All	1540/1540 (100%)	1081 (70%)	459 (30%)	0	1

5 of 459 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	194	SER
4	H	153	THR
3	E	199	GLU
4	H	140	LEU
3	G	190	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
3	E	137	ASN
4	F	212	HIS
3	E	171	GLN
4	F	53	ASN
3	G	31	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	215	1	14,14,15	0.41	0	17,19,21	0.62	1 (5%)
5	NAG	C	214	1	14,14,15	0.45	0	17,19,21	0.62	0
5	NAG	A	214	1	14,14,15	1.08	1 (7%)	17,19,21	1.08	1 (5%)
5	NAG	D	248	2	14,14,15	0.38	0	17,19,21	0.55	0
5	NAG	D	250	2	14,14,15	0.43	0	17,19,21	0.63	0
5	NAG	B	251	2	14,14,15	0.52	0	17,19,21	0.64	0
5	NAG	B	250	2	14,14,15	0.40	0	17,19,21	0.86	1 (5%)
5	NAG	B	248	2	14,14,15	0.46	0	17,19,21	0.83	1 (5%)
5	NAG	D	251	2	14,14,15	0.51	0	17,19,21	0.63	0
5	NAG	C	215	1	14,14,15	0.43	0	17,19,21	0.67	0
5	NAG	A	216	1	14,14,15	0.53	0	17,19,21	0.78	0
5	NAG	D	249	2	14,14,15	0.53	0	17,19,21	0.67	0
5	NAG	C	216	1	14,14,15	4.48	6 (42%)	17,19,21	2.19	6 (35%)
5	NAG	B	249	2	14,14,15	0.37	0	17,19,21	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	215	1	-	2/6/23/26	0/1/1/1
5	NAG	C	214	1	1/1/5/7	2/6/23/26	0/1/1/1
5	NAG	A	214	1	-	1/6/23/26	0/1/1/1
5	NAG	D	248	2	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	D	250	2	-	2/6/23/26	0/1/1/1
5	NAG	B	251	2	-	3/6/23/26	0/1/1/1
5	NAG	B	250	2	-	3/6/23/26	0/1/1/1
5	NAG	B	248	2	1/1/5/7	0/6/23/26	0/1/1/1
5	NAG	D	251	2	-	2/6/23/26	0/1/1/1
5	NAG	C	215	1	1/1/5/7	4/6/23/26	0/1/1/1
5	NAG	A	216	1	-	2/6/23/26	0/1/1/1
5	NAG	D	249	2	-	3/6/23/26	0/1/1/1
5	NAG	C	216	1	1/1/5/7	3/6/23/26	0/1/1/1
5	NAG	B	249	2	-	2/6/23/26	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	216	NAG	C1-C2	-11.73	1.36	1.52
5	C	216	NAG	O5-C1	7.89	1.56	1.43
5	C	216	NAG	C4-C5	-5.54	1.41	1.53
5	C	216	NAG	C8-C7	4.47	1.59	1.50
5	C	216	NAG	C3-C2	-3.84	1.44	1.52

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	216	NAG	O5-C5-C4	4.83	122.58	110.83
5	C	216	NAG	C4-C3-C2	3.92	116.77	111.02
5	C	216	NAG	C1-C2-N2	-3.36	105.14	110.43
5	A	214	NAG	C2-N2-C7	-3.19	118.62	122.90
5	C	216	NAG	C6-C5-C4	2.96	120.28	113.02

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	248	NAG	C1
5	C	214	NAG	C1
5	C	215	NAG	C1
5	C	216	NAG	C1

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	216	NAG	C1-C2-N2-C7
5	A	215	NAG	C4-C5-C6-O6
5	C	214	NAG	C4-C5-C6-O6
5	C	216	NAG	O5-C5-C6-O6
5	A	215	NAG	O5-C5-C6-O6

There are no ring outliers.

9 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	214	NAG	3	0
5	A	214	NAG	2	0
5	D	248	NAG	2	0
5	B	251	NAG	1	0
5	B	248	NAG	3	0
5	A	216	NAG	2	0
5	D	249	NAG	1	0
5	C	216	NAG	5	0
5	B	249	NAG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.