



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 28, 2024 – 03:01 pm GMT

PDB ID : 5NFY
Title : SARS-CoV nsp10/nsp14 dynamic complex
Authors : Ferron, F.; Gluais, L.; Vonnrhein, C.; Bricogne, G.; Canard, B.; Imbert, I.
Deposited on : 2017-03-16
Resolution : 3.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1137 (3.44-3.32)
Clashscore	180529	1165 (3.44-3.32)
Ramachandran outliers	177936	1164 (3.44-3.32)
Sidechain outliers	177891	1164 (3.44-3.32)
RSRZ outliers	164620	1137 (3.44-3.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>..</div> </div> </div>
1	B	534	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>10%</div> <div>..</div> </div> </div>
1	C	534	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>9%</div> <div>..</div> </div> </div>
1	D	534	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>9%</div> <div>..</div> </div> </div>
2	M	138	<div> <div>0%</div> <div> <div></div> <div>88%</div> <div>8%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	N	138	
2	O	138	
2	P	138	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ZN	A	603	-	-	X	-
3	ZN	C	601	-	-	X	-
3	ZN	P	201	-	-	X	-
4	PEG	A	604	-	-	X	-
4	PEG	C	604	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyprotein 1ab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	512	Total	C	N	O	S	0	0	0
			4098	2626	702	734	36			
1	B	514	Total	C	N	O	S	0	0	0
			4112	2634	704	738	36			
1	C	514	Total	C	N	O	S	0	0	0
			4112	2634	704	738	36			
1	D	514	Total	C	N	O	S	0	0	0
			4112	2634	704	738	36			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP Q1T6X8
A	-5	HIS	-	expression tag	UNP Q1T6X8
A	-4	HIS	-	expression tag	UNP Q1T6X8
A	-3	HIS	-	expression tag	UNP Q1T6X8
A	-2	HIS	-	expression tag	UNP Q1T6X8
A	-1	HIS	-	expression tag	UNP Q1T6X8
A	0	HIS	-	expression tag	UNP Q1T6X8
B	-6	MET	-	initiating methionine	UNP Q1T6X8
B	-5	HIS	-	expression tag	UNP Q1T6X8
B	-4	HIS	-	expression tag	UNP Q1T6X8
B	-3	HIS	-	expression tag	UNP Q1T6X8
B	-2	HIS	-	expression tag	UNP Q1T6X8
B	-1	HIS	-	expression tag	UNP Q1T6X8
B	0	HIS	-	expression tag	UNP Q1T6X8
C	-6	MET	-	initiating methionine	UNP Q1T6X8
C	-5	HIS	-	expression tag	UNP Q1T6X8
C	-4	HIS	-	expression tag	UNP Q1T6X8
C	-3	HIS	-	expression tag	UNP Q1T6X8
C	-2	HIS	-	expression tag	UNP Q1T6X8
C	-1	HIS	-	expression tag	UNP Q1T6X8
C	0	HIS	-	expression tag	UNP Q1T6X8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-6	MET	-	initiating methionine	UNP Q1T6X8
D	-5	HIS	-	expression tag	UNP Q1T6X8
D	-4	HIS	-	expression tag	UNP Q1T6X8
D	-3	HIS	-	expression tag	UNP Q1T6X8
D	-2	HIS	-	expression tag	UNP Q1T6X8
D	-1	HIS	-	expression tag	UNP Q1T6X8
D	0	HIS	-	expression tag	UNP Q1T6X8

- Molecule 2 is a protein called Polyprotein 1ab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	132	Total	C	N	O	S	0	0	0
			969	601	165	187	16			
2	N	133	Total	C	N	O	S	0	0	0
			979	607	168	188	16			
2	O	133	Total	C	N	O	S	0	0	0
			979	607	168	188	16			
2	P	133	Total	C	N	O	S	0	0	0
			979	607	168	188	16			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	-6	MET	-	initiating methionine	UNP Q1T6X8
M	-5	HIS	-	expression tag	UNP Q1T6X8
M	-4	HIS	-	expression tag	UNP Q1T6X8
M	-3	HIS	-	expression tag	UNP Q1T6X8
M	-2	HIS	-	expression tag	UNP Q1T6X8
M	-1	HIS	-	expression tag	UNP Q1T6X8
M	0	HIS	-	expression tag	UNP Q1T6X8
N	-6	MET	-	initiating methionine	UNP Q1T6X8
N	-5	HIS	-	expression tag	UNP Q1T6X8
N	-4	HIS	-	expression tag	UNP Q1T6X8
N	-3	HIS	-	expression tag	UNP Q1T6X8
N	-2	HIS	-	expression tag	UNP Q1T6X8
N	-1	HIS	-	expression tag	UNP Q1T6X8
N	0	HIS	-	expression tag	UNP Q1T6X8
O	-6	MET	-	initiating methionine	UNP Q1T6X8
O	-5	HIS	-	expression tag	UNP Q1T6X8
O	-4	HIS	-	expression tag	UNP Q1T6X8
O	-3	HIS	-	expression tag	UNP Q1T6X8
O	-2	HIS	-	expression tag	UNP Q1T6X8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
O	-1	HIS	-	expression tag	UNP Q1T6X8
O	0	HIS	-	expression tag	UNP Q1T6X8
P	-6	MET	-	initiating methionine	UNP Q1T6X8
P	-5	HIS	-	expression tag	UNP Q1T6X8
P	-4	HIS	-	expression tag	UNP Q1T6X8
P	-3	HIS	-	expression tag	UNP Q1T6X8
P	-2	HIS	-	expression tag	UNP Q1T6X8
P	-1	HIS	-	expression tag	UNP Q1T6X8
P	0	HIS	-	expression tag	UNP Q1T6X8

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total 3	Zn 3	0	0
3	B	3	Total 3	Zn 3	0	0
3	C	3	Total 3	Zn 3	0	0
3	D	3	Total 3	Zn 3	0	0
3	M	2	Total 2	Zn 2	0	0
3	N	2	Total 2	Zn 2	0	0
3	O	2	Total 2	Zn 2	0	0
3	P	2	Total 2	Zn 2	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		
4	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	59	Total	O	0	0
			59	59		
5	B	56	Total	O	0	0
			56	56		
5	C	67	Total	O	0	0
			67	67		
5	D	34	Total	O	0	0
			34	34		
5	M	7	Total	O	0	0
			7	7		
5	N	6	Total	O	0	0
			6	6		
5	O	7	Total	O	0	0
			7	7		
5	P	8	Total	O	0	0
			8	8		

- Molecule 1: Polypeptide 1ab







● Molecule 1: Polyprotein 1ab





● Molecule 2: Polyprotein 1ab





● Molecule 2: Polyprotein 1ab



● Molecule 2: Polyprotein 1ab



● Molecule 2: Polyprotein 1ab



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	185.88Å 189.79Å 196.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	66.40 – 3.38 66.40 – 3.38	Depositor EDS
% Data completeness (in resolution range)	99.5 (66.40-3.38) 99.5 (66.40-3.38)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 3.41Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.193 , 0.235 0.201 , 0.235	Depositor DCC
R_{free} test set	92531 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	100.4	Xtriage
Anisotropy	0.206	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 76.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.004 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	20625	wwPDB-VP
Average B, all atoms (Å ²)	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.85 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.4350e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.82	1/4215 (0.0%)	0.96	7/5727 (0.1%)
1	B	0.79	0/4229	0.94	8/5746 (0.1%)
1	C	0.81	0/4229	0.96	12/5746 (0.2%)
1	D	0.78	0/4229	0.92	5/5746 (0.1%)
2	M	0.71	0/992	0.89	1/1350 (0.1%)
2	N	0.74	0/1003	0.85	0/1365
2	O	0.67	0/1003	0.83	1/1365 (0.1%)
2	P	0.71	0/1003	0.89	1/1365 (0.1%)
All	All	0.79	1/20903 (0.0%)	0.93	35/28410 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	99	ASP	CB-CG	5.29	1.62	1.51

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	62	MET	CG-SD-CE	-7.70	87.88	100.20
1	A	213	ARG	NE-CZ-NH2	-7.49	116.55	120.30
1	C	301	ASP	N-CA-C	-7.41	91.00	111.00
1	A	210	CYS	CA-CB-SG	7.14	126.86	114.00
1	C	423	LYS	N-CA-C	-7.11	91.80	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4098	0	3972	91	1
1	B	4112	0	3986	69	1
1	C	4112	0	3987	72	1
1	D	4112	0	3985	87	1
2	M	969	0	921	4	0
2	N	979	0	929	11	0
2	O	979	0	928	34	0
2	P	979	0	931	23	0
3	A	3	0	0	2	0
3	B	3	0	0	0	0
3	C	3	0	0	2	0
3	D	3	0	0	0	0
3	M	2	0	0	0	0
3	N	2	0	0	0	0
3	O	2	0	0	0	0
3	P	2	0	0	2	0
4	A	14	0	20	8	0
4	C	7	0	10	3	0
5	A	59	0	0	1	0
5	B	56	0	0	2	0
5	C	67	0	0	4	0
5	D	34	0	0	0	0
5	M	7	0	0	2	0
5	N	6	0	0	0	0
5	O	7	0	0	0	0
5	P	8	0	0	0	0
All	All	20625	0	19669	388	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 388 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:O:117:CYS:CB	2:O:124:LYS:HD2	1.45	1.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:477:CYS:SG	3:A:603:ZN:ZN	1.10	1.38
2:O:110:PHE:HE2	2:O:123:TRP:NE1	1.24	1.29
1:D:18:LEU:HD21	1:D:53:ARG:NE	1.45	1.29
2:O:117:CYS:CA	2:O:124:LYS:HD2	1.61	1.28

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:ARG:NH1	1:D:515:ASP:OD2[4_557]	1.35	0.85
1:B:515:ASP:OD2	1:C:52:ARG:NH2[4_456]	1.81	0.39

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/534 (95%)	450 (89%)	44 (9%)	14 (3%)	4	19
1	B	510/534 (96%)	456 (89%)	37 (7%)	17 (3%)	3	17
1	C	510/534 (96%)	449 (88%)	44 (9%)	17 (3%)	3	17
1	D	510/534 (96%)	461 (90%)	38 (8%)	11 (2%)	5	24
2	M	130/138 (94%)	116 (89%)	11 (8%)	3 (2%)	5	23
2	N	131/138 (95%)	119 (91%)	9 (7%)	3 (2%)	5	23
2	O	131/138 (95%)	115 (88%)	13 (10%)	3 (2%)	5	23
2	P	131/138 (95%)	122 (93%)	8 (6%)	1 (1%)	16	45
All	All	2561/2688 (95%)	2288 (89%)	204 (8%)	69 (3%)	4	20

5 of 69 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	LYS
1	A	222	ASP
1	A	250	THR
1	A	483	VAL
1	B	15	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	450/470 (96%)	442 (98%)	8 (2%)	54	73
1	B	451/470 (96%)	437 (97%)	14 (3%)	35	61
1	C	451/470 (96%)	443 (98%)	8 (2%)	54	73
1	D	451/470 (96%)	436 (97%)	15 (3%)	33	59
2	M	107/113 (95%)	104 (97%)	3 (3%)	38	63
2	N	108/113 (96%)	107 (99%)	1 (1%)	75	86
2	O	108/113 (96%)	105 (97%)	3 (3%)	38	63
2	P	108/113 (96%)	106 (98%)	2 (2%)	52	72
All	All	2234/2332 (96%)	2180 (98%)	54 (2%)	44	67

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	479	LEU
1	D	18	LEU
2	O	87	LYS
1	C	483	VAL
1	D	7	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	408	ASN
2	N	85	ASN
2	P	0	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 20 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	605	-	6,6,6	0.33	0	5,5,5	0.32	0
4	PEG	A	604	-	6,6,6	0.33	0	5,5,5	0.19	0
4	PEG	C	604	-	6,6,6	0.41	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	605	-	-	4/4/4/4	-
4	PEG	A	604	-	-	3/4/4/4	-
4	PEG	C	604	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	604	PEG	O1-C1-C2-O2
4	C	604	PEG	C4-C3-O2-C2
4	A	604	PEG	C1-C2-O2-C3
4	A	604	PEG	O1-C1-C2-O2
4	A	605	PEG	C4-C3-O2-C2

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	604	PEG	8	0
4	C	604	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	512/534 (95%)	-0.04	10 (1%) 64 55	61, 89, 127, 156	0
1	B	514/534 (96%)	-0.05	18 (3%) 47 40	61, 87, 128, 192	0
1	C	514/534 (96%)	-0.08	11 (2%) 63 53	55, 93, 147, 171	0
1	D	514/534 (96%)	0.20	24 (4%) 37 33	66, 99, 142, 193	0
2	M	132/138 (95%)	-0.06	1 (0%) 82 76	69, 103, 129, 165	0
2	N	133/138 (96%)	0.08	2 (1%) 71 63	70, 92, 123, 162	0
2	O	133/138 (96%)	0.54	7 (5%) 33 30	83, 121, 159, 181	0
2	P	133/138 (96%)	-0.13	2 (1%) 71 63	66, 88, 124, 151	0
All	All	2585/2688 (96%)	0.03	75 (2%) 54 46	55, 94, 141, 193	0

The worst 5 of 75 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	482	ALA	6.6
1	D	468	LEU	6.3
1	B	481	GLY	5.8
1	D	481	GLY	5.8
1	D	483	VAL	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PEG	C	604	7/7	0.51	0.47	92,105,127,133	0
3	ZN	O	202	1/1	0.63	0.30	175,175,175,175	0
4	PEG	A	604	7/7	0.82	0.25	83,87,103,115	0
3	ZN	N	202	1/1	0.86	0.25	148,148,148,148	0
4	PEG	A	605	7/7	0.88	0.29	99,103,118,128	0
3	ZN	A	601	1/1	0.89	0.10	161,161,161,161	0
3	ZN	D	603	1/1	0.90	0.20	115,115,115,115	0
3	ZN	D	601	1/1	0.90	0.09	144,144,144,144	0
3	ZN	B	603	1/1	0.92	0.13	119,119,119,119	0
3	ZN	O	201	1/1	0.93	0.14	196,196,196,196	0
3	ZN	C	603	1/1	0.93	0.17	168,168,168,168	0
3	ZN	P	202	1/1	0.93	0.23	142,142,142,142	0
3	ZN	B	601	1/1	0.96	0.16	123,123,123,123	0
3	ZN	P	201	1/1	0.96	0.11	161,161,161,161	0
3	ZN	M	202	1/1	0.96	0.20	155,155,155,155	0
3	ZN	C	601	1/1	0.97	0.06	155,155,155,155	0
3	ZN	C	602	1/1	0.97	0.18	107,107,107,107	0
3	ZN	B	602	1/1	0.99	0.17	111,111,111,111	0
3	ZN	M	201	1/1	0.99	0.14	112,112,112,112	0
3	ZN	A	603	1/1	0.99	0.06	92,92,92,92	0
3	ZN	N	201	1/1	0.99	0.16	88,88,88,88	0
3	ZN	A	602	1/1	0.99	0.14	102,102,102,102	0
3	ZN	D	602	1/1	0.99	0.14	107,107,107,107	0

6.5 Other polymers ⓘ

There are no such residues in this entry.