



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 27, 2024 – 03:05 PM EDT

PDB ID : 1NTZ  
Title : Crystal Structure of Mitochondrial Cytochrome bc1 Complex Bound with Ubiquinone  
Authors : Gao, X.; Wen, X.; Esser, L.; Quinn, B.; Yu, L.; Yu, C.-A.; Xia, D.  
Deposited on : 2003-01-30  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	446	
2	B	439	
3	C	379	
4	D	241	
5	E	196	
6	F	110	
7	G	81	
8	H	78	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
9	I	57	
10	J	62	
11	K	56	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	UQ2	C	383	-	-	X	-

## 2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 16896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquinol-cytochrome C reductase complex core protein I, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	446	3458	2161	609	668	20	0	0	0

- Molecule 2 is a protein called Ubiquinol-cytochrome C reductase complex core protein 2, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	423	3172	1993	562	610	7	0	0	0

- Molecule 3 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	378	3003	2013	471	501	18	0	0	0

- Molecule 4 is a protein called cytochrome c1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	241	1918	1225	330	348	15	0	0	0

- Molecule 5 is a protein called UBIQUINOL-CYTOCHROME C REDUCTASE IRON-SULFUR SUBUNIT, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	196	1519	957	263	291	8	0	0	0

- Molecule 6 is a protein called Ubiquinol-cytochrome C reductase complex 14 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	105	911	576	165	168	2	0	0	0

- Molecule 7 is a protein called Ubiquinol-cytochrome C reductase complex ubiquinone-binding protein QP-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	75	628	410	118	99	1	0	0	0

- Molecule 8 is a protein called Ubiquinol-cytochrome C reductase complex 11 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	70	575	347	102	121	5	0	0	0

- Molecule 9 is a protein called Ubiquinol-cytochrome C reductase 8 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	57	406	253	77	74	2	0	0	0

- Molecule 10 is a protein called Ubiquinol-cytochrome C reductase complex 7.2 kDa protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
10	J	61	483	316	82	85	0	0	0

- Molecule 11 is a protein called Ubiquinol-cytochrome C reductase complex 6.4 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	K	53	437	292	78	66	1	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	22	GLN	SER	SEE REMARK 999	UNP P07552


- Molecule 12 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
13	C	1	23	19	4	0	0

- Molecule 14 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
14	E	1	4	2	2	0	0

- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
15	A	29	29	29	0	0
15	B	39	39	39	0	0
15	C	78	78	78	0	0
15	D	30	30	30	0	0
15	E	3	3	3	0	0
15	F	14	14	14	0	0
15	G	4	4	4	0	0

Continued on next page...

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
15	H	1	Total O 1 1	0	0
15	I	4	Total O 4 4	0	0
15	J	2	Total O 2 2	0	0
15	K	3	Total O 3 3	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Ubiquinol-cytochrome C reductase complex core protein I, mitochondrial




- Molecule 2: Ubiquinol-cytochrome C reductase complex core protein 2, mitochondrial




- Molecule 3: Cytochrome b






- Molecule 4: cytochrome c1

Chain D: 55% 33% 11%




- Molecule 5: UBIQUINOL-CYTOCHROME C REDUCTASE IRON-SULFUR SUBUNIT, mitochondrial

Chain E: 78% 20%



- Molecule 6: Ubiquinol-cytochrome C reductase complex 14 kDa protein

Chain F: 72% 15% 8% 5%



- Molecule 7: Ubiquinol-cytochrome C reductase complex ubiquinone-binding protein QP-C

Chain G: 65% 20% 6% 7%



- Molecule 8: Ubiquinol-cytochrome C reductase complex 11 kDa protein



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.83Å 153.83Å 596.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.99 – 2.60	Depositor
% Data completeness (in resolution range)	100.0 (28.99-2.60)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REFMAC 5.0	Depositor
R, $R_{free}$	0.247 , 0.283	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	16896	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FES, UQ2, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.91	2/3531 (0.1%)	0.88	17/4792 (0.4%)
2	B	1.11	6/3232 (0.2%)	0.90	11/4386 (0.3%)
3	C	0.89	3/3100 (0.1%)	0.82	7/4242 (0.2%)
4	D	0.76	0/1977	0.89	8/2684 (0.3%)
5	E	0.75	1/1553 (0.1%)	0.81	5/2100 (0.2%)
6	F	1.11	2/930 (0.2%)	0.94	4/1246 (0.3%)
7	G	0.98	0/649	0.77	0/878
8	H	0.69	0/580	0.98	7/777 (0.9%)
9	I	1.29	2/411 (0.5%)	1.28	4/558 (0.7%)
10	J	0.78	0/495	0.79	1/672 (0.1%)
11	K	0.76	0/453	0.72	1/621 (0.2%)
All	All	0.93	16/16911 (0.1%)	0.88	65/22956 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0
2	B	1	0
3	C	1	2
4	D	2	0
7	G	1	0
8	H	1	0
9	I	1	0
10	J	1	0
All	All	9	2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	379	TRP	CB-CG	-8.67	1.34	1.50
9	I	25	ALA	CA-CB	-7.45	1.36	1.52
2	B	424	MET	SD-CE	-7.45	1.36	1.77
2	B	309	VAL	CB-CG1	-7.40	1.37	1.52
1	A	281	ASP	CB-CG	-6.75	1.37	1.51

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	221	HIS	N-CA-C	7.66	131.69	111.00
1	A	42	ASP	CB-CG-OD2	7.55	125.10	118.30
4	D	122	GLY	N-CA-C	-7.31	94.83	113.10
3	C	252	ASP	CB-CG-OD2	7.04	124.64	118.30
6	F	85	GLU	N-CA-C	-7.00	92.09	111.00

5 of 9 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	443	TRP	CA
2	B	305	GLN	CA
3	C	221	HIS	CA
4	D	145	GLU	CA
4	D	169	LEU	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	221	HIS	Mainchain
3	C	345	HIS	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3458	0	3356	69	0
2	B	3172	0	3152	69	0
3	C	3003	0	3065	76	0
4	D	1918	0	1870	65	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	E	1519	0	1504	20	0
6	F	911	0	904	15	0
7	G	628	0	636	15	0
8	H	575	0	550	14	0
9	I	406	0	437	61	0
10	J	483	0	465	20	0
11	K	437	0	439	7	0
12	C	86	0	60	5	0
12	D	43	0	30	3	0
13	C	46	0	52	19	0
14	E	4	0	0	0	0
15	A	29	0	0	3	0
15	B	39	0	0	2	0
15	C	78	0	0	6	0
15	D	30	0	0	1	0
15	E	3	0	0	0	0
15	F	14	0	0	3	0
15	G	4	0	0	0	0
15	H	1	0	0	0	0
15	I	4	0	0	1	0
15	J	2	0	0	0	0
15	K	3	0	0	0	0
All	All	16896	0	16520	376	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 376 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:11:MET:SD	3:C:11:MET:CE	2.09	1.41
4:D:37:CYS:SG	12:D:242:HEM:HAB	1.90	1.12
10:J:18:SER:HA	11:K:24:TRP:HZ3	1.19	1.03
1:A:325:VAL:HG21	9:I:43:LEU:HD12	1.44	0.98
3:C:221:HIS:CG	3:C:221:HIS:O	2.13	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/446 (100%)	409 (92%)	21 (5%)	14 (3%)	3	5
2	B	421/439 (96%)	393 (93%)	20 (5%)	8 (2%)	6	13
3	C	376/379 (99%)	349 (93%)	18 (5%)	9 (2%)	5	9
4	D	239/241 (99%)	183 (77%)	36 (15%)	20 (8%)	0	0
5	E	194/196 (99%)	175 (90%)	15 (8%)	4 (2%)	5	11
6	F	103/110 (94%)	98 (95%)	3 (3%)	2 (2%)	6	13
7	G	73/81 (90%)	65 (89%)	3 (4%)	5 (7%)	1	1
8	H	68/78 (87%)	49 (72%)	10 (15%)	9 (13%)	0	0
9	I	55/57 (96%)	26 (47%)	16 (29%)	13 (24%)	0	0
10	J	59/62 (95%)	45 (76%)	9 (15%)	5 (8%)	0	0
11	K	51/56 (91%)	42 (82%)	4 (8%)	5 (10%)	0	0
All	All	2083/2145 (97%)	1834 (88%)	155 (7%)	94 (4%)	2	2

5 of 94 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	159	GLN
1	A	220	SER
1	A	227	ALA
2	B	305	GLN
3	C	222	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/370 (100%)	339 (92%)	31 (8%)	9	19
2	B	332/343 (97%)	301 (91%)	31 (9%)	7	15
3	C	326/327 (100%)	305 (94%)	21 (6%)	14	32
4	D	206/206 (100%)	173 (84%)	33 (16%)	2	3
5	E	168/168 (100%)	157 (94%)	11 (6%)	14	31
6	F	96/98 (98%)	86 (90%)	10 (10%)	5	11
7	G	66/71 (93%)	62 (94%)	4 (6%)	15	34
8	H	67/74 (90%)	55 (82%)	12 (18%)	1	2
9	I	44/44 (100%)	31 (70%)	13 (30%)	0	0
10	J	46/52 (88%)	41 (89%)	5 (11%)	5	10
11	K	42/46 (91%)	38 (90%)	4 (10%)	7	14
All	All	1763/1799 (98%)	1588 (90%)	175 (10%)	6	13

5 of 175 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	228	SER
8	H	14	VAL
5	E	12	ASP
6	F	53	ASN
8	H	65	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	385	GLN
3	C	267	HIS
9	I	31	GLN
2	B	401	GLN
3	C	68	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
12	HEM	D	242	4	42,50,50	3.00	20 (47%)	46,82,82	2.44	19 (41%)
14	FES	E	200	5	0,4,4	-	-	-	-	-
13	UQ2	C	384	-	23,23,23	2.62	5 (21%)	30,31,31	1.52	5 (16%)
13	UQ2	C	383	-	23,23,23	2.38	6 (26%)	30,31,31	1.78	7 (23%)
12	HEM	C	382	3	42,50,50	3.01	20 (47%)	46,82,82	2.30	16 (34%)
12	HEM	C	381	3	42,50,50	2.99	20 (47%)	46,82,82	2.38	16 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	HEM	D	242	4	-	5/12/54/54	-
14	FES	E	200	5	-	-	0/1/1/1
13	UQ2	C	384	-	-	3/15/39/39	0/1/1/1
13	UQ2	C	383	-	-	5/15/39/39	0/1/1/1
12	HEM	C	382	3	-	2/12/54/54	-
12	HEM	C	381	3	-	6/12/54/54	-

The worst 5 of 71 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	C	384	UQ2	C6-C5	10.22	1.53	1.35
13	C	383	UQ2	C6-C5	8.18	1.49	1.35
12	D	242	HEM	C4D-C3D	-7.32	1.32	1.45
12	C	381	HEM	C3C-C2C	-7.24	1.30	1.40
12	D	242	HEM	C1B-C2B	-6.47	1.31	1.44

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	C	382	HEM	C4C-CHD-C1D	-8.40	111.47	122.56
12	C	381	HEM	C4B-CHC-C1C	-7.18	113.08	122.56
12	C	382	HEM	C4B-CHC-C1C	-6.75	113.65	122.56
12	D	242	HEM	C4B-CHC-C1C	-6.45	114.05	122.56
12	C	381	HEM	C4C-CHD-C1D	-6.22	114.36	122.56

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
12	C	381	HEM	C2B-C3B-CAB-CBB
12	D	242	HEM	C2B-C3B-CAB-CBB
13	C	383	UQ2	C1-C6-C7-C8
13	C	384	UQ2	C1-C6-C7-C8
13	C	384	UQ2	C5-C6-C7-C8


There are no ring outliers.



5 monomers are involved in 27 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	D	242	HEM	3	0
13	C	384	UQ2	8	0
13	C	383	UQ2	11	0
12	C	382	HEM	2	0
12	C	381	HEM	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.