



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 18, 2024 – 09:53 PM EDT

PDB ID : 4O4X
Title : Crystal structure of the vaccine antigen Transferrin Binding Protein B (TbpB) double mutant Tyr-167-Ala and Trp-176-Ala from Haemophilus parasuis Hp5
Authors : Calmettes, C.; Yu, R.H.; Schryvers, A.B.; Moraes, T.F.
Deposited on : 2013-12-19
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	532	
1	B	532	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	604	-	-	X	-
2	GOL	A	605	-	-	-	X
2	GOL	A	608	-	-	-	X
2	GOL	A	609	-	-	X	-
3	SO4	A	611	-	-	X	X
3	SO4	B	603	-	-	X	-
3	SO4	B	605	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TbpB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	503	Total	C	N	O	S	0	1	0
			3892	2452	651	779	10			
1	B	502	Total	C	N	O	S	0	0	0
			3877	2442	648	777	10			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	37	Total	O	0	0
			37	37		
5	B	13	Total	O	0	0
			13	13		

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	239.89Å 42.24Å 114.02Å 90.00° 92.42° 90.00°	Depositor
Resolution (Å)	42.18 – 2.90 42.18 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (42.18-2.90) 99.4 (42.18-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.90Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.216 , 0.234 0.214 , 0.232	Depositor DCC
R_{free} test set	1293 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	57.7	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.004 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7924	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/3968	0.47	0/5346
1	B	0.28	0/3950	0.45	0/5323
All	All	0.30	0/7918	0.46	0/10669

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	348	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3892	0	3788	108	0
1	B	3877	0	3767	63	0
2	A	42	0	56	11	0
2	B	12	0	16	3	0
3	A	20	0	0	3	0
3	B	30	0	0	5	0
4	A	1	0	0	0	0
5	A	37	0	0	4	0
5	B	13	0	0	1	0
All	All	7924	0	7627	168	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:346:ILE:CG2	1:A:347[B]:ASN:H	1.52	1.22
1:A:346:ILE:CG2	1:A:347[A]:ASN:H	1.52	1.21
1:A:327:ASP:HB2	1:A:346:ILE:HG21	1.43	1.00
1:A:346:ILE:HG22	1:A:347[A]:ASN:H	0.82	0.97
1:A:346:ILE:HG22	1:A:347[B]:ASN:H	0.82	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/532 (94%)	472 (95%)	25 (5%)	1 (0%)	47	78
1	B	496/532 (93%)	470 (95%)	25 (5%)	1 (0%)	47	78
All	All	994/1064 (93%)	942 (95%)	50 (5%)	2 (0%)	47	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	B	328	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/439 (94%)	400 (96%)	15 (4%)	35	69
1	B	413/439 (94%)	402 (97%)	11 (3%)	44	77
All	All	828/878 (94%)	802 (97%)	26 (3%)	40	74

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	515	ARG
1	B	175	THR
1	B	416	ASN
1	B	103	GLU
1	B	180	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	442	ASN
1	B	513	GLN
1	A	256	GLN
1	A	377	GLN
1	B	81	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 1 is monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	601	-	5,5,5	0.37	0	5,5,5	0.20	0
3	SO4	B	602	-	4,4,4	0.29	0	6,6,6	0.16	0
3	SO4	A	611	-	4,4,4	0.38	0	6,6,6	0.19	0
2	GOL	A	609	-	5,5,5	0.18	0	5,5,5	0.34	0
3	SO4	A	606	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	B	605	-	4,4,4	0.25	0	6,6,6	0.34	0
2	GOL	A	604	-	5,5,5	0.10	0	5,5,5	0.62	0
2	GOL	A	608	-	5,5,5	0.16	0	5,5,5	0.49	0
3	SO4	B	604	-	4,4,4	0.28	0	6,6,6	0.33	0
3	SO4	A	612	-	4,4,4	0.36	0	6,6,6	0.19	0
2	GOL	B	608	-	5,5,5	0.24	0	5,5,5	0.43	0
3	SO4	B	606	-	4,4,4	0.23	0	6,6,6	0.40	0
2	GOL	A	602	-	5,5,5	0.13	0	5,5,5	0.70	0
3	SO4	B	601	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	A	607	-	4,4,4	0.32	0	6,6,6	0.22	0
2	GOL	A	605	-	5,5,5	0.38	0	5,5,5	0.23	0
2	GOL	A	603	-	5,5,5	0.37	0	5,5,5	0.36	0
2	GOL	B	607	-	5,5,5	0.12	0	5,5,5	0.50	0
3	SO4	B	603	-	4,4,4	0.25	0	6,6,6	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	605	-	-	2/4/4/4	-
2	GOL	B	608	-	-	3/4/4/4	-
2	GOL	A	602	-	-	4/4/4/4	-
2	GOL	A	601	-	-	2/4/4/4	-
2	GOL	A	603	-	-	0/4/4/4	-
2	GOL	B	607	-	-	0/4/4/4	-
2	GOL	A	604	-	-	4/4/4/4	-
2	GOL	A	608	-	-	3/4/4/4	-
2	GOL	A	609	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	O1-C1-C2-C3
2	A	602	GOL	O1-C1-C2-O2
2	A	602	GOL	O1-C1-C2-C3
2	A	604	GOL	O1-C1-C2-C3
2	A	604	GOL	C1-C2-C3-O3

There are no ring outliers.

10 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	602	SO4	1	0
3	A	611	SO4	2	0
2	A	609	GOL	4	0
2	A	604	GOL	5	0
2	A	608	GOL	1	0
3	A	612	SO4	1	0
2	B	608	GOL	2	0
2	A	605	GOL	1	0
2	B	607	GOL	1	0
3	B	603	SO4	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	503/532 (94%)	0.13	13 (2%) 56 52	19, 47, 91, 116	0
1	B	502/532 (94%)	0.27	10 (1%) 65 63	23, 58, 93, 128	0
All	All	1005/1064 (94%)	0.20	23 (2%) 60 58	19, 53, 92, 128	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	348	GLY	6.4
1	A	520	ASN	6.1
1	A	481	LEU	5.0
1	B	521	GLY	4.4
1	B	520	ASN	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	608	6/6	0.73	0.62	57,69,72,74	0
2	GOL	A	604	6/6	0.74	0.20	48,51,65,66	0
3	SO4	A	607	5/5	0.75	0.36	55,60,69,84	5
3	SO4	A	606	5/5	0.76	0.37	46,49,51,60	5
2	GOL	B	607	6/6	0.76	0.27	57,64,77,79	0
3	SO4	A	611	5/5	0.76	0.51	45,48,51,54	5
2	GOL	A	605	6/6	0.79	0.41	52,63,71,76	0
3	SO4	B	605	5/5	0.79	0.40	41,44,51,53	5
3	SO4	B	602	5/5	0.80	0.21	102,108,119,132	0
3	SO4	A	612	5/5	0.81	0.38	47,56,65,80	5
3	SO4	B	604	5/5	0.85	0.22	69,72,82,89	5
3	SO4	B	606	5/5	0.85	0.29	53,55,62,62	5
3	SO4	B	603	5/5	0.86	0.25	46,47,54,54	5
3	SO4	B	601	5/5	0.88	0.26	62,63,68,77	5
2	GOL	A	609	6/6	0.88	0.17	64,68,71,77	0
2	GOL	A	603	6/6	0.88	0.15	52,56,59,62	0
2	GOL	A	602	6/6	0.89	0.24	31,37,43,47	6
2	GOL	B	608	6/6	0.89	0.26	52,60,64,65	0
2	GOL	A	601	6/6	0.89	0.27	36,58,64,65	0
4	NA	A	610	1/1	0.94	0.23	65,65,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.