



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2024 – 08:07 pm BST

PDB ID : 5OAW
Title : Crystal structure of Aspergillus fumigatus N-acetylphosphoglucosamine mutase in complex with GlcNAc-6P and magnesium
Authors : Raimi, O.G.; Hurtado-Guerrero, R.
Deposited on : 2017-06-24
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2747 (2.36-2.32)
Clashscore	180529	2936 (2.36-2.32)
Ramachandran outliers	177936	2912 (2.36-2.32)
Sidechain outliers	177891	2912 (2.36-2.32)
RSRZ outliers	164620	2747 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	 % 80% 15% . .
1	B	549	 2% 75% 15% . 9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoacetylglucosamine mutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	534	Total	C	N	O	P	S	0	0	0
			4043	2541	700	785	1	16			
1	B	499	Total	C	N	O	S		0	1	0
			3753	2362	650	727	14				


There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	LYS	ARG	conflict	UNP A0A0S7E9S6
A	142	ILE	VAL	conflict	UNP A0A0S7E9S6
A	161	PHE	TYR	conflict	UNP A0A0S7E9S6
A	276	ILE	VAL	conflict	UNP A0A0S7E9S6
A	291	LEU	VAL	conflict	UNP A0A0S7E9S6
A	371	MET	LEU	conflict	UNP A0A0S7E9S6
A	467	GLU	ASP	conflict	UNP A0A0S7E9S6
B	35	LYS	ARG	conflict	UNP A0A0S7E9S6
B	142	ILE	VAL	conflict	UNP A0A0S7E9S6
B	161	PHE	TYR	conflict	UNP A0A0S7E9S6
B	276	ILE	VAL	conflict	UNP A0A0S7E9S6
B	291	LEU	VAL	conflict	UNP A0A0S7E9S6
B	371	MET	LEU	conflict	UNP A0A0S7E9S6
B	467	GLU	ASP	conflict	UNP A0A0S7E9S6

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-acetamido-2-deoxy-6-O-phosphono-alpha-D-glucopyranose (three-letter code: 16G) (formula: C₈H₁₆NO₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			19	8	1	9	1		
3	B	1	Total	C	N	O	P	0	0
			19	8	1	9	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	139	Total	O	0	0
			139	139		
5	B	164	Total	O	0	0
			164	164		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphoacetylglucosamine mutase



• Molecule 1: Phosphoacetylglucosamine mutase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.31Å 84.81Å 186.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.25 – 2.34 93.20 – 2.34	Depositor EDS
% Data completeness (in resolution range)	98.9 (93.25-2.34) 99.4 (93.20-2.34)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.14 (at 2.33Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.210 , 0.278 0.213 , 0.277	Depositor DCC
R_{free} test set	486 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å ²)	46.0	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8156	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 16G, MG, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/4098	0.74	2/5554 (0.0%)
1	B	0.53	0/3812	0.74	0/5171
All	All	0.52	0/7910	0.74	2/10725 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	427	ASP	CB-CG-OD1	5.13	122.92	118.30
1	A	480	LEU	CA-CB-CG	5.10	127.04	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	38	LEU	Peptide
1	B	325	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4043	0	3983	41	0
1	B	3753	0	3694	43	0
2	A	1	0	0	0	0
3	A	19	0	14	0	0
3	B	19	0	14	0	0
4	A	6	0	8	0	0
4	B	12	0	16	3	0
5	A	139	0	0	4	0
5	B	164	0	0	4	0
All	All	8156	0	7729	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:ILE:HD12	1:A:248:ILE:H	1.31	0.94
1:A:248:ILE:HG13	1:B:247:VAL:HG11	1.56	0.85
1:B:480:LEU:H	1:B:487:GLN:HE21	1.32	0.76
1:B:284:ASP:OD2	5:B:701:HOH:O	2.08	0.71
1:B:326:ILE:HD11	1:B:396:THR:HB	1.72	0.70
5:A:704:HOH:O	4:B:602:GOL:C1	2.40	0.69
1:A:248:ILE:H	1:A:248:ILE:CD1	1.99	0.69
5:A:704:HOH:O	4:B:602:GOL:H11	1.91	0.68
1:B:304:LEU:HD23	1:B:308:ARG:HG2	1.74	0.68
5:A:769:HOH:O	4:B:602:GOL:H12	1.95	0.67
1:B:286[A]:ASP:OD1	5:B:701:HOH:O	2.14	0.64
1:B:480:LEU:H	1:B:487:GLN:NE2	1.97	0.63
1:A:11:SER:CB	1:A:105:GLU:OE2	2.48	0.62
1:A:39:LEU:HD23	1:A:141:LEU:HD21	1.84	0.60
1:A:359:THR:OG1	1:A:366:LEU:HD23	2.04	0.58
1:A:310:ALA:HB2	1:A:379:PHE:HB2	1.85	0.58
1:A:248:ILE:HD13	5:A:704:HOH:O	2.04	0.57
1:A:445:THR:HG22	1:A:447:LYS:H	1.70	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:128:VAL:HG11	1:B:149:LEU:HD22	1.87	0.56
1:A:346:GLU:HG3	1:A:350:LYS:HD3	1.89	0.55
1:B:467:GLU:HB2	5:B:702:HOH:O	2.06	0.55
1:A:304:LEU:HD23	1:A:308:ARG:HG2	1.89	0.54
1:A:68:ALA:HB2	1:A:76:ASN:HA	1.90	0.54
1:A:480:LEU:H	1:A:487:GLN:HE21	1.55	0.54
1:B:377:VAL:HG22	1:B:387:ILE:HD12	1.89	0.54
1:A:248:ILE:HG13	1:B:247:VAL:CG1	2.33	0.53
1:B:427:ASP:HB3	1:B:430:SER:HB3	1.91	0.53
1:A:335:VAL:HG12	1:A:366:LEU:HD22	1.89	0.53
1:A:10:ILE:HD11	1:A:152:THR:HG22	1.91	0.53
1:B:452:THR:HG23	1:B:453:TYR:CD2	2.44	0.52
1:A:199:PHE:HA	1:A:202:VAL:HG12	1.92	0.51
1:A:480:LEU:H	1:A:487:GLN:NE2	2.08	0.51
1:A:452:THR:HG23	1:A:453:TYR:CD2	2.46	0.51
1:A:43:VAL:HG21	1:A:141:LEU:HG	1.92	0.50
1:A:19:LYS:H	1:A:20:PRO:HA	1.76	0.50
1:A:202:VAL:O	1:A:441:HIS:HE1	1.94	0.50
1:A:490:ILE:HG12	1:A:532:VAL:HG13	1.94	0.50
1:B:129:VAL:HG12	1:B:157:ILE:HB	1.92	0.50
1:A:392:ASN:O	1:A:396:THR:HG23	2.11	0.50
1:B:229:ILE:HA	1:B:232:LEU:HD22	1.94	0.50
1:B:494:GLN:NE2	1:B:502:SER:H	2.10	0.50
1:B:51:THR:HG21	1:B:152:THR:OG1	2.12	0.49
1:B:132:ARG:HD3	1:B:142:ILE:HG12	1.95	0.49
1:B:389:PHE:H	1:B:423:GLN:HE22	1.61	0.48
1:B:494:GLN:HE21	1:B:501:ARG:HA	1.77	0.48
1:A:3:SER:HB3	1:A:6:VAL:HB	1.96	0.47
1:B:317:ILE:HG21	1:B:375:VAL:HG11	1.95	0.47
1:A:371:MET:HA	1:A:388:THR:HG21	1.96	0.47
1:A:466:ALA:HA	1:A:467:GLU:HA	1.69	0.46
1:A:19:LYS:N	1:A:20:PRO:HA	2.31	0.45
1:B:40:ASN:HD22	1:B:144:VAL:HG11	1.80	0.45
1:A:201:ARG:NH2	1:A:416:ALA:HA	2.31	0.45
1:A:316:PHE:CD2	1:A:438:ILE:HG12	2.52	0.45
1:B:466:ALA:HA	1:B:467:GLU:HA	1.71	0.45
1:A:34:MET:HG2	1:A:39:LEU:HD13	1.98	0.45
1:A:494:GLN:NE2	1:A:502:SER:H	2.15	0.45
1:A:461:VAL:HG13	1:A:515:VAL:HB	1.99	0.45
1:B:528:LEU:O	1:B:532:VAL:HG23	2.16	0.45
1:A:53:ARG:CZ	1:A:57:LEU:HD11	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:ILE:HD12	1:A:248:ILE:N	2.14	0.44
1:B:82:ASP:HB2	1:B:86:GLU:HG2	2.00	0.43
1:B:281:ALA:HA	1:B:291:LEU:O	2.18	0.43
1:B:46:VAL:HG13	1:B:80:LEU:HD13	2.01	0.43
1:B:494:GLN:HE21	1:B:502:SER:H	1.66	0.43
1:B:202:VAL:O	1:B:441:HIS:HE1	2.02	0.43
1:B:494:GLN:NE2	1:B:501:ARG:HA	2.34	0.43
1:B:384:HIS:HE1	5:B:737:HOH:O	2.01	0.43
1:B:202:VAL:O	1:B:441:HIS:CE1	2.72	0.43
1:A:494:GLN:HE21	1:A:502:SER:H	1.67	0.42
1:B:323:SER:HB2	1:B:410:SER:OG	2.20	0.42
1:A:306:GLY:HA3	1:A:380:GLU:O	2.19	0.42
1:B:306:GLY:HA3	1:B:380:GLU:O	2.20	0.42
1:B:288:ASP:HB3	1:B:384:HIS:HD2	1.85	0.41
1:B:310:ALA:HB2	1:B:379:PHE:HB2	2.02	0.41
1:B:473:ALA:HB1	1:B:477:GLU:HA	2.02	0.41
1:A:296:ASP:HB3	1:A:298:GLY:H	1.85	0.41
1:A:202:VAL:O	1:A:441:HIS:CE1	2.73	0.41
1:A:296:ASP:OD2	1:A:300:VAL:HB	2.21	0.41
1:B:248:ILE:O	1:B:250:PRO:HD3	2.21	0.41
1:B:389:PHE:H	1:B:423:GLN:NE2	2.18	0.40
1:B:63:GLY:HA2	1:B:129:VAL:O	2.21	0.40
1:B:330:LEU:HD21	1:B:396:THR:OG1	2.21	0.40
1:A:134:THR:HG22	1:B:248:ILE:HG21	2.04	0.40
1:B:225:LEU:HD21	1:B:432:MET:CE	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	523/549 (95%)	505 (97%)	15 (3%)	3 (1%)	22 23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	484/549 (88%)	463 (96%)	19 (4%)	2 (0%)	30	33
All	All	1007/1098 (92%)	968 (96%)	34 (3%)	5 (0%)	25	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	19	LYS
1	A	18	ALA
1	B	120	VAL
1	B	537	ARG
1	A	4	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/442 (94%)	374 (90%)	41 (10%)	6	5
1	B	387/442 (88%)	357 (92%)	30 (8%)	10	10
All	All	802/884 (91%)	731 (91%)	71 (9%)	8	7

All (71) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	49	LEU
1	A	79	LYS
1	A	84	MET
1	A	86	GLU
1	A	112	ASP
1	A	128	VAL
1	A	129	VAL
1	A	141	LEU
1	A	167	LEU
1	A	206	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	215	ASP
1	A	225	LEU
1	A	235	ASP
1	A	238	LEU
1	A	239	ASP
1	A	248	ILE
1	A	265	LYS
1	A	291	LEU
1	A	296	ASP
1	A	302	ARG
1	A	308	ARG
1	A	330	LEU
1	A	350	LYS
1	A	359	THR
1	A	361	THR
1	A	364	LYS
1	A	372	ARG
1	A	396	THR
1	A	403	GLN
1	A	434	LEU
1	A	450	LEU
1	A	461	VAL
1	A	467	GLU
1	A	472	LYS
1	A	480	LEU
1	A	486	LEU
1	A	489	LYS
1	A	499	LYS
1	A	513	VAL
1	A	522	ARG
1	B	11	SER
1	B	16	GLN
1	B	41	THR
1	B	49	LEU
1	B	56	LYS
1	B	97	THR
1	B	111	TYR
1	B	112	ASP
1	B	114	LEU
1	B	128	VAL
1	B	129	VAL
1	B	204	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	225	LEU
1	B	232	LEU
1	B	235	ASP
1	B	266	GLN
1	B	289	ARG
1	B	291	LEU
1	B	302	ARG
1	B	308	ARG
1	B	320	LEU
1	B	322	ARG
1	B	364	LYS
1	B	414	LEU
1	B	434	LEU
1	B	450	LEU
1	B	461	VAL
1	B	486	LEU
1	B	513	VAL
1	B	522	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	ASN
1	A	336	GLN
1	A	367	HIS
1	A	487	GLN
1	A	494	GLN
1	B	40	ASN
1	B	76	ASN
1	B	180	GLN
1	B	189	GLN
1	B	336	GLN
1	B	384	HIS
1	B	407	GLN
1	B	423	GLN
1	B	487	GLN
1	B	494	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	69	2,1	8,9,10	0.79	0	8,12,14	2.72	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	69	2,1	-	4/5/8/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	SEP	OG-CB-CA	6.89	114.85	108.14
1	A	69	SEP	P-OG-CB	2.16	124.25	118.30

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	69	SEP	CB-OG-P-O1P
1	A	69	SEP	CB-OG-P-O2P
1	A	69	SEP	CB-OG-P-O3P
1	A	69	SEP	N-CA-CB-OG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	16G	B	601	-	19,19,19	1.09	2 (10%)	28,28,28	1.08	2 (7%)
4	GOL	A	603	-	5,5,5	0.31	0	5,5,5	0.56	0
4	GOL	B	603	-	5,5,5	0.44	0	5,5,5	0.41	0
4	GOL	B	602	-	5,5,5	0.42	0	5,5,5	0.50	0
3	16G	A	602	-	19,19,19	1.26	3 (15%)	28,28,28	1.39	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	16G	B	601	-	-	2/10/30/30	0/1/1/1
4	GOL	A	603	-	-	4/4/4/4	-
4	GOL	B	603	-	-	2/4/4/4	-
4	GOL	B	602	-	-	0/4/4/4	-
3	16G	A	602	-	-	0/10/30/30	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	16G	P-O2P	2.34	1.63	1.54
3	B	601	16G	C1-C2	2.33	1.55	1.52
3	A	602	16G	P-O1P	2.32	1.58	1.50
3	A	602	16G	C1-C2	2.06	1.55	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	16G	P-O2P	2.02	1.62	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	16G	O5-C1-C2	4.40	113.94	109.52
3	A	602	16G	O3P-P-O6	3.13	115.05	106.73
3	B	601	16G	O6-P-O1P	2.88	114.56	106.47
3	B	601	16G	C1-C2-C3	2.79	114.35	110.54
3	A	602	16G	C6-C5-C4	-2.13	107.64	112.09

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	601	16G	C6-O6-P-O1P
3	B	601	16G	C6-O6-P-O3P
4	A	603	GOL	O1-C1-C2-C3
4	B	603	GOL	C1-C2-C3-O3
4	A	603	GOL	O1-C1-C2-O2
4	A	603	GOL	C1-C2-C3-O3
4	A	603	GOL	O2-C2-C3-O3
4	B	603	GOL	O2-C2-C3-O3


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	602	GOL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	533/549 (97%)	-0.18	4 (0%) 82 85	32, 51, 84, 103	0
1	B	499/549 (90%)	-0.19	11 (2%) 62 67	23, 47, 81, 113	1 (0%)
All	All	1032/1098 (93%)	-0.19	15 (1%) 71 75	23, 49, 83, 113	1 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	110	VAL	4.3
1	A	105	GLU	4.0
1	B	120	VAL	3.6
1	B	119	ASP	3.4
1	B	122	MET	3.0
1	B	467	GLU	2.7
1	B	124	ASN	2.5
1	A	110	VAL	2.4
1	B	111	TYR	2.3
1	B	5	ALA	2.2
1	B	121	SER	2.2
1	B	151	ALA	2.1
1	A	109	ASP	2.1
1	B	125	PRO	2.0
1	A	104	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	SEP	A	69	10/11	0.77	0.14	54,67,87,92	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	B	603	6/6	0.85	0.12	46,55,56,61	0
4	GOL	B	602	6/6	0.91	0.10	59,62,68,75	0
2	MG	A	601	1/1	0.92	0.06	49,49,49,49	0
4	GOL	A	603	6/6	0.95	0.06	44,49,52,56	0
3	16G	A	602	19/19	0.95	0.06	42,46,50,51	0
3	16G	B	601	19/19	0.95	0.07	58,65,75,76	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 16G A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 16G B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.