



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 01:04 PM EDT

PDB ID : 4OC5
Title : X-ray structure of of human glutamate carboxypeptidase II (GCPII) in a complex with CHIBzL, a urea-based inhibitor N 2 -{[(S)-carboxy(4-hydroxyphenyl)methyl]carbamoyl}-N 6 -(4-iodobenzoyl)-L-lysine
Authors : Pavlicek, J.; Ptacek, J.; Cerny, J.; Byun, Y.; Skultetyova, L.; Pomper, M.; Lubkowski, J.; Barinka, C.
Deposited on : 2014-01-08
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	709	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>8%</div> <div>..</div> </div> </div>
2	B	2	<div> <div>50%</div> <div>50%</div> </div>
2	C	2	<div> <div>100%</div> </div>
2	D	2	<div> <div>100%</div> </div>
3	E	4	<div> <div>100%</div> </div>

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 6466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	44	0
			5754	3700	962	1069	23			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ARG	-	expression tag	UNP Q04609
A	43	SER	-	expression tag	UNP Q04609

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		


- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0

- Molecule 8 is N 2 -{[(S)-carboxy(4-hydroxyphenyl)methyl]carbamoyl}-N 6 -(4-iodobenzoyl)-L-lysine (three-letter code: 2QM) (formula: C₂₂H₂₄IN₃O₇).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C I N O 33 22 1 3 7	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	499	Total O 499 499	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate carboxypeptidase 2




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:

100%

MAG1
MAG2
BMA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.58Å 130.45Å 159.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.70 20.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.7 (20.00-1.70) 97.6 (20.00-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.47 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.169 , 0.204 0.169 , 0.203	Depositor DCC
R_{free} test set	1139 reflections (1.01%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6466	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN, NAG, MAN, BMA, CL, 2QM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	2/6022 (0.0%)	0.86	9/8154 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	434	TRP	CE3-CZ3	5.36	1.47	1.38
1	A	164	PHE	CE2-CZ	5.26	1.47	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	688[A]	ARG	NE-CZ-NH2	-6.63	116.99	120.30
1	A	688[B]	ARG	NE-CZ-NH2	-6.63	116.99	120.30
1	A	673	ARG	NE-CZ-NH1	6.26	123.43	120.30
1	A	673	ARG	NE-CZ-NH2	-6.15	117.22	120.30
1	A	174[A]	LEU	CA-CB-CG	5.75	128.52	115.30
1	A	174[B]	LEU	CA-CB-CG	5.75	128.52	115.30
1	A	664	MET	CG-SD-CE	-5.25	91.79	100.20
1	A	370	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	A	440	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5754	0	5633	45	0
2	B	28	0	25	1	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
3	E	50	0	43	0	0
4	A	42	0	39	0	0
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	33	0	21	1	0
9	A	499	0	0	8	2
All	All	6466	0	5811	46	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:660[A]:VAL:O	1:A:664:MET:HG2	1.38	1.22
1:A:147:PRO:HG2	1:A:152:GLU:HG2	1.46	0.97
1:A:89[B]:ASN:HD21	1:A:378:ARG:HH21	1.08	0.97
1:A:89[B]:ASN:ND2	1:A:378:ARG:HH21	1.79	0.78
1:A:90:PHE:CE2	1:A:94:LYS:HE2	2.20	0.76
1:A:733:TYR:HE2	9:A:1244:HOH:O	1.70	0.74
1:A:58[A]:MET:CE	1:A:586:GLU:HG2	2.18	0.72
1:A:131:ILE:HG22	1:A:137:GLU:HG2	1.72	0.72
1:A:641:GLU:HG3	9:A:1253:HOH:O	1.91	0.69
1:A:58[A]:MET:HE1	1:A:586:GLU:HG2	1.74	0.68
1:A:89[B]:ASN:HD21	1:A:378:ARG:NH2	1.87	0.66
1:A:400:ARG:O	1:A:404[B]:THR:HG23	1.96	0.65
1:A:539:LYS:HD2	1:A:541:TRP:HB3	1.80	0.63
1:A:278:ALA:HB3	1:A:280[A]:ARG:NH1	2.15	0.62
1:A:308:LYS:HB2	9:A:1397:HOH:O	2.01	0.60
1:A:58[A]:MET:HE2	1:A:586:GLU:HG2	1.86	0.57
1:A:177:VAL:HG12	1:A:188[A]:LEU:HD11	1.89	0.55
1:A:188[B]:LEU:HD11	1:A:329:VAL:HG11	1.89	0.54
1:A:465:ASP:OD1	1:A:513:SER:HB2	2.08	0.53
1:A:579:VAL:O	1:A:583[B]:MET:HG2	2.08	0.53
1:A:155:SER:O	1:A:156:ASP:HB2	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58[A]:MET:HE1	1:A:586:GLU:CG	2.45	0.45
1:A:177:VAL:CG2	1:A:201[B]:VAL:HG13	2.46	0.45
1:A:177:VAL:HG22	1:A:201[B]:VAL:HG13	1.98	0.45
1:A:444:GLU:HA	1:A:444:GLU:OE1	2.17	0.44
1:A:249:PRO:HG2	1:A:556:TYR:CE2	2.53	0.43
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.53	0.43
1:A:610:LYS:HE2	9:A:1080:HOH:O	2.18	0.43
1:A:131:ILE:CG2	1:A:137:GLU:HG2	2.43	0.43
1:A:517:SER:HB2	1:A:694:PRO:HG3	2.00	0.43
1:A:505:GLU:HG3	1:A:506:PHE:CE1	2.54	0.42
1:A:90:PHE:CZ	1:A:94:LYS:HE2	2.54	0.42
1:A:236:ALA:O	1:A:239[A]:VAL:HG13	2.19	0.42
1:A:539:LYS:CD	1:A:541:TRP:HB3	2.49	0.41
1:A:534:ARG:HG3	8:A:818:2QM:IAH	2.91	0.41
1:A:733:TYR:CE2	9:A:1244:HOH:O	2.55	0.41
1:A:526[A]:GLN:NE2	9:A:1374:HOH:O	2.37	0.41
9:A:1144:HOH:O	2:B:2:NAG:H81	2.21	0.41
1:A:320[A]:ARG:HG2	1:A:327:TYR:HB2	2.02	0.41
1:A:653:PHE:HE1	1:A:661[A]:LEU:HB2	1.86	0.40
1:A:688[B]:ARG:NH1	9:A:970:HOH:O	2.32	0.40
1:A:517:SER:OG	1:A:518:GLY:N	2.52	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:1115:HOH:O	9:A:1364:HOH:O[2_565]	1.97	0.23
9:A:1255:HOH:O	9:A:1399:HOH:O[2_565]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	731/709 (103%)	714 (98%)	16 (2%)	1 (0%)	48 32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	382	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	633/605 (105%)	618 (98%)	15 (2%)	44 27

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LYS
1	A	68[A]	GLU
1	A	68[B]	GLU
1	A	152	GLU
1	A	155	SER
1	A	171	GLU
1	A	239[A]	VAL
1	A	239[B]	VAL
1	A	505	GLU
1	A	519	ASN
1	A	537	TYR
1	A	600	TYR
1	A	662[A]	ARG
1	A	662[B]	ARG
1	A	673	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	HIS
1	A	124	HIS
1	A	347	HIS
1	A	443	GLN
1	A	618	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.51	0	17,19,21	1.06	1 (5%)
2	NAG	B	2	2	14,14,15	0.74	0	17,19,21	1.20	2 (11%)
2	NAG	C	1	1,2	14,14,15	0.82	1 (7%)	17,19,21	0.86	1 (5%)
2	NAG	C	2	2	14,14,15	0.43	0	17,19,21	1.16	1 (5%)
2	NAG	D	1	1,2	14,14,15	0.96	1 (7%)	17,19,21	1.29	3 (17%)
2	NAG	D	2	2	14,14,15	0.53	0	17,19,21	1.21	3 (17%)
3	NAG	E	1	1,3	14,14,15	0.52	0	17,19,21	1.27	2 (11%)
3	NAG	E	2	3	14,14,15	0.71	0	17,19,21	1.51	2 (11%)
3	BMA	E	3	3	11,11,12	0.47	0	15,15,17	0.86	1 (6%)
3	MAN	E	4	3	11,11,12	0.55	0	15,15,17	0.99	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	NAG	O7-C7	2.54	1.28	1.23
2	C	1	NAG	C1-C2	2.03	1.55	1.52

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	1	NAG	O5-C1-C2	-3.23	106.30	111.29
3	E	2	NAG	C3-C4-C5	-2.91	104.96	110.23
2	D	1	NAG	O4-C4-C3	-2.76	103.87	110.38
2	B	2	NAG	C1-C2-N2	2.75	114.77	110.43
2	D	2	NAG	C8-C7-N2	2.59	120.41	116.12
2	B	2	NAG	C2-N2-C7	2.53	126.30	122.90
2	D	1	NAG	O5-C5-C4	-2.43	104.92	110.83
2	C	2	NAG	O5-C5-C6	2.43	112.38	107.66
2	D	2	NAG	O7-C7-C8	-2.41	117.77	122.05
3	E	2	NAG	C8-C7-N2	2.39	120.08	116.12
2	B	1	NAG	O3-C3-C4	-2.26	105.05	110.38
2	D	2	NAG	C1-O5-C5	2.26	115.21	112.19
2	C	1	NAG	O5-C1-C2	-2.20	107.89	111.29
3	E	1	NAG	C6-C5-C4	-2.16	107.72	113.02
3	E	4	MAN	O5-C5-C6	2.14	111.82	107.66
3	E	3	BMA	O3-C3-C2	-2.10	105.76	110.05
2	D	1	NAG	O3-C3-C2	-2.01	105.22	109.40

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	806	1	14,14,15	0.50	0	17,19,21	1.14	2 (11%)
4	NAG	A	803	1	14,14,15	0.68	0	17,19,21	1.54	2 (11%)
4	NAG	A	807	1	14,14,15	0.67	0	17,19,21	1.70	6 (35%)
8	2QM	A	818	5	34,34,34	1.41	6 (17%)	43,45,45	1.83	9 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	806	1	-	0/6/23/26	0/1/1/1
4	NAG	A	803	1	-	2/6/23/26	0/1/1/1
4	NAG	A	807	1	-	0/6/23/26	0/1/1/1
8	2QM	A	818	5	-	3/32/32/32	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	818	2QM	CBC-IAH	3.34	2.18	2.10
8	A	818	2QM	CBD-CBA	3.23	1.57	1.50
8	A	818	2QM	CAN-CAL	2.80	1.43	1.38
8	A	818	2QM	CAK-CAM	2.54	1.42	1.38
8	A	818	2QM	OXT-C	2.32	1.29	1.22
8	A	818	2QM	OAF-CAY	2.03	1.28	1.22

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	818	2QM	CAL-CBC-CAK	-6.24	112.39	120.64
4	A	803	NAG	C1-C2-N2	-4.10	103.97	110.43
8	A	818	2QM	CAM-CAK-CBC	3.83	124.46	119.55
8	A	818	2QM	CAN-CAL-CBC	3.29	123.77	119.55
4	A	807	NAG	O5-C1-C2	-3.22	106.31	111.29
8	A	818	2QM	CAK-CBC-IAH	3.10	124.32	119.69
8	A	818	2QM	CAO-CBE-CBG	-3.05	115.90	120.78
4	A	807	NAG	O5-C5-C6	2.91	113.33	107.66
4	A	803	NAG	O3-C3-C2	-2.77	103.64	109.40
8	A	818	2QM	CAN-CBD-CAM	-2.74	115.08	118.57
4	A	806	NAG	C1-O5-C5	2.69	115.80	112.19
8	A	818	2QM	CAP-CAJ-CBB	-2.67	117.06	119.88
4	A	807	NAG	C1-O5-C5	2.56	115.62	112.19
4	A	807	NAG	O7-C7-C8	-2.48	117.63	122.05
8	A	818	2QM	CAL-CBC-IAH	2.40	123.28	119.69
4	A	807	NAG	C1-C2-N2	2.14	113.80	110.43
4	A	806	NAG	O5-C5-C6	2.07	111.69	107.66
4	A	807	NAG	C3-C4-C5	-2.06	106.49	110.23
8	A	818	2QM	CAI-CAO-CBE	-2.04	119.15	121.18

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	803	NAG	C8-C7-N2-C2
4	A	803	NAG	O7-C7-N2-C2
8	A	818	2QM	CAY-CBG-NAW-CAZ
8	A	818	2QM	CAO-CBE-CBG-CAY
8	A	818	2QM	CAP-CBE-CBG-CAY

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	818	2QM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/709 (97%)	-0.13	17 (2%) 59 62	9, 28, 48, 65	42 (6%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	GLU	5.6
1	A	656[A]	SER	3.9
1	A	541	TRP	3.8
1	A	652	ASP	3.2
1	A	653	PHE	3.0
1	A	506	PHE	2.9
1	A	155	SER	2.8
1	A	543	THR	2.7
1	A	154	VAL	2.5
1	A	544	ASN	2.4
1	A	505	GLU	2.3
1	A	719	VAL	2.2
1	A	135	GLY	2.2
1	A	133	GLU	2.1
1	A	191	ASP	2.0
1	A	189	GLU	2.0
1	A	507	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	2	14/15	0.77	0.12	45,53,57,58	0
2	NAG	C	2	14/15	0.80	0.12	60,62,67,69	0
3	MAN	E	4	11/12	0.81	0.12	50,52,56,57	0
3	NAG	E	2	14/15	0.82	0.13	39,43,52,52	0
2	NAG	D	2	14/15	0.85	0.11	36,43,49,51	0
3	BMA	E	3	11/12	0.88	0.09	42,44,45,47	0
2	NAG	C	1	14/15	0.88	0.10	40,44,48,54	0
2	NAG	D	1	14/15	0.92	0.08	31,34,42,44	0
2	NAG	B	1	14/15	0.93	0.09	35,43,48,52	0
3	NAG	E	1	14/15	0.94	0.09	25,29,39,45	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	A	803	14/15	0.77	0.13	46,53,56,58	0
4	NAG	A	806	14/15	0.84	0.11	60,68,70,71	0
4	NAG	A	807	14/15	0.84	0.12	33,49,55,59	0
7	CL	A	817	1/1	0.97	0.04	29,29,29,29	0
8	2QM	A	818	33/33	0.98	0.09	22,29,38,45	0
6	CA	A	816	1/1	1.00	0.01	18,18,18,18	0
5	ZN	A	814	1/1	1.00	0.02	23,23,23,23	0
5	ZN	A	815	1/1	1.00	0.02	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.