



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 05:50 am BST

PDB ID : 5OD9
Title : Structure of the engineered metalloesterase MID1sc9
Authors : Studer, S.; Mittl, P.R.E.; Hilvert, D.
Deposited on : 2017-07-05
Resolution : 1.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1168 (1.14-1.10)
Clashscore	141614	1205 (1.14-1.10)
Ramachandran outliers	138981	1168 (1.14-1.10)
Sidechain outliers	138945	1165 (1.14-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	97	 74% 20% . .
1	B	97	 74% 13% .. 9%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 1860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MID1sc9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	95	Total	C	N	O	S	0	9	0
			823	510	156	153	4			
1	B	88	Total	C	N	O	S	0	7	0
			781	488	145	144	4			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Zn	0	0
			4	4		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	N	0	0
			5	3	2		

- Molecule 7 is (2 {S})-2-phenylpropanoic acid (three-letter code: 9RW) (formula: $C_9H_{10}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			11	9	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	126	Total O 126 126	0	0
8	B	97	Total O 97 97	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MID1sc9

Chain A:  74% 20% . .



- Molecule 1: MID1sc9

Chain B:  74% 13% . . 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	43.82Å 104.96Å 33.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.00 – 1.13 40.44 – 1.13	Depositor EDS
% Data completeness (in resolution range)	99.0 (52.00-1.13) 99.4 (40.44-1.13)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 1.13Å)	Xtriage
Refinement program	SHELXL	Depositor
R, R_{free}	(Not available) , (Not available) 0.152 , 0.191	Depositor DCC
R_{free} test set	2928 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	10.9	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 88.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	1860	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 9RW, CL, ZN, IMD, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/855	1.43	14/1148 (1.2%)
1	B	0.75	0/797	1.64	13/1073 (1.2%)
All	All	0.77	0/1652	1.53	27/2221 (1.2%)

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2077[A]	ARG	NE-CZ-NH2	-17.31	111.65	120.30
1	B	2077[B]	ARG	NE-CZ-NH2	-17.31	111.65	120.30
1	A	1072[A]	ARG	NE-CZ-NH2	-11.14	114.73	120.30
1	A	1072[C]	ARG	NE-CZ-NH2	-11.14	114.73	120.30
1	A	1074	ASP	CB-CG-OD1	9.17	126.56	118.30
1	B	2028	ARG	NE-CZ-NH1	7.96	124.28	120.30
1	B	2014	PHE	CB-CG-CD1	7.35	125.94	120.80
1	B	2077[A]	ARG	NE-CZ-NH1	6.92	123.76	120.30
1	B	2077[B]	ARG	NE-CZ-NH1	6.92	123.76	120.30
1	A	1074	ASP	CB-CG-OD2	-6.90	112.09	118.30
1	B	2068	ARG	NE-CZ-NH2	-6.66	116.97	120.30
1	A	1078	ARG	NE-CZ-NH2	-6.62	116.99	120.30
1	B	2056	GLN	CA-CB-CG	6.50	127.69	113.40
1	A	1095	ASP	CB-CG-OD1	6.48	124.13	118.30
1	A	1091	PHE	CB-CG-CD1	-6.19	116.47	120.80
1	B	2023	ARG	NE-CZ-NH2	6.04	123.32	120.30
1	B	2046	GLU	CA-C-O	5.93	132.56	120.10
1	A	1017	GLN	CG-CD-OE1	5.77	133.15	121.60
1	B	2059	TYR	CB-CG-CD1	5.71	124.43	121.00
1	A	1025	ASP	CB-CG-OD2	-5.51	113.34	118.30
1	A	1017	GLN	OE1-CD-NE2	-5.43	109.41	121.90
1	B	2028	ARG	NH1-CZ-NH2	-5.35	113.51	119.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2078	ARG	NE-CZ-NH2	5.18	122.89	120.30
1	A	1044	GLN	CA-CB-CG	5.14	124.70	113.40
1	A	1072[A]	ARG	NH1-CZ-NH2	5.12	125.03	119.40
1	A	1072[C]	ARG	NH1-CZ-NH2	5.12	125.03	119.40
1	A	1014	PHE	CB-CG-CD2	5.03	124.32	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	823	0	814	18	0
1	B	781	0	762	13	0
2	A	4	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	2	0	0	0	0
5	A	8	0	12	1	0
6	A	5	0	5	1	0
7	A	11	0	0	2	0
8	A	126	0	0	8	0
8	B	97	0	0	9	0
All	All	1860	0	1593	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1056:GLN:HB2	8:A:3161:HOH:O	1.90	0.72
1:B:2077[A]:ARG:O	1:B:2081:GLU:HG3	1.90	0.71
1:A:1073:MET:O	1:A:1077[B]:ARG:HG3	2.03	0.59
1:A:1078:ARG:HH12	7:A:3010:9RW:CD	2.16	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2085[A]:GLN:HG2	8:B:2207:HOH:O	2.03	0.58
1:A:1082:ASN:HD22	7:A:3010:9RW:C3	2.17	0.57
1:B:2043:GLN:HG3	8:B:2216:HOH:O	2.04	0.57
1:B:2077[B]:ARG:O	1:B:2081:GLU:HG3	2.04	0.57
1:B:2093:GLN:HB3	8:B:2275:HOH:O	2.07	0.55
1:A:1043:GLN:OE1	1:B:2068:ARG:HD3	2.08	0.54
1:A:1092:GLN:O	1:A:1095:ASP:HB2	2.10	0.52
1:A:1066:GLN:HG3	8:B:2222:HOH:O	2.12	0.50
1:A:1072[A]:ARG:HG2	8:A:3122:HOH:O	2.10	0.49
5:A:3007:EDO:H22	8:A:3178:HOH:O	2.12	0.49
1:A:1066:GLN:HG3	8:A:3191:HOH:O	2.12	0.48
1:B:2077[B]:ARG:NH2	8:B:2208:HOH:O	2.47	0.48
1:A:1088:LYS:NZ	8:A:3109:HOH:O	2.47	0.48
1:B:2093:GLN:NE2	8:B:2209:HOH:O	2.47	0.47
1:B:2042:PHE:CE1	1:B:2046:GLU:HB2	2.50	0.46
1:A:1072[C]:ARG:NH1	1:A:1075:GLU:HG3	2.30	0.46
1:B:2073:MET:O	1:B:2077[B]:ARG:HG3	2.15	0.46
1:A:1066:GLN:NE2	8:A:3112:HOH:O	2.49	0.45
1:B:2043:GLN:HA	8:B:2202:HOH:O	2.16	0.45
1:A:1031:GLN:OE1	1:A:1068[B]:ARG:NH2	2.50	0.44
1:A:1017:GLN:NE2	8:A:3111:HOH:O	2.49	0.44
1:A:1077[A]:ARG:NH1	8:A:3114:HOH:O	2.51	0.43
1:B:2042:PHE:CZ	1:B:2046:GLU:HG3	2.54	0.43
1:A:1031:GLN:OE1	1:A:1068[B]:ARG:NH1	2.52	0.42
6:A:3009:IMD:H2	8:B:2243:HOH:O	2.19	0.42
1:A:1028[A]:ARG:NE	1:A:1068[A]:ARG:HD3	2.35	0.42
1:A:1004:LEU:HD22	1:A:1037[A]:LEU:CD1	2.50	0.41
1:B:2088:LYS:HE2	8:B:2213:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/97 (105%)	101 (99%)	1 (1%)	0	100	100
1	B	91/97 (94%)	90 (99%)	1 (1%)	0	100	100
All	All	193/194 (100%)	191 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/80 (110%)	88 (100%)	0	100	100
1	B	83/80 (104%)	82 (99%)	1 (1%)	71	35
All	All	171/160 (107%)	170 (99%)	1 (1%)	86	61

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2046	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1066	GLN
1	A	1082	ASN
1	B	2006	GLN
1	B	2033	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 9 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	9RW	A	3010	-	11,11,11	2.42	1 (9%)	14,14,14	1.64	3 (21%)
5	EDO	A	3008	-	3,3,3	0.43	0	2,2,2	0.18	0
5	EDO	A	3007	-	3,3,3	0.59	0	2,2,2	0.65	0
6	IMD	A	3009	-	3,5,5	0.72	0	4,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	9RW	A	3010	-	-	2/8/8/8	0/1/1/1
5	EDO	A	3008	-	-	0/1/1/1	-
5	EDO	A	3007	-	-	0/1/1/1	-
6	IMD	A	3009	-	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	3010	9RW	CB-CG	-7.44	1.40	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
7	A	3010	9RW	CB-CG-CD	3.55	118.94	108.60
7	A	3010	9RW	C8-C9-CB	2.53	123.78	120.65
7	A	3010	9RW	OE2-CD-OE1	-2.40	118.63	124.09

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	3010	9RW	C9-CB-CG-CD
7	A	3010	9RW	CA-CB-CG-CD

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	3010	9RW	2	0
5	A	3007	EDO	1	0
6	A	3009	IMD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.